

378

**CLINICAL AND IMMUNOLOGICAL ACTIVITY IN POLISH COHORT OF SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS TREATED WITH GLUCOCORTICOIDS**

K Pawlak-Bus\*, P Leszczynski. *Poznan University of Medical Sciences, Rheumatology and Rehabilitation, Poznan, Poland*

10.1136/lupus-2017-000215.378

**Background and aims** Nowadays the lupus treatment strategy is based on background therapy, immunosuppressive drugs and glucocorticoids (GC). Using minimal effective dose of GC only in flares is a recommendation for preventing complications which increase mortality. The Aim of the study was to evaluate SLE clinical and immunological activity in lupus patients during the standard clinical care and analyse GC treatment.

**Methods** We observed Polish cohort of patients with SLE recognised and confirmed by SLICC classification criteria 2012. 127 patients (118 female and 9 male) with average age 43±6 years (range 18–64 years), average disease duration 7.8±5.6 years (range 1.0–15.0 years). All of them were treated with oral and pulse GC and standard immunosuppressive therapies (CTX, MMF, AZT, MTX, CsA). As a background therapy 77% of these patients were on chloroquine or hydroxychloroquine (CQ/HCQ) Table 1. All patients were assessed according to SLEDAI (Gladman *et al*, 2002) and divided into 5 groups: no GC, low dose, medium dose, high dose and puls GC therapy group. Immunological activity was assessed by anti-dsDNA and C3 and C4 complements levels.

**Results:** Results Tables 2 and 3.

**Conclusions** In this Polish cohort lupus patients GC doses depended on lupus activity. Minimizing glucocorticoid exposure is an important part of appropriate management of lupus patients. Proper assessment of clinical and immunological lupus activity is critically for treatment decisions, especially for long-term GC use.

**Abstract 378 Table 1** Baseline characteristic.

Gender	Female 93% (n=118)	Male 7% (n=9)			
SLEDAI score	<b>Very high &gt; 20</b> 35% (44)	<b>High 13-20</b> 16% (20)	<b>Moderate 6-12</b> 36% (46)	<b>Low &lt; 6</b> 13% (17)	
Serology	<b>+dsDNA;</b> <b>↓C3/C4</b> 41% (n=52)	<b>+dsDNA;</b> <b>N C3/C4</b> 14% (n=18)	<b>-dsDNA;</b> <b>↓C3/C4</b> 10% (n=13)	<b>-dsDNA;</b> <b>N C3/C4</b> 35% (n=44)	
Treatment	<b>CTX</b> 13% (n=17)	<b>MMF</b> 21% (n=27)	<b>AZT</b> 23% (n=29)	<b>CsA/MTX</b> 16% (n=20)	<b>CQ/HCQ</b> 77% (n=98)
Glucocorticoids	<b>High dose</b> 21% (n=27)	<b>Medium dose</b> 16% (n=20)	<b>Low dose</b> 41% (n=52)	<b>No steroids</b> 22% (n=28)	

+dsDNA - antibodies anti-dsDNA are present; -dsDNA - antibodies anti-dsDNA are not present; ↓C3/C4 - C3 or C4 complement level is below normal range; N C3/C4 - C3 or C4 complement is in normal range; CTX - cyclophosphamide; MMF - mycophenolate mofetil; AZT - azathioprine; CsA - cyclosporin A; MTX - methotrexate; CQ - chloroquine; HCQ - hydroxychloroquine

**Abstract 378 Table 2** Clinical lupus activity and glucocorticoids (GC) doses\*.

Glucocorticoids doses	Patients	Mean SLEDAI score ± SD	SLE activity by SLEDAI scores	
Group 1 no GC	22% (n=28)	7 ± 3	SLEDAI <6	21% (n=6)
			SLEDAI 6-12	68% (n=19)
			SLEDAI 13-20	11% (n=3)
			SLEDAI >20	(n=0)
Group 2 < 7,5mg.	41% (n=52)	13 ± 7	SLEDAI <6	17% (n=9)
			SLEDAI 6-12	31% (n=16)
			SLEDAI 13-20	23% (n=12)
			SLEDAI >20	29% (n=15)
Group 3 7,5-10 mg.	16% (n=20)	19 ± 15	SLEDAI <6	15% (n=3)
			SLEDAI 6-12	25% (n=5)
			SLEDAI 13-20	20% (n=4)
			SLEDAI >20	40% (n=8)
Group 4 >10 mg.	4% (n=5)	30 ± 13	SLEDAI <6	(n=0)
			SLEDAI 6-12	(n=0)
Group 5 puls therapy iv.	17% (n=22)	21% (n=27)	SLEDAI 13-20	15% (n=4)
			SLEDAI >20	85% (n=23)

\*GC dose of prednison

**Abstract 378 Table 3** Immunological state and GC doses.

GC dose	Immunological SLE activity			
	+dsDNA; ↓C3/C4	+dsDNA; N C3/C4	-dsDNA; ↓C3/C4	-dsDNA; N C3/C4
Group 1	18% (n=5)	14% (n=4)	18% (n=5)	50% (n=14)
Group 2	29% (n=15)	17% (n=9)	8% (n=4)	46% (n=24)
Group 3	55% (n=11)	15% (n=3)	10% (n=2)	20% (n=4)
Group 4	100% (n=5)	n=0	n=0	n=0
Group 5	64% (n=14)	13% (n=3)	5% (n=1)	18% (n=4)

+dsDNA - antibodies anti-dsDNA are present; -dsDNA - antibodies anti-dsDNA are not present; ↓C3/C4 - C3 or C4 complement level is low; N C3/C4 - C3 or C4 complement is in normal range; Groups 1 - no GC; 2 - low dose of GC; 3 - medium dose of GC; 4 - high dose of GC; 5 - GC puls therapy