

Association of the interferon signature metric with serological disease manifestations but not global activity scores in multiple cohorts of patients with SLE

William P Kennedy,¹ Romeo Maciuca,² Kristen Wolslegel,³ Wei Tew,³ Alexander R Abbas,⁴ Christina Chaivorapol,⁴ Alyssa Morimoto,⁵ Jacqueline M McBride,⁶ Paul Brunetta,⁷ Bruce C Richardson,⁸ John C Davis Jr,¹ Timothy W Behrens,⁹ Michael J Townsend³

ABSTRACT

To cite: Kennedy WP, Maciuca R, Wolslegel K, *et al.* Association of the interferon signature metric with serological disease manifestations but not global activity scores in multiple cohorts of patients with SLE. *Lupus Science & Medicine* 2015;**2**:e000080. doi:10.1136/ lupus-2014-000080

► Additional material is available. To view please visit the journal (http://dx.doi.org/ 10.1136/lupus-2014-000080).

Received 23 December 2014 Revised 3 February 2015 Accepted 11 March 2015



► http://dx.doi.org/10.1136/ annrheumdis-2014-206090



For numbered affiliations see end of article.

Correspondence to Dr Michael J Townsend; townsend.michael@gene.com **Objectives:** The interferon (IFN) signature (IS) in patients with systemic lupus erythematosus (SLE) includes over 100 genes induced by type I IFN pathway activation. We developed a method to quantify the IS using three genes—the IS metric (ISM)—and characterised the clinical characteristics of patients with SLE with different ISM status from multiple clinical trials.

Methods: Blood microarray expression data from a training cohort of patients with SLE confirmed the presence of the IS and identified surrogate genes. We assayed these genes in a quantitative PCR (qPCR) assay, yielding an ISM from the IS. The association of ISM status with clinical disease characteristics was assessed in patients with extrarenal lupus and lupus nephritis from four clinical trials.

Results: Three genes, HERC5, EPSTI and CMPK2, correlated well with the IS (p>0.96), and composed the ISM qPCR assay. Using the 95th centile for healthy control data, patients with SLE from different studies were classified into two ISM subsets—ISM-Low and ISM-High—that are longitudinally stable over 36 weeks. Significant associations were identified between ISM-High status and higher titres of anti-dsDNA antibodies, presence of anti extractable nuclear antigen autoantibodies, elevated serum B cell activating factor of the tumour necrosis factor family (BAFF) levels, and hypocomplementaemia. However, measures of overall clinical disease activity were similar for ISM-High and ISM-Low groups.

Conclusions: The ISM is an IS biomarker that divides patients with SLE into two subpopulations—ISM-High and ISM-Low—with differing serological manifestations. The ISM does not distinguish between high and low disease activity, but may have utility in identifying patients more likely to respond to treatment(s) targeting IFN- α .

Clinicaltrials.gov registration number: NCT00962832.

KEY MESSAGES

- A three-gene surrogate (interferon signature metric, ISM) for the interferon signature in patients with systemic lupus erythematosus (SLE) has a distinct bimodal profile that is similar across different interventional clinical SLE studies.
- The ISM profile distinguishes patients with SLE on serological characteristics but not on clinical phenotype.
- Patients with an ISM-High status have elevated anti-dsDNA and ENA autoantibodies, hypocomplementaemia, and elevated serum BAFF levels.

INTRODUCTION

Systemic lupus erythematosus (SLE) is a chronic autoimmune disease of unknown aetiology. It is characterised by the presence of pathogenic autoantibodies to nuclear antigens, elevated immune complexes (ICs), and immunological abnormalities, including lymphopenia, elevated immunoglobulin and autoantibody levels, complement activation, and abnormal activation of the innate and adaptive immune system, and often leads to multisystem organ damage.^{1–3}

Development of novel therapies to manage SLE has been hampered by several challenges, including poorly understood pathogenesis, the heterogeneity of disease activity across and within patient populations, and difficulties conducting interventional studies.^{4 5} One approach for development of successful therapies may follow from efforts to identify useful biomarkers that can classify patients with SLE into more homogenous subsets for use in clinical trials and clinical practice.





1

Lupus Science & Medicine

Activation of the type I interferon (IFN) pathway has been implicated in the initiation of SLE and perpetuation of the resultant clinical disease.⁶ Elevated levels of IFN- α are associated with disease activity, flares and tissue injury, especially of the skin, kidney and nervous system.^{7 8} Genetic association studies show polymorphisms in SLE associated with kinase cascade signalling genes for type I IFNs and in IFN-regulated genes (IRGs).⁹ ¹⁰ Further, gene expression profiling studies have identified an IFN signature (IS), or a prominent upregulation of mRNA transcripts encoded by IRGs, in a subset of patients with SLE and other autoimmune diseases.¹⁰⁻¹² It has been hypothesised that dysregulation of the type I IFN signalling pathway could override immune tolerance mechanisms, leading to formation of autoreactive antibodies.¹³ Therefore, blockade of the type I IFN pathway, especially of IFN- α , may be a strategy for selective immunosuppression in patients with SLE with elevated IS expression whose pathophysiology of disease is driven by IFN dysregulation.¹⁴

We previously described an application of a quantitative reverse transcription PCR (qPCR biomarker assay to accurately measure the IFN signature in patients with SLE using a three-gene surrogate called the IS metric (ISM).¹⁵ Here, we describe the development of the ISM assay, its validation in multiple SLE clinical trial cohorts, and characterise the clinical and serological features associated with the ISM-Low and ISM-High subsets of patients with SLE.

PATIENTS AND METHODS

Cohorts of patients with SLE

All patients met the American College of Rheumatology criteria for SLE. This trial is registered (NCT00962832) on the ClinicalTrials.gov website. For purposes of executing clinical trials with different end points, patients with SLE are characterised predominantly as patients with extrarenal lupus (ERL) or as patients with lupus nephritis (LN). The following cohorts were evaluated: 61 patients with ERL in the University of Michigan observational cohort, 60 patients with mild ERL enrolled in the rontalizumab Phase I trial,¹⁶ 135 patients with moderate-severe ERL in the EXPLORER rituximab trial,¹⁷ 80 patients with moderate-severe LN in the LUNAR rituximab trial¹⁸ and 238 patients with moderate-severe ERL in the ROSE rontalizumab (anti-IFN- α monoclonal antibody) trial.¹⁹ Further descriptions are in the online supplementary methods. Healthy control subjects (n=85) were recruited by the Genentech blood donation programme for research use of blood samples, and were age matched and gender matched to the lupus trial patients.

Blood RNA extraction and microarray analysis

Detailed methodology is presented in the online supplementary methods. RNA from SLE cohorts was isolated from peripheral blood mononuclear cells (PBMCs) and/ or PAXgene whole blood samples. Genome-wide expression data were obtained using Affymetrix Human Genome U133 Plus V.2.0 expression microarrays, and data analysis was performed using Bioconductor R packages.²⁰ Microarray ISM scores were derived from the scaled geometric mean across included genes for each sample subjected to exponentiation by 2. Microarray data have been submitted to the gene expression omnibus (GEO) repository under the accession number GSE50772.

Development of the ISM PCR assay

The qPCR assays were developed with primers and fluorescent dye-labelled probes for IRGs and housekeeping genes and run on the Agilent Mx2500P and the Applied Biosystems 7900HT platforms. The three-gene ISM score was calculated using expression values from the gene complementary DNAs (cDNAs) CMPK2, EPSTI1, HERC5, and normalised using the housekeeping gene TFRC. The ISM score was calculated from the mean of the CMPK2, EPSTI1 and HERC5 cycle threshold (Ct)—TFRC C_t (Δ Ct) values and multiplied by -1 to give the correct directionality of relative log₂-scaled expression. Baseline ROSE trial samples were assessed using this assay developed on the Cobas 4800 platform (Roche Molecular Systems) as a prototype diagnostic test.

Measurement of serum analytes

Levels of serum creatine were determined with Roche Modular Analysers using a modified Jaffe reaction. Levels of complement C3 and C4 components were determined using standard immunonephelometry methods. AntidsDNA, anti-SSA/Ro, anti-SSB/La, anti-ribonucleoprotein (RNP) and anti-Sm were measured using a bead-based immunoassay (AtheNA Multi-Lyte antinuclear antibodies Test System). Serum concentrations of B cell-activating factor of the tumour necrosis factor family (BAFF) were determined by an ELISA using monoclonal antibodies developed at Genentech, with a standard curve using recombinant human BAFF. Serum bioactivity was determined using U-937 cells stably transfected with the pGL4.14 vector containing a 1.5 kb fragment of the myxovirus influenza resistance 1 promoter,²¹ where samples were added to reporter cells for 24 h alongside a standard curve of recombinant IFN-a for quantification. Luciferase levels were measured by Luminometer instrumentation.

Statistical analysis

Demographics and key baseline patient characteristics were compared using descriptive statistics, that is, differences between ISM-Low and ISM-High patients in mean values. Statistical testing was performed using the Wilcoxon rank-sum test or the Fisher's exact test as appropriate. No adjustments for multiple comparisons were performed for individual statistical tests. The longitudinal stability of the ISM was evaluated in patients who received placebo, and data were used to establish the probability of reproducibility (P_{rep}) of ISM-Low or ISM-High categorisation (range 0–100%) when samples

were measured from the same patient at baseline, Week 2 on treatment and monthly visits through Week 36. P_{rep} was derived as the average proportion of times that the ISM status observed at baseline was maintained across all available time points. The effect of rontalizumab treatment on the ISM was evaluated with a longitudinal mixed-effects linear model with an autoregressive covariance matrix. The model included the fixed effects of treatment, visit, baseline ISM score and baseline ISM score-by-treatment interaction. Multivariate model selection for determining covariates associated with the ISM score at baseline was performed using stepwise selection with PROC GLMSELECT in Statistical Analysis System (SAS) V.9.2 (SAS Institute, Cary, North Carolina, USA).

RESULTS

Derivation of the ISM biomarker

Using unsupervised hierarchical clustering of samples and probe sets derived from expression microarray analysis of PBMCs in the University of Michigan SLE cohort, we observed that patients with SLE (n=61) and control patients (n=20) had distinct gene expression at the genome-wide level (see online supplementary figure S1). A cluster of 128 genes had high representation of IRGs, relating to the previously described IS, which was elevated in $\sim 50\%$ of patients with SLE (figure 1A and see online supplementary table S4). Composites of only a few genes could yield scores that were highly correlated to the score calculated from the 128-gene cluster (figure 1B). Quantitative PCR of whole blood RNA samples obtained from the same patients in this training cohort for a threegene combination (HERC5, EPSTI1 and TYKI/CMPK2) was a robust surrogate for the original 128-gene IS observed in the matching PBMC RNA samples (figure 1C) -defined hereafter as the ISM. Therefore, the magnitude of the microarray IS that was originally identified by microarray analysis of PBMC RNA samples could be measured accurately using a three-gene qPCR assay of corresponding whole blood RNA samples.

ISM profile in healthy controls and different lupus populations

The ISM scores were generated from whole blood RNA samples from a cohort of 85 additional healthy control subjects and pretreatment samples from patients enrolled in the Phase I trial of rontalizumab (anti-IFN- α) in mild SLE, the EXPLORER trial of rituximab (anti-CD20) in moderate-severe ERL and from the LUNAR trial of rituximab in moderate-severe LN (see Methods). Consistent with the Michigan SLE cohort data, a subset of patients with ERL and LN had significantly elevated ISM scores while other patients had ISM scores that overlapped those of healthy control subjects (figure 2A). The lupus populations exhibited a strong bimodal ISM separation, indicating two distinct patient subsets. A threshold for the ISM score at the 95th centile of the healthy subject ISM scores was calculated to be at an ISM score of 1, separating patients with

lupus into ISM-Low and ISM-High subsets. Compared with the ISM-Low subjects, whose median ISM score of -0.8overlapped with the control population, the ISM-High subset had a median ISM score ≥ 2.3 , corresponding to >eightfold higher expression of the ISM genes (figure 2B) in ISM-High versus ISM-Low subsets. The bimodal distribution and elevation of ISM was consistent in patients from the ROSE Phase II trial (figure 2C), with the median for ISM-High patients indicating >sevenfold higher IS expression than in ISM-Low patients (figure 2D).

Baseline demographics by ISM across four SLE populations

In a mild ERL population enrolled in a Phase I study,¹⁶ the ratio of ISM-Low to ISM-High patients was approximately 1:1. In contrast, in a moderate-to-severe ERL population (EXPLORER) and an LN population (LUNAR), 70-75% of patients were ISM-High (table 1) at baseline. Furthermore, 76% of patients in the ROSE moderate-to-severe SLE trial were designated ISM-High. Gender did not differ between ISM-Low and ISM-High patients in any of the cohorts. However, in the Phase I EXPLORER and LUNAR trials, the ISM-Low patients tended to be older than the ISM-High patients-a trend that was also seen in the ROSE trial-with a 5-year mean difference between the groups. While there was a trend for increased representation of the African-American ethnicity in ISM-Low versus ISM-High subsets of the ERL cohorts, the prevalence of such patients in the studies was low and the trend was not observed in the LUNAR trial.

Clinical disease activity by ISM across four lupus cohorts

Within each SLE trial, global disease activity did not differ between ISM-Low and ISM-High patient subsets as defined by the British Isles Lupus Assessment Group (BILAG) activity index, Safety of Estrogen in Lupus Erythematosus-National Assessment (SELENA)-Systemic Lupus Erythematosus Disease Activity Index (SLEDAI) score, or Physician Global Assessment (table 1). Within the context of enrolment criteria based on SELENA-SLEDAI or BILAG measures and/or renal impairment, disease activity at baseline did not differ between ISM-Low and ISM-High patients with SLE. Furthermore, mucocutaneous involvement did not differ between the two ISM subsets across the four studies. There was a consistent trend for increased musculoskeletal involvement in the ISM-Low versus ISM-High patients, but this was not statistically significant.

In addition, data was collected in the ROSE trial for Cutaneous Lupus Erythematosus Disease Area and Severity Index (CLASI) as well as for the Functional Assessment of Chronic Illness Therapy (FACIT) measurement system of fatigue. Of note, we observed a significant association between ISM-High status and the CLASI damage score; ISM-High (n=180) mean 2.1, SD 5.7, versus ISM-Low (n=58) mean 0.3, SD 1.0, Wilcoxon p=0.0209. However, we did not see a significant association between ISM status and the CLASI activity score; ISM-High mean 5.8 (SD 6.5) versus ISM-Low mean 7.0



Figure 1 Derivation of the blood interferon signature (IS) using a small subset of interferon-regulated genes (IRGs). (A) Visualisation of the 128 IRG cluster after unsupervised hierarchical clustering of genome-wide microarray expression data derived from peripheral blood mononuclear cell (PBMC) samples in either the University of Michigan systemic lupus erythematosus (SLE) cohort or healthy controls. SLE or control samples are indicated by the top colour bar. The heat map colour scale reflects z-score values of gene expression levels (colour bar on left). (B) Spearman's correlations between the whole 128-gene interferon signature metric (ISM) and ISMs calculated from subsets of these 128 genes of sizes ranging from 1 to 128, showing that small numbers of genes yield ISMs very similar to the 128-gene ISM. The genes that compose a subset of size n were selected by first calculating each of the 128 genes' correlation with the 128-gene ISM, and then picking the genes defining n with the highest correlations to the whole 128-gene ISM. (C) ISM from the microarray expression data of 128 interferon signature genes correlates highly with the qPCR ISM generated from whole blood RNA samples using three selected genes. ρ is from Spearman's correlation.

(SD 8.0), p=0.32. Further, there was a trend observed for elevated FACIT fatigue scores in ISM-High patients; ISM-High mean 27.5 (SD 1.43) versus ISM-Low mean 24.3 (SD 13.7), p=0.074.

Increased serological manifestations in ISM-High patients with lupus

In the ERL trials, the ISM-High subset had decreased levels of C3 and C4 complement components relative to the ISM-Low subset, and these differences were consistent in the ROSE trial (table 2). In contrast, C3 and C4 levels did not differ in the ISM subpopulations in the LUNAR LN trial. ISM-High status was associated with elevation of anti-dsDNA titres (figure 3A) and increased incidence of positive extractable nuclear antigen antibody status (figure 3B) as compared with ISM-Low patients with ERL. Levels of these autoantibodies did not significantly differ between ISM-High and ISM-Low patients with LN. Serum BAFF levels were elevated in the ISM-High versus ISM-Low subsets in patients with ERL and LN (figure 3C, table 2). In patients with ERL, type I IFN serum bioactivity was also elevated in the ISM-High



Figure 2 Comparison of the interferon (IFN) signature metric (ISM) magnitude and distribution in healthy controls versus patients with systemic lupus erythematosus (SLE). (A) Distribution of pretreatment ISM values in the previously described healthy control cohort versus bimodal ISM distributions in a cohort of patients with mild-SLE (rontalizumab Phase I trial), the rituximab EXPLORER SLE trial and the rituximab LUNAR lupus nephritis (LN) trial. The dashed line indicates cut-off used for ISM-Low versus ISM-High patients and incidences of ISM values across the patient cohorts are plotted as densities. (B) Median, IQRs, and 5th and 95th centile values for the pretreatment ISM in a healthy control cohort (n=85), and in ISM-Low versus ISM-High patient subsets in a moderate-to-severe extrarenal lupus (ERL) cohort (rituximab EXPLORER trial, n=131) and a moderate-to-severe LN cohort (rituximab LUNAR trial, n=80). (C) Bimodal ISM distribution observed in the moderate-to-severe SLE cohort (ROSE rontalizumab Phase II trial, n=238) with cut-off value indicated with dashed line. (D) Median, IQRs, and 5th and 95th centile values for the pretreatment ISM in the ROSE trial (n=238).

patients relative to ISM-Low patients (figure 3D); this was not assessed in patients with LN. We further assessed C reactive protein (CRP) levels between ISM-High and ISM-Low patients, and noted that while CRP levels were elevated in ISM-High patients in the ROSE trial; ISM-High mean 0.81 mg/dL (SD 1.43) versus ISM-Low 0.42 mg/dL (SD 0.6), p=0.019, there were no significant elevations in CRP levels between ISM-High patients versus ISM-Low patients across the other trials (table 1). It was further noteworthy that across the four trials, total leucocyte counts and lymphocyte, neutrophil, monocyte and eosinophil counts were consistently lower in ISM-High patients (see online supplementary table S2).

Multivariate analysis of the ISM score

Absolute counts of CD4, anti extractable nuclear antigen status, levels of BAFF and anti-dsDNA, and SLE disease

duration were independently associated with the ISM score (see online supplementary table S3). Longer disease duration was the only non-serological component of the model associated with lower ISM scores. A statistically non-significant trend for an association between the ISM score and age at baseline was noted.

Determination of longitudinal stability of the ISM

Patients receiving placebo (n=61) had a 94% probability of being consistently categorised in the same ISM subset throughout the 36-week period (see online supplementary table S4). Patients with an ISM score closer to the cut-off of 1 (>–0.5 to ≤ 2 , n=17) had a lower likelihood (81% to 85%) of maintaining their ISM status, while patients with ISM scores further from the cut-off (\leq –0.5 or >2, n=44) had a higher (97% to 100%)

Total (N=135) 39.9 (11.2)

8.9 (7.7)

94

24.4

15.6

56.3

14.5 (5.2)

10.9 (6.4)

54.1 (17.0)

3.7

83.7

74.8

0.7 (0.2)

0.81 (1.41)

1.5

EXPLORER trial—moderate-to-severe SLE

ISM-Low (n=40)

42.8 (11.4)

8.2 (7.9)

100

12.5

12.5

72.5

2.5

92.5

60.0

0.0

13.1 (4.7)

50.5 (20.4)

9.3 (5.2)

0.8 (0.2)

0.89 (1.33)

ISM-High (n=95)

38.8 (10.9)

9.2 (8.0)

92

29.5

16.8

49.5

80.0

81.1

2.1

15.2 (5.2)

9.2 (8.0)

0.7 (0.2)

0.79 (1.44)

55.7 (15.3)

4.2

	Phase I rontalizum	ab trial-mild S
Parameter	ISM-Low (n=32)	ISM-High (I
Age (years), mean (SD)	48.2 (9.7)	46.5 (10.5)
SLE duration (years), mean (SD)	7.5 (5.6)	10.4 (7.1)
Female, %	97	93
Race/ethnicity, %		
Black	12.5	42.9
Hispanic*	0	7.1
White	87.5	50
Other	0	0
BILAG index global, mean (SD)	N/A	N/A
SELENA-SLEDAI, mean (SD)	4.1 (2.6)	2.9 (2.7)
PGA, mean (SD)	25.5 (15.1)	17.3 (17.6)
Musculoskeletal involvement, %†	37.5	28.6
Mucocutaneous involvement, %†	84.4	64.3
Proteinuria (>0.5 g), %	0.0	0.0
Serum creatine (mg/dL) mean (SD)	0.8 (0.1)	0.9(0.2)

Table 1 Demographics and clinical baseline characteristics by ISM status of patients in four lupus studies

	LUNAR Trial‡—LN			Phase II Rontalizumab ROSE Trial—Moderate-to SLE		
	ISM-Low (n=21)	ISM-High (n=59)	Total (N=80)	ISM-Low (n=58)	ISM-High (n=180)	Total (N=238)
Age (years), mean (SD)	31.4 (10.9)	30.0 (9.1)	30.4 (9.6)	42.6 (11.8)	37.6§ (10.9)	38.8 (11.3)
SLE duration (years), mean (SD)	6.2 (6.7)	4.7 (4.6)	5.1 (5.2)	7.2 (7.7)	6.2 (5.8)	6.5 (6.3)
Female, %	81	93	90	97	93	94
Race/ethnicity, %						
Black	19.1	20.3	20	8.6	16.1	14.3
Hispanic*	57.1	40.7	45	25.9	36.1	33.6
White	23.8	37.3	33.8	55.2	43.3	46.2
Other	0	1.7	1.3	10.3	4.4	5.9
BILAG index global, mean (SD)	15.8 (5.8)	14.3 (6.5)	14.7 (6.3)	11.9 (4.9)	11.0 (4.6)	11.2 (4.7)
SELENA-SLEDAI, mean (SD)	N/A	N/A	N/A	9.2 (2.6)	10.0 (3.5)	9.8 (3.3)
PGA, mean (SD)	51.6 (24.2)	49.4 (21.2)	50.0 (21.9)	52.1 (17.2)	57.5 (16.0)	56.2 (16.4)
Musculoskeletal involvement, %†	28.6	18.6	21.3	98.3	95.6	96.2
Mucocutaneous involvement, %†	38.1	32.2	33.8	77.6	75.6	76.1
Proteinuria (>0.5 g), %	100	94.9	96.3	0.0	1.7	1.3
Serum creatine (mg/dL), mean (SD)	1.0 (0.4)	1.0 (0.6)	1.0 (0.5)	0.8 (0.2)	0.8 (0.2)	0.8 (0.2)
Serum C reactive protein (mg/dL), mean (SD)	0.4 (0.56)	0.63 (1.08)	0.57 (0.97)	0.42 (0.6)	0.81 (1.43)	0.71 (1.29)

0.34 (0.35)

LE

າ=28)

Total (N=60)

47.4 (10.0)

8.8 (6.4)

95

26.7

3.3

70

0

N/A

33.3

75.0

0.0

3.6 (2.7)

0.8 (0.2)

0.47 (0.58)

21.7 (16.7)

*Hispanic defined as a person of self-declared race other than black or white, and self-declared ethnicity Hispanic or Latino.

0.59 (0.71)

†BILAG index scores of A, B or present on SELENA-SLEDAI.

Serum C reactive protein (mg/dL), mean (SD)

‡LUNAR patients missing an ISM score were not included.

§P=0.0042, ISM-Low versus ISM-High.

BILAG, British Isles Lupus Assessment Group; ISM, interferon signature metric; LN, lupus nephritis; N/A, not applicable; PGA, Physician Global Assessment; SELENA, Safety of Estrogen in Lupus Erythematosus-National Assessment; SLE, systemic lupus erythematosus; SLEDAI, Systemic Lupus Erythematosus Disease Activity Index.

Kennedy WP, Maciuca R, Wolslegel K, et al. Lupus Science & Medicine 2015;2:e000080. doi:10.1136/upus-2014-000080

Table 2 Baseline serological characteristics versus ISM status for patients from the Rontalizumab Phase I, rituximab EXPLORER, rituximab LUNAR and rontalizumab Phase II ROSE studies

	Phase I Study				EXPLORER Stud	у	
Serology parameter	ISM-Low (n=32)	ISM-High (n=28)	Total (N=60)	Serology Parameter	ISM-Low (n=40)	ISM-High (n=95)	Total (N=135)
C3 (mg/dL), mean (SD)	133.5 (29.6)	120.6 (36.1)	128.7 (28.1)	C3 (mg/dL), mean (SD)	120.6 (36.1)	92.7 (30.9)	100.9 (34.9)
C4 (mg/dL), mean (SD)	25.3 (7.7)	21.2 (9.1)	23.4 (7.4)	C4 (mg/dL), mean (SD)	20.2 (9.1)	13.6 (7.2)	15.5 (8.3)
Anti-dsDNA (IU/mL), median (IQR)	12 (12–15)	15 (12–45)	12 (12–23)	Anti-dsDNA (IU/mL), median (IQR)	12 (12–79)	71 (29–239)	44 (13–224)
Anti-dsDNA+ (≥30 U/mL), %	9.4	32.1	20.0	Anti-dsDNA+ (≥30 U/mL), %	35.0	72.6	61.5
Anti-ENA+, %*	6.3	60.7	31.7	Anti-ENA+, %*	10.0	59.6	44.8
Anti-SSA/Ro52+ (≥120 IU/mL), %	3.1	46.4	23.2	Anti-SSA/Ro52+ (≥120 IU/mL), %	7.5	41.5	31.3
Anti-SSB/La+ (≥120 IU/mL), %	0.0	7.1	3.3	Anti-SSB/La+ (≥120 IU/mL), %	0.0	17.0	11.9
Anti-RNP+ (≥120 IU/mL), %	3.1	21.4	11.7	Anti-RNP+ (≥120 IU/mL), %	2.5	26.6	19.4
Anti-Sm+ (≥120 IU/mL), %	0.0	10.7	5.0	Anti-Sm+ (≥ 120 IU/mL), %	2.5	23.4	17.2
BAFF† (IU/mL), median (IQR)	1782	2542	1975	BAFF† (IU/mL), median (IQR)	2085	3750	2870
	(1468–2202)	(1842–3031)	(1589–2862)		(1630–2735)	(2250–6690)	(2000–4960)

LUNAR study				ROSE study				
Serology parameter	ISM-Low (n=21)	ISM-High (n=59)	Total (N=80)	Serology Parameter	ISM-Low (n=58)	ISM-High (n=180)	Total (N=238)	p Value‡
C3 (mg/dL), mean (SD)	77.7 (32.0)	75.0 (30.2)	75.7 (30.5)	C3 (mg/dL), mean (SD)	121.7 (30.7)	95.8 (32.6)	102.1 (34.0)	<0.0001
C4 (mg/dL), mean (SD)	13.7 (8.4)	14.3 (7.8)	14.2 (7.9)	C4 (mg/dL), mean (SD)	21.4 (8.5)	14.5 (9.9)	16.2 (10.0)	<0.0001
Anti-dsDNA (IU/mL), median (IQR)	66 (23–209)	154 (57–347)	116 (39–303)	Anti-dsDNA (IU/mL), median (IQR)	12 (12–36)	91 (24–273)	53 (14–193)	<0.0001
Anti-dsDNA+ (≥30 U/mL), %	57.1	84.4	77.5	Anti-dsDNA+ (≥ 30 U/mL), %	34.5	70.9	62.0	<0.0001
Anti-ENA+, %*	52.4	57.6	56.3	Anti-ENA+, %*	19.0	73.3	60.1	<0.0001
Anti-SSA/Ro52+ (≥120 IU/mL), %	23.8	25.4	25.0	Anti-SSA/Ro52+ (≥ 120 IU/mL), %	13.8	49.4	40.8	<0.0001
Anti-SSB/La+ (≥120 IU/mL), %	9.5	8.5	8.8	Anti-SSB/La+ (≥ 120 IU/mL), %	5.2	16.1	13.4	0.0443
Anti-RNP+ (≥120 IU/mL), %	23.8	35.6	32.5	Anti-RNP+ (≥ 120 IU/mL), %	8.6	37.8	30.7	<0.0001
Anti-Sm+ (≥120 IU/mL), %	28.6	30.5	30.0	Anti-Sm+ (≥120 IU/mL), %	0.0	28.3	21.4	<0.0001
BAFF† (IU/mL), median (IQR)	1860	3550	3010	BAFF (IU/mL), median (IQR)	1935	2990	2660	<0.0001
	(1590–2940)	(2330–5700)	(1960–5080)		(1565–2590)	(2180–4120)	(1940–3890)	

ENA, extractable nuclear antigen; ISM, interferon signature metric; IU, international unit. *Positive for two or more autoantigen reactivities of the following: SM+, RNP+, SSA+, SSB+. †For Phase I, BAFF data are available for n=28 patients (14 ISM-Low, 14 ISM-High). ‡p Value represents ISM-Low versus ISM-High.

Biomarker studies



Figure 3 Association of pretreatment interferon (IFN) signature metric (ISM) status with serum autoantibodies, BAFF levels and type I IFN bioactivity. Patients with extrarenal lupus (ERL) from the rontalizumab Phase I and Phase IIa ROSE trials and from the rituximab EXPLORER trial were pooled and stratified by ISM-Low (n=118) or High (n=307) status. Patients with renal lupus from the rituximab LUNAR trial were also examined (ISM-Low n=21; ISM-High n=59). (A) Levels of anti-dsDNA antibodies plotted as median, IQRs, and 5th and 95th centiles. Statistical significance was determined using the Wilcoxon test. (B) Presence or absence of extractable nuclear antigen (ENA) antibodies plotted as percentages of the patient subgroups. Statistical significance was determined using Fisher's exact test. (C) Levels of serum BAFF plotted as median, IQRs, and 5th and 95th centiles. Statistical significance was determined using fisher's exact test. (D) Levels of serum myxovirus influenza resistance 1 (MX1)-induced bioactivity in ROSE patients with ERL are plotted as median, IQRs, and 5th and 95th centiles. Statistical significance was determined using the Wilcoxon test.

likelihood of maintaining their status over the same period of time.

Decrease in ISM score after dosing of patients with SLE with rontalizumab (anti-IFN- α)

Treatment with rontalizumab decreased the ISM score relative to placebo during the 24-week treatment period based upon a longitudinal model incorporating all of the time points that demonstrated an intercept that was statistically significantly lower for the rontalizumab groups versus the placebo groups (p<0.01), reflecting a pharmacodynamic response for IFN- α blockade. Application of a mixed-effect model also supported an interaction between baseline ISM score and treatment group (p<0.01), where differences between active treatment and placebo were greater for patients with lower versus higher baseline ISM scores. Despite this pharmacodynamic response, treatment of ISM-High patients with rontalizumab did not decrease the ISM-High scores down to the levels observed in ISM-Low patients (figure 4).

DISCUSSION

In this study, we confirmed the presence of an IFN-inducible gene expression signature (IS) in the peripheral blood of patients with SLE and developed a novel and simplified qPCR-based biomarker (ISM) test to quantitate the IS. Importantly, we showed that the magnitude of the ISM biomarker was highly similar between PBMCs and whole blood RNA samples obtained from the same subjects. At baseline, patients

6



Figure 4 Decreased interferon (IFN) signature metric (ISM) magnitude after dosing patients with systemic lupus erythematosus (SLE) with rontalizumab (anti-IFN- α). Patients with SLE enrolled in the ROSE trial were dosed over 24 weeks with rontalizumab or placebo (see Methods) and ISM values were determined from pretreatment and post-treatment blood RNA samples collected at monthly frequencies. Blood ISM scores were calculated for pretreatment and post-treatment time points and plotted versus time as mean±SEM for the ISM-High (n=55 for placebo and 120 for rontalizumab) and ISM-Low (n=24 for placebo and 33 for rontalizumab) patient groups as determined at baseline. The ISM scale (v axis) is split between 0 and 2 for clarity, and data from the intravenous (750 mg every 4 weeks) and subcutaneous (300 mg every 2 weeks) dose arms for rontalizumab versus the corresponding placebo arms were pooled for this analysis. Assessment of statistical differences between placebo and rontalizumab groups was based upon a longitudinal model assessing all time points.

with SLE from four different clinical trials consistently fall into one of two subsets—ISM-High and ISM-Low and the bimodal distribution of patients with SLE by the ISM agrees with previous reports.¹

The bimodal distribution expression of the IRG in ISM-Low and ISM-High patients with SLE is a distinct feature of this biomarker. High-ISM patients with SLE had markedly elevated IRG expression, while patients in the ISM-Low subset had a distribution of gene expression that overlapped with the predominantly unimodal ISM measured in healthy subjects, despite the presence of active disease. The relative distribution of patients who are ISM-High or ISM-Low varied, in part, with disease activity based on BILAG or SELENA-SLEDAI measures, but this appears to be a consequence of the trial inclusion criteria. Thus, ISM status across SLE trial populations varied from an incidence of ~50% ISM-High scores in a mild disease population in a Phase I study¹⁶ to ~75% in moderate-to-severe SLE disease

populations in Phase II/III studies that represent patients with ERL and LN.

Despite an apparently increased incidence of ISM-High patients in Phase II/III trials in populations with moderate-to-severe disease activity compared with Phase I studies in populations with mild disease activity, the ISM status within each trial's cohorts did not correlate with differences in disease activity. Our observations are based on the consistent use of a platform-the qPCR-based ISMacross different trial populations, in contrast to reports of IS associations with increased SLE disease activity based on differing methods applied across different cohorts to measure the IS.¹ ¹⁰ ¹⁴ For example, in the Phase II ROSE study with patients with moderate-to-severe SLE, ISM status did not correlate with differences in mean measures of disease activity assessed by BILAG or SELENA-SLEDAI.¹⁹ Further, the skewed incidence of ISM-High patients in moderate-to-severe disease was fairly similar in the ERL and LN populations, indicating that the ISM distribution is not broadly specific with end-organ activity in most patients with SLE. In the EXPLORER¹⁷ and ROSE¹⁹ studies, we did not observe consistent correlations between mucocutaneous disease manifestations and ISM status, and we observed only a modest trend in elevated musculoskeletal disease manifestations in ISM-Low compared with ISM-High patients across the four trials. Of note, we did observe increased CLASI damage scores, but not CLASI activity scores, in the ROSE trial where this instrument was used that suggests the possibility of an association of the ISM biomarker with aspects of cutaneous disease. We also observed a trend for elevated FACIT fatigue scores in ISM-High patients in the ROSE trial. The possibility still remains, however, that there are distinct clinical SLE pathotypes within the ISM subsets that have not yet been identified. We did observe that the ISM-High subset was slightly younger overall across the trials and, in at least the ERL populations, had a lower frequency of white patients, consistent with previous observations.¹⁰

ISM status does correlate with serological differences between the two subpopulations across multiple trial cohorts with extrarenal SLE. The presence of characteristic SLE autoantibodies, hypocomplementaemia, cytokine production and serum IFN bioactivity varied between ISM-High and ISM-Low patients with lupus. These findings are consistent with previous reports^{1 10 16} and underscore the role of type I IFNs in systemic immune activation, including extrafollicular differentiation of autoreactive B cells to antibody-secreting plasmablasts via regulation of toll-like receptors (TLRs) and increased survival via BAFF-induced signalling.⁶ Thus, the ISM could be a biomarker of systemic immune activation that comanifests with serological abnormalities. Indeed, our multivariate analysis confirmed independent association of ISM magnitude with T cell counts, autoantibodies, cytokine levels and disease duration parameters.

We found no significant differences between the ISM-Low and ISM-High subpopulations in autoantibody presence or titre in patients with LN, and observed

Lupus Science & Medicine

comparable levels of hypocomplementaemia in these subpopulations. So, although immune-serology differences exist between ISM-High and ISM-Low patients with ERL, additional drivers of the IS likely exist in patients with LN, giving rise to a bimodal ISM distribution. This suggests that the regulation of the IRGs could be due in part to an amplification loop where, upon reaching a threshold, genes could be upregulated by additional upstream signals. These signals may include, but are not limited to, nucleic acid-containing IC activation of endosomal TLR7 and TLR9 nucleic acid sensors such as the retinoic acid-inducible gene I-like receptors, nucleotide oligomerisation domain-like receptors, and cyclic GMP-AMP synthase, which lead to type I IFN expression.^{6 24 25}

Although our analysis was not designed to address the stability of the ISM on a particular background of immunosuppressants, it appears that the most frequently used immunosuppressants (ie, hydroxychloroquine and oral corticosteroids) in lupus management do not significantly impact the ISM status of patients with SLE. In contrast, others have shown that high-dose, pulse intravenous corticosteroids markedly and transiently attenuate the IS in patients with SLE.^{13 26} Further, blockade of IFN- α with rontalizumab in patients with ERL¹⁹ only leads to a partial reduction in ISM score by 2–4 weeks of treatment. This is consistent with observations in patients with SLE treated with a different anti-IFN- α antibody, sifalimumab.²⁷

One potential contributing factor to this partial ISM decrease could include suboptimal dosing of patients with anti-IFN-a antibodies, and treatment of patients with higher amounts of this therapy could result in a greater magnitude of decrease of the ISM score. It is also important to consider the complexity of the genome-wide gene expression profile, and the potential limitations of our approach to derive a simplified biomarker that integrates the magnitude of a large number of genes. Indeed, a recent study using modular transcriptional analysis has demonstrated that different a priori-defined IFN-inducible modules have differential associations with disease activity such as SELENA-SLEDAI scores, serology and presence of flare.²⁸ Further, the different IFN-inducible modules had varying degrees of longitudinal stability, and likely reflected differential contributions of IFN-β and IFN-γ in addition to IFN- α . Clearly, therefore, the IS in patients with SLE is driven by factors beyond IFN-a signalling and may include overlapping Jak-Stat signalling mechanisms used by other cytokines²⁹ and/or sustained IC derived from pre-existing plasma cells. In this regard, it has recently been reported that blockade of the Lymphotoxin/LIGHT pathway using baminercept, a lymphotoxin-β receptor-immunoglobulin fusion protein, reduced the IFN signature in the blood of patients with rheumatoid arthritis with elevated baseline signatures³⁰ suggesting a role of the Lymphotoxin/LIGHT pathway as an upstream modulator of IFNs in RA and potentially SLE. Furthermore in this study, patients with RA and SLE had lymphopenia associated with elevation of blood IFN signature, and treatment of patients with RA

with baminercept subsequently increased lymphocyte counts. Consistent with this observation, it was noteworthy that the ISM-High patients within the trials examined in this study also had reduced circulating leucocyte counts including lymphocytes, monocytes, neutrophils and eosinophils as compared with ISM-Low patients. These data are consistent with a role for IFN signalling in retaining immune cells in tissues during inflammatory disease and therefore contributing to lymphopenia. Additional mechanistic studies are warranted to investigate upstream signals that can specifically induce genes also induced by type I IFNs.

In conclusion, we have characterised the ISM as a surrogate of the peripheral blood global IS, first identified in patients with SLE.^{10–12} The method identifies distinct ISM-Low and ISM-High SLE populations, and has the potential to provide a useful approach to identify and stratify the heterogeneous SLE population in the context of randomised control clinical trials and clinical practice. The correlation of baseline ISM status and clinical response has recently been evaluated in the ROSE Phase II randomised, controlled trial of the safety and efficacy of rontalizumab in extrarenal lupus.¹⁹

Author affiliations

¹Departments of Early Clinical Development, Genentech, South San Francisco, California, USA

²Department of Biostatistics, Genentech, South San Francisco, California, USA ³Department of ITGR Diagnostics Discovery, Genentech, South San Francisco, California, USA

⁴Department of Bioinformatics and Computational Biology, Genentech, South San Francisco, California, USA

⁵Department of Bioanalytical Sciences, Genentech, South San Francisco, California, USA

⁶Department of Pharmacodynamic Biomarkers ITGR, Genentech, South San Francisco, California, USA

⁷Department of Late Stage Immunology Product Development, Genentech, South San Francisco, California, USA

⁸Department of Medicine, University of Michigan and the Ann Arbor VA Hospital, Ann Arbor, Michigan, USA

⁹Department of ITGR Human Genetics, Genentech, South San Francisco, California, USA

Acknowledgements The authors thank Rachel Brock and Sushma Yarlagadda for assistance in processing patient samples from the Michigan SLE cohort, Lauren Nguyen and Zachary Martinez for immunoassay analysis of patient serum samples, and Dr Jane Grogan for scientific discussion of the data. Writing assistance was provided by Genentech.

Contributors All authors made substantial contributions to the conception or design of the work; or the acquisition, analysis or interpretation of data for the work; and drafting the work or revising it critically for important intellectual content; and final approval of the version to be published; and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding This study was financially supported by Genentech, a member of the Roche Group.

Competing interests WPK, RM, KW, WT, ARA, CC, AM, JMM, PB, JCD., TWB, and MJT: Employees and shareholders of Roche. BCR: Research support from Genentech. All of the authors with the exception of BCR are current or former employees of Genentech, a member of the Roche group.

Kennedy WP, Maciuca R, Wolslegel K, et al. Lupus Science & Medicine 2015;2:e000080. doi:10.1136/lupus-2014-000080

Biomarker studies

Ethics approval The study was approved by the relevant Institutional Review Boards/Independent Ethics Committees and conducted in accordance with the ethical principles of the Declaration of Helsinki.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Microarray data have been submitted to the GEO repository and has the accession number GSE50772.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http:// creativecommons.org/licenses/by-nc/4.0/

REFERENCES

- Koffler D, Agnello V, Thoburn R, *et al.* Systemic lupus erythematosus: prototype of immune complex nephritis in man. *J Exp Med* 1971;134:169s–79s.
- Truedsson L, Bengtsson AA, Sturfelt G. Complement deficiencies and systemic lupus erythematosus. *Autoimmunity* 2007;40:560–6.
- Yung S, Chan TM. Anti-DNA antibodies in the pathogenesis of lupus nephritis--the emerging mechanisms. *Autoimmun Rev* 2008;7:317–21.
- Strand V, Chu AD. Measuring outcomes in systemic lupus erythematosus clinical trials. *Exp Rev Pharmacoec Outcomes Res* 2011;11:455–68.
- Wollaston SJ, Farewell VT, Isenberg DA, et al. Defining response in systemic lupus erythematosus: a study by the Systemic Lupus International Collaborating Clinics group. J Rheum 2004;31:2390–4.
- 6. Ronnblom L, Eloranta ML. The interferon signature in autoimmune diseases. *Curr Opin Rheum* 2013;25:248–53.
- Banchereau J, Pascual V. Type I interferon in systemic lupus erythematosus and other autoimmune diseases. *Immunity* 2006;25:383–92.
- Crow MK. Type I interferon in organ-targeted autoimmune and inflammatory diseases. *Arth Res Ther* 2010;12(Suppl 1):S5.
- Bronson PG, Chaivorapol C, Ortmann W, *et al.* The genetics of type I interferon in systemic lupus erythematosus. *Curr Opin Immun* 2012;24:530–7.
- Baechler EC, Batliwalla FM, Karypis G, et al. Interferon-inducible gene expression signature in peripheral blood cells of patients with severe lupus. Proc Nat Acad Sci 2003;100:2610–5.
- Bennett L, Palucka AK, Arce E, *et al.* Interferon and granulopoiesis signatures in systemic lupus erythematosus blood. *J Exp Med* 2003;197:711–23.
- Higgs BW, Liu Z, White B, *et al.* Patients with systemic lupus erythematosus, myositis, rheumatoid arthritis and scleroderma share activation of a common type I interferon pathway. *Ann Rheum Dis* 2011;70:2029–36.

- Ronnblom L, Eloranta ML, Alm GV. The type I interferon system in systemic lupus erythematosus. *Arthritis Rheum* 2006;54:408–20.
- Lichtman EI, Helfgott SM, Kriegel MA. Emerging therapies for systemic lupus erythematosus–focus on targeting interferon-alpha. *Clin Immunol* 2012;143:210–21.
- Morimoto AM, Flesher DT, Yang J, *et al.* Association of endogenous anti-interferon-alpha autoantibodies with decreased interferonpathway and disease activity in patients with systemic lupus erythematosus. *Arthritis Rheum* 2011;63:2407–15.
- McBride JM, Jiang J, Abbas AR, *et al.* Safety and pharmacodynamics of rontalizumab in patients with systemic lupus erythematosus: results of a phase I, placebo-controlled, double-blind, dose-escalation study. *Arthritis Rheum* 2012;64:3666–76.
- Merrill JT, Neuwelt CM, Wallace DJ, *et al.* Efficacy and safety of rituximab in moderately-to-severely active systemic lupus erythematosus: the randomized, double-blind, phase II/III systemic lupus erythematosus evaluation of rituximab trial. *Arthritis Rheum* 2010;62:222–33.
- Merrill J, Buyon J, Furie R, *et al.* Assessment of flares in lupus patients enrolled in a phase II/III study of rituximab (EXPLORER). *Lupus* 2011;20:709–16.
- Kalunian KC, Merrill JT, Maciuca R, *et al.* A Phase II study of the efficacy and safety of rontalizumab (rhuMAb interferon-α) in systemic lupus erythematosus patients (ROSE). Ann Rheum Dis 2015. doi: 10.1136/annrheumdis-2014-206090.
- Bourgon R, Gentleman R, Huber W. Independent filtering increases detection power for high-throughput experiments. *Proc Natl Acad Sci* 2010;107:9546–51.
- 21. Ronni T, Matikainen S, Lehtonen A, *et al.* The proximal interferon-stimulated response elements are essential for interferon responsiveness: a promoter analysis of the antiviral MxA gene. *J Interferon Cytokine Res* 1998;18:773–81.
- Ronnblom L, Alm GV. A pivotal role for the natural interferon alpha-producing cells (plasmacytoid dendritic cells) in the pathogenesis of lupus. *J Exp Med* 2001;194:F59–63.
- Means TK, Latz E, Hayashi F, et al. Human lupus autoantibody-DNA complexes activate DCs through cooperation of CD32 and TLR9. J Clin Invest 2005;115:407–17.
- 24. Takeuchi O, Akira S. Innate immunity to virus infection. *Immunol Rev* 2009;227:75–86.
- Sun L, Wu J, Du F, *et al.* Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* 2013;339:786–91.
- Guiducci C, Gong M, Xu Z, et al. TLR recognition of self nucleic acids hampers glucocorticoid activity in lupus. *Nature* 2010;465:937–41.
- Petri M, Wallace DJ, Spindler A, *et al.* Sifalimumab, a human anti-interferon-alpha monoclonal antibody, in systemic lupus erythematosus: a phase I randomized, controlled, dose-escalation study. *Arthritis Rheum* 2013;65:1011–21.
- Chiche L, Jourde-Chiche N, Whalen E, et al. Modular transcriptional repertoire analyses of adults with systemic lupus erythematosus reveal distinct type I and type II interferon signatures. Arthritis Rheumatol 2014;66:1583–95.
- 29. Stark GR, Darnell JE Jr. The JAK-STAT pathway at twenty. Immunity 2012;36:503–14.
- Bienkowska J, Allaire N, Thai A, *et al.* Lymphotoxin-LIGHT pathway regulates the interferon signature in rheumatoid arthritis. *PLoS One* 2014;9:e112545.

Association of the Interferon Signature Metric with serological disease manifestations but not global activity scores in multiple SLE patient cohorts

William P. Kennedy,¹ Romeo Maciuca,² Kristen Wolslegel,³ Wei Tew,³ Alexander R. Abbas,⁴ Christina Chaivorapol,⁴ Alyssa Morimoto,⁵ Jacqueline M. McBride,⁶ Paul Brunetta,⁷ Bruce C. Richardson,⁸ John C. Davis, Jr.,¹ Timothy W. Behrens,⁹ Michael J. Townsend³

¹Departments of Early Clinical Development, Genentech, South San Francisco, CA, USA
²Biostatistics, Genentech, South San Francisco, CA, USA
³ITGR Diagnostics Discovery, Genentech, South San Francisco, CA, USA
⁴Bioinformatics and Computational Biology, Genentech, South San Francisco, CA, USA
⁵Bioanalytical Sciences, Genentech, South San Francisco, CA, USA
⁶Pharmacodynamic Biomarkers ITGR, Genentech, South San Francisco, CA, USA
⁷Late Stage Immunology Product Development, Genentech, South San Francisco, CA, USA
⁹ITGR Human Genetics, Genentech, South San Francisco, CA, USA
⁸Department of Medicine, University of Michigan and the Ann Arbor VA Hospital, Ann Arbor, MI, USA

Corresponding author:

Michael J. Townsend, Ph.D. ITGR Diagnostics Discovery Genentech, Inc. 1 DNA Way South San Francisco, CA 94080 Tel: 650-225-3239 Email: townsend.michael@gene.com

Key Words: Interferon Signature, SLE, lupus, disease activity, serological manifestations

SUPPLEMENTAL METHODS

SLE patient cohorts

University of Michigan SLE cohort

SLE patients (n=61) were enrolled at the University of Michigan into an observational cohort. Patient enrollment and sample collection were carried out under the supervision of the Institutional Review Board of the University of Michigan Medical School (IRBMED: protocol reference 2002-0339). Demographics and baseline disease characteristics are summarized in Supplemental Table 1. Healthy control subjects (n=20) were age-matched.

Rontalizumab Phase I trial¹⁶

Results of the trial have been published and are present on ClinicalTrials.gov (Identifier NCT00541749). Sixty patients were enrolled with mild disease activity as defined by SELENA-SLEDAI, and continued their pre-enrollment background medications, including nonsteroidal anti-inflammatory drugs (NSAIDs), anti-malarials, and steroids up to 20 mg/day prednisone equivalent, for lupus management.

EXPLORER: Assessment of flares in lupus patients enrolled in a Phase II/III study of rituximab¹⁷

Patients had moderate to severe active extra-renal lupus (ERL), with active disease defined according to the British Isles Lupus Disease Activity Group (BILAG) index, and stable use of 1 immunosuppressive drug at entry. Patients with severe central nervous system or organ-threatening lupus were excluded.

LUNAR: Lupus Nephritis Assessment with Rituximab¹⁸

Patients had Class III/IV LN according to the 2003 International Society of Nephrology/Renal Pathology Society criteria determined by renal biopsy within previous 12 months and presence of proteinuria.

ROSE: ROntalizumab Phase II Trial in Systemic lupus Erythematosus¹⁹

The trial is registered on ClinicalTrials.gov (identifier NCT00962832). Patients had moderate to severe active ERL (active disease defined according to the BILAG index; 1 BILAG A or 2 BILAG B domains of disease involvement and severity), and positive antinuclear antibody (ANA). Patients were excluded if they had lupus nephritis (LN), unstable neuropsychiatric SLE, or a recent history of severe anti-phospholipid antibody syndrome.

Blood RNA extraction and microarray analysis

Samples of RNA from all SLE cohorts were extracted from peripheral blood mononuclear cells (PBMCs) isolated by Ficoll gradient and/or from whole blood samples collected in PAXgene tubes using the RNeasy Mini Kit (Qiagen, Valencia, CA). PBMC RNA samples from the University of Michigan cohort were profiled on Affymetrix Human Genome U133 Plus 2.0 expression microarrays. The microarrays were MAS 5.0 normalized in R 3.0.0; then probe sets were filtered using the default settings of the featureFilter function in the geneFilter Bioconductor package²⁰ to result in one probe set per gene. Samples and filtered probe sets were hierarchically clustered using a Euclidean distance metric and Ward linkage method using hclust (R stats package). Twenty clusters were derived using cutree (k=20) on the probe set dendrogram. Heatmap plots were generated in R. Probe sets were mapped to genes using the hgu133plus2.db (v. 2.9.0) Bioconductor annotation package. Differential expression of genes was assessed by linear modeling via the Limma package (Bioconductor). Moderated *t*statistics from modeling were used to calculate the adjusted *p* values, using the Benjamini-Hochberg correction. Differential expression was defined as gene expression greater than 1.5fold and the adjusted *p* value for the difference in gene expression was less than 0.2. Microarray data have been deposited in the Gene Expression Omnibus data repository (GEO accession number GSE50772).

Supplemental Figure 1: Unsupervised hierarchical clustering of microarray expression data from the University of Michigan SLE cohort. Filtered microarray probes generated from microarray analysis of 20 healthy control subjects and 61 SLE patients underwent unsupervised clustering. The left sided figure bar is the heatmap color scale and refers to z-score values. The location of the Interferon Regulated Gene cluster is demarcated with an open box. The bar above the heatmap refers to diagnosis (SLE in yellow, healthy control in green) for each subject sample (pertaining to each column of the heatmap).



	University of Michigan SLE cohort				
Parameter	ISM-Low (n=28)	ISM-High (n=33)	Total (N=61)		
Age (years), mean (SD)	39 (9.7)	37.6 (11.05)	38.2 (10.4)		
SLE duration (years), mean (SD)	8.2 (8.3)	11.6 (8.3)	9.7 (8.3)		
Female, %	96	97	97		
Race/Ethnicity, %					
Black	29	33	31		
Hispanic	4	0	2		
White	64	67	66		
Other	3	0	1		
SELENA-SLEDAI, mean (SD)	5.8 (3.1)	5.6 (3.2)	5.7 (3.1)		
Proteinuria (> 0.5 g), %	13.6	14.3	14		
Serum creatinine (mg/dL), mean (SD)	1.03 (0.43)	0.85 (0.28)	0.94 (0.37)		
Prednisone usage (mg), mean (SD)	20.3 (21.3)	18.7 (17.7)	19.4 (19.3)		

Supplemental Table 1: Demographics and clinical characteristics of the University of Michigan SLE observational cohort

Supplemental Table 2: Baseline peripheral blood populations in SLE trials stratified by ISM status

Parameter	Trial	ISM-Low	ISM-High	p-value
White Blood Cell				
Count (x10 ⁹ /L)	LUNAR Trial (N=80)	9.88 (2.6)	7.07 (3.6)	0.0012
	EXPLORER Trial (N=135)	9.04 (4.2)	6.47 (3.5)	0.0003
	ROSE Trial (N=238)	7.20 (2.8)	5.54 (2.5)	<0.0001
	Phase I Rontalizumab Trial (N=60)	6.37 (2.2)	5.26 (1.9)	0.0323
Lymphocytes				
(absolute counts				
x10 ⁹ /L)	LUNAR Trial (N=80)	2.46 (1.7)	1.37 (0.9)	0.0158
	EXPLORER Trial (N=135)	1.54 (1.0)	1.09 (0.6)	0.0154
	ROSE Trial (N=238)	1.76 (0.8)	1.15 (0.6)	<0.0001
	Phase I Rontalizumab Trial (N=60)	1.99 (0.8)	1.59 (0.6)	0.0608
Neutrophils		, ,	, ,	
(absolute counts				
x10°/L)	LUNAR Trial (N=80)	6.78 (1.8)	5.22 (3.0)	0.0107
	EXPLORER Trial (N=135)	7.02 (3.9)	4.99 (3.2)	0.0036
	ROSE Trial (N=238)	4.94 (2.6)	4.0 (2.2)	0.0087
	Phase I Rontalizumab Trial (N=60)	3.87 (1.6)	3.19 (1.5)	0.0781
Monocytes (absolute				
counts x10 ⁹ /L)	LUNAR Trial (N=80)	0.45 (0.19)	0.34 (0.2)	0.0149
	EXPLORER Trial (N=135)	0.34 (0.2)	0.30 (0.3)	0.0341
	ROSE Trial (N=238)	0.34 (0.1)	0.28 (0.2)	0.0007
	Phase I Rontalizumab Trial (N=60)	0.32 (0.1)	0.32 (0.1)	0.81
Eosinophils (absolute counts	, , , , , , , , , , , , , , , , , , ,			
$x10^{9}/L$	LUNAR Trial (N=80)	0.09 (0.08)	0.08 (0.07)	0.5662
	EXPLORER Trial (N=135)	0.08 (0.08)	0.04 (0.06)	0.001
	BOSE Trial (N=238)	0.10 (0.07)	0.07 (0.06)	<0.0001
	Phase I Rontalizumab Trial	0.10 (0.07)	0.07 (0.00)	NO.0001
	(N=60)	0.14 (0.11)	0.11 (0.09)	0.14

Data shown as mean (SD) for baseline values as defined by last pre-treatment value. P values calculated using the Wilcoxon test.

Supplemental Table 3: Multivariate linear model for the ISM

Intercept refers to the free term of the linear regression model: Predicted ISM = -4.045 -0.0013 * CD4 + 0.810 * (anti-ENA status = Positive) + 0.706*Ln (BAFF) + 0.205*Ln (anti-dsDNA) - 0.049 *(SLE duration)

Parameter	Coefficient Estimate	t statistic	p-value
Intercept	-4.045	-2.68	0.0079
CD4 counts (/µL)	-0.0013	-4.48	<0.0001
anti-ENA status (Positive vs. Negative)	0.810	3.71	0.0003
Ln(BAFF (IU/mL))	0.706	3.86	0.0002
Ln(anti-dsDNA (IU/mL))	0.205	3.10	0.0022
SLE duration (years)	-0.049	-3.16	0.0018

The t-statistic equals the coefficient estimate divided by the standard error used to derive a pvalue for each coefficient.

Supplemental Table 4: Reproducibility of ISM^a categorization over 36 weeks in the placebo arm of the ROSE study

ISM Value	Patients (n)	P _{rep} %
≤ -0.5	11	97
> –0.5, ≤ 1	6	81
> 1 to ≤ 2	11	85
> 2 to ≤ 3	19	98
≥ 3	14	100
All patients	61	94 (95% CI: 91 to 97)

All patients61(95% Cl; 91 to 97)CI, confidence interval; Prep, probability of reproducibility
a Data generated using a research-grade assay

Supplemental Table 5: Identity of Affymetrix probes for 128 Interferon-regulated gene cluster identified in the University of Michigan SLE cohort PBMC microarray analysis

Probe Set ID	Gene Symbol	Gene Title
1552623_at	HSH2D	hematopoietic SH2 domain containing
1555251_a_ at	OTOF	otoferlin
1556643_at	LOC1005 07535	uncharacterized LOC100507535
1568592_at	TRIM69	tripartite motif containing 69
200814_at	PSME1	proteasome (prosome, macropain) activator subunit 1 (PA28 alpha)
200887_s_at	STAT1	signal transducer and activator of transcription 1, 91kDa
200923_at	LGALS3B P	lectin, galactoside-binding, soluble, 3 binding protein
200986_at	SERPING 1	serpin peptidase inhibitor, clade G (C1 inhibitor), member 1
201315_x_at	IFITM2	interferon induced transmembrane protein 2
201641_at	BST2	bone marrow stromal cell antigen 2
201649_at	UBE2L6	ubiquitin-conjugating enzyme E2L 6
201762_s_at	PSME2	proteasome (prosome, macropain) activator subunit 2 (PA28 beta)
201786_s_at	ADAR	adenosine deaminase, RNA-specific
202086_at	MX1	myxovirus (influenza virus) resistance 1, interferon- inducible protein p78 (mouse)
202145_at	LY6E	lymphocyte antigen 6 complex, locus E
202307_s_at	TAP1	transporter 1, ATP-binding cassette, sub-family B (MDR/TAP)
202411_at	IFI27	interferon, alpha-inducible protein 27
202446_s_at	PLSCR1	phospholipid scramblase 1
202688_at	TNFSF10	tumor necrosis factor (ligand) superfamily, member 10
202869_at	OAS1	2'-5'-oligoadenylate synthetase 1, 40/46kDa
203153_at	IFIT1	interferon-induced protein with tetratricopeptide repeats 1
203236_s_at	LGALS9	lectin, galactoside-binding, soluble, 9
203258_at	DRAP1	DR1-associated protein 1 (negative cofactor 2 alpha)
203595_s_at	IFIT5	interferon-induced protein with tetratricopeptide repeats 5
203773_x_at	BLVRA	biliverdin reductase A
203882_at	IRF9	interferon regulatory factor 9
203964_at	NMI	N-myc (and STAT) interactor
204279_at	PSMB9	proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional peptidase 2)
204415_at	IFI6	interferon, alpha-inducible protein 6
204439_at	IFI44L	interferon-induced protein 44-like
204698_at	ISG20	interferon stimulated exonuclease gene 20kDa

204804_at	TRIM21	tripartite motif containing 21
204858_s_at	TYMP	thymidine phosphorylase
204972_at	OAS2	2'-5'-oligoadenylate synthetase 2, 69/71kDa
204994_at	MX2	myxovirus (influenza virus) resistance 2 (mouse)
205098_at	CCR1	chemokine (C-C motif) receptor 1
205126_at	VRK2	vaccinia related kinase 2
205241_at	SCO2	SCO cytochrome oxidase deficient homolog 2 (yeast)
205483_s_at	ISG15	ISG15 ubiquitin-like modifier
205569_at	LAMP3	lysosomal-associated membrane protein 3
205660_at	OASL	2'-5'-oligoadenylate synthetase-like
206513_at	AIM2	absent in melanoma 2
207777_s_at	SP140	SP140 nuclear body protein
208436_s_at	IRF7	interferon regulatory factor 7
208966_x_at	IFI16	interferon, gamma-inducible protein 16
209417_s_at	IFI35	interferon-induced protein 35
209546_s_at	APOL1	apolipoprotein L, 1
209568_s_at	RGL1	ral guanine nucleotide dissociation stimulator-like 1
209593_s_at	TOR1B	torsin family 1, member B (torsin B)
209762_x_at	SP110	SP110 nuclear body protein
210873_x_at	APOBEC3 A	apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3A
211012_s_at	PML	promyelocytic leukemia
211012_s_at 211138_s_at	PML KMO	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase)
211012_s_at 211138_s_at 211267_at	PML KMO HESX1	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1
211012_s_at 211138_s_at 211267_at 212203_x_at	PML KMO HESX1 IFITM3	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at	PML KMO HESX1 IFITM3 FTSJD2	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at 213293_s_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRIM22	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at 213293_s_at 213294_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRIM22 EIF2AK2	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at 213293_s_at 213294_at 213361_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRANK1 TRIM22 EIF2AK2 TDRD7	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2 tudor domain containing 7
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at 213293_s_at 213294_at 213361_at 214022_s_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRANK1 TRIM22 EIF2AK2 TDRD7 IFITM1	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2 tudor domain containing 7 interferon induced transmembrane protein 1
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at 213293_s_at 213294_at 213361_at 214022_s_at 214453_s_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRANK1 TRIM22 EIF2AK2 TDRD7 IFITM1 IFI44	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2 tudor domain containing 7 interferon induced transmembrane protein 1 interferon-induced protein 44
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at 213293_s_at 213294_at 213361_at 214022_s_at 214453_s_at 214511_x_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRIM22 EIF2AK2 TDRD7 IFITM1 IFI44 FCGR1B	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2 tudor domain containing 7 interferon induced transmembrane protein 1 interferon-induced protein 44 Fc fragment of IgG, high affinity Ib, receptor (CD64)
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at 213293_s_at 213294_at 213294_at 214022_s_at 214453_s_at 214511_x_at 217933_s_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRIM22 EIF2AK2 TDRD7 IFITM1 IFI44 FCGR1B LAP3	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2 tudor domain containing 7 interferon induced transmembrane protein 1 interferon-induced protein 44 Fc fragment of IgG, high affinity lb, receptor (CD64) leucine aminopeptidase 3
211012_s_at 211138_s_at 211267_at 212203_x_at 212203_x_at 212657_s_at 213051_at 213261_at 213293_s_at 213294_at 213361_at 214022_s_at 214453_s_at 214511_x_at 217933_s_at 217986_s_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRIM22 EIF2AK2 TDRD7 IFITM1 IFI44 FCGR1B LAP3 BAZ1A	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2 tudor domain containing 7 interferon induced transmembrane protein 1 interferon-induced protein 44 Fc fragment of IgG, high affinity Ib, receptor (CD64) leucine aminopeptidase 3 bromodomain adjacent to zinc finger domain, 1A
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at 213294_at 213294_at 213051_at 214022_s_at 214022_s_at 214511_x_at 217933_s_at 217986_s_at 218076_s_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRIM22 EIF2AK2 TDRD7 IFITM1 IFI44 FCGR1B LAP3 BAZ1A ARHGAP 17	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2 tudor domain containing 7 interferon induced transmembrane protein 1 interferon-induced protein 44 Fc fragment of IgG, high affinity lb, receptor (CD64) leucine aminopeptidase 3 bromodomain adjacent to zinc finger domain, 1A Rho GTPase activating protein 17
211012_s_at 211138_s_at 211267_at 212203_x_at 212203_x_at 212657_s_at 213051_at 213261_at 213293_s_at 213294_at 213361_at 214022_s_at 214453_s_at 214511_x_at 217933_s_at 217986_s_at 218076_s_at 218231_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRIM22 EIF2AK2 TDRD7 IFITM1 IFI44 FCGR1B LAP3 BAZ1A ARHGAP 17 NAGK	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2 tudor domain containing 7 interferon induced transmembrane protein 1 interferon-induced protein 44 Fc fragment of IgG, high affinity Ib, receptor (CD64) leucine aminopeptidase 3 bromodomain adjacent to zinc finger domain, 1A Rho GTPase activating protein 17 N-acetylglucosamine kinase
211012_s_at 211138_s_at 211267_at 212203_x_at 212380_at 212657_s_at 213051_at 213261_at 213294_at 213294_at 213294_at 214022_s_at 214453_s_at 214511_x_at 217933_s_at 217986_s_at 218076_s_at 218231_at 218400_at	PML KMO HESX1 IFITM3 FTSJD2 IL1RN ZC3HAV1 TRANK1 TRIM22 EIF2AK2 TDRD7 IFITM1 IFI44 FCGR1B LAP3 BAZ1A ARHGAP 17 NAGK OAS3	promyelocytic leukemia kynurenine 3-monooxygenase (kynurenine 3-hydroxylase) HESX homeobox 1 interferon induced transmembrane protein 3 FtsJ methyltransferase domain containing 2 interleukin 1 receptor antagonist zinc finger CCCH-type, antiviral 1 tetratricopeptide repeat and ankyrin repeat containing 1 tripartite motif containing 22 eukaryotic translation initiation factor 2-alpha kinase 2 tudor domain containing 7 interferon induced transmembrane protein 1 interferon-induced protein 44 Fc fragment of IgG, high affinity lb, receptor (CD64) leucine aminopeptidase 3 bromodomain adjacent to zinc finger domain, 1A Rho GTPase activating protein 17 N-acetylglucosamine kinase 2'-5'-oligoadenylate synthetase 3, 100kDa

218543_s_at	PARP12	poly (ADP-ribose) polymerase family, member 12
218943_s_at	DDX58	DEAD (Asp-Glu-Ala-Asp) box polypeptide 58
218986_s_at	DDX60	DEAD (Asp-Glu-Ala-Asp) box polypeptide 60
219014_at	PLAC8	placenta-specific 8
219062_s_at	ZCCHC2	zinc finger, CCHC domain containing 2
219209_at	IFIH1	interferon induced with helicase C domain 1
219211_at	USP18	ubiquitin specific peptidase 18
219352_at	HERC6	HECT and RLD domain containing E3 ubiquitin protein ligase family member 6
219356_s_at	CHMP5	charged multivesicular body protein 5
219364_at	DHX58	DEXH (Asp-Glu-X-His) box polypeptide 58
219371_s_at	KLF2	Kruppel-like factor 2 (lung)
219519_s_at	SIGLEC1	sialic acid binding Ig-like lectin 1, sialoadhesin
219684_at	RTP4	receptor (chemosensory) transporter protein 4
219863_at	HERC5	HECT and RLD domain containing E3 ubiquitin protein ligase 5
220146_at	TLR7	toll-like receptor 7
221680_s_at	ETV7	ets variant 7
221816_s_at	PHF11	PHD finger protein 11
221827_at	RBCK1	RanBP-type and C3HC4-type zinc finger containing 1
222154_s_at	SPATS2L	spermatogenesis associated, serine-rich 2-like
222986_s_at	SHISA5	shisa homolog 5 (Xenopus laevis)
223220_s_at	PARP9	poly (ADP-ribose) polymerase family, member 9
223322_at	RASSF5	Ras association (RalGDS/AF-6) domain family member 5
223501_at	TNFSF13 B	tumor necrosis factor (ligand) superfamily, member 13b
223599_at	TRIM6	tripartite motif containing 6
224701_at	PARP14	poly (ADP-ribose) polymerase family, member 14
225076_s_at	ZNFX1	zinc finger, NFX1-type containing 1
225291_at	PNPT1	polyribonucleotide nucleotidyltransferase 1
225344_at	NCOA7	nuclear receptor coactivator 7
225415_at	DTX3L	deltex 3-like (Drosophila)
225636_at	STAT2	signal transducer and activator of transcription 2, 113kDa
226603_at	SAMD9L	sterile alpha motif domain containing 9-like
226702_at	CMPK2	cytidine monophosphate (UMP-CMP) kinase 2, mitochondrial
226748_at	LYSMD2	LysM, putative peptidoglycan-binding, domain containing 2
226757_at	IFIT2	interferon-induced protein with tetratricopeptide repeats 2
227609_at	EPSTI1	epithelial stromal interaction 1 (breast)
228152_s_at	DDX60L	DEAD (Asp-Glu-Ala-Asp) box polypeptide 60-like
228230_at	PRIC285	peroxisomal proliferator-activated receptor A interacting complex 285
228439_at	BATF2	basic leucine zipper transcription factor, ATF-like 2

228531_at	SAMD9	sterile alpha motif domain containing 9
228617_at	XAF1	XIAP associated factor 1
229350_x_at	PARP10	poly (ADP-ribose) polymerase family, member 10
229450_at	IFIT3	interferon-induced protein with tetratricopeptide repeats 3
230166_at	KIAA1958	KIAA1958
230405_at	C5orf56	chromosome 5 open reading frame 56
231455_at	LINC0048 7	long intergenic non-protein coding RNA 487
231577_s_at	GBP1	guanylate binding protein 1, interferon-inducible
231747_at	CYSLTR1	cysteinyl leukotriene receptor 1
231769_at	FBXO6	F-box protein 6
232222_at	C18orf49	chromosome 18 open reading frame 49
236285_at	KLHDC7B	kelch domain containing 7B
238327_at	ODF3B	outer dense fiber of sperm tails 3B
238439_at	ANKRD22	ankyrin repeat domain 22
242020_s_at	ZBP1	Z-DNA binding protein 1
242625_at	RSAD2	radical S-adenosyl methionine domain containing 2
32069_at	N4BP1	NEDD4 binding protein 1
35254_at	TRAFD1	TRAF-type zinc finger domain containing 1
38269_at	PRKD2	protein kinase D2