

Background Past studies suggest that males with lupus nephritis (LN) may have increased rates of end-stage renal disease (ESRD) and mortality compared to females. However studies included few males and were focused on biological differences, not health-care use. In a nationwide cohort of SLE patients with incident LN, we investigated LN-related outcomes and utilisation by sex.

Materials and methods We used the Medicaid Analytic eXtract (MAX) with nationwide billing claims to identify individuals 5–65 years with LN (2000–2004) using a validated algorithm (PPV 80%) and required 12 months without any LN codes to define incident cases. MAX data were linked to the U.S. Renal Data System (USRDS) 2000–2006 to identify ESRD onset. Mortality was determined using National and Social Security Death Index Files (2000–2006). We assessed sex-specific incidence rates and adjusted incidence rate ratios (IRRs) for healthcare utilisation, medications, preventive care and renal biopsies using Poisson regression. We used Fine and Grey proportional hazard models to compare the subdistribution hazard ratios (HRsd) of ESRD by sex accounting for the competing risk of death, and Cox models to compare hazard ratios (HR) of death, adjusted for age and race/ethnicity.

Results Of 2576 patients with incident LN, 230 (9%) were male. Mean follow-up was 2.8 (SD 1.5) years for both sexes. Mean age was 30 (SD 16) years among males and 34 (SD 14) years among females ($p < 0.001$). 31% of males and 36% of females underwent renal biopsy ($p = 0.06$). Other than azathioprine use, which was more frequent among females ($p = 0.02$), were no differences in medications or preventive care. Adjusted rates of outpatient and emergency department (ED) visits were lower for males compared to females (IRR 0.81, 95% CI: 0.68–0.98 and 0.88, 95% CI: 0.79–0.99, respectively); hospitalizations were comparable. The five-year cumulative incidence of ESRD was 13% and the HRsd of ESRD for males compared to females was 0.86 (95% CI: 0.53–1.38) in the first two years and 2.21 (95% CI: 1.2–4.1) in the subsequent two years. The five-year cumulative incidence of death was 15% with no difference in HR by sex.

Conclusions In this incident LN Medicaid cohort, we found high rates of ESRD and mortality overall, with no differences in ESRD by sex in the first two years but more than twice the risk among males thereafter. Males had lower rates of outpatient and ED visits compared to females. Further studies are needed to understand the relationship between utilisation and long-term outcomes.

Acknowledgements Drs. Feldman and Broder contributed equally.

CE-05 A LONGITUDINAL ANALYSIS OF OUTCOMES OF LUPUS NEPHRITIS IN AN INTERNATIONAL INCEPTION COHORT USING A MULTISTATE MODEL APPROACH

John G Hanly*. Dalhousie University and Nova Scotia Health Authority, Halifax, NS, Canada

10.1136/lupus-2016-000179.84

Background Patients with lupus nephritis (LN) may have improvement or deterioration in renal status over time. To capture bidirectional change we used a reversible multistate Markov model to study transitions in glomerular filtration rate (GFR) and proteinuria (PrU) in a prospective, international, inception cohort of SLE patients receiving standard of care.

Materials and methods Patients were evaluated at enrolment and annually. GFR states were defined: state 1 (eGFR: >60 ml/

min); state 2 (eGFR: 30–60 ml/min); and state 3 (eGFR: <30 ml/min). Similarly, PrU states were defined: state 1 (ePrU: <0.25 gr/day); state 2 (ePrU: 0.25–3.0 gr/day); and state 3 (ePrU: >3.0 gr/day). Multistate models provided estimates of relative transition rates and state occupancy probabilities.

Results Of 1,826 SLE patients, 89% were female, 49.2% Caucasian with mean \pm SD age 35.1 ± 13.3 years. The mean disease duration at enrollment was 0.5 ± 0.3 years and follow-up was 4.6 ± 3.4 years. LN occurred in 700/1,826 (38.3%) patients. The likelihood of improvement in eGFR and ePrU (states 2 \rightarrow 1 and 3 \rightarrow 2) was greater than deterioration (states 1 \rightarrow 2 and 2 \rightarrow 3). After 5 years, the estimated transition to ESRD was 62% of patients initially in eGFR state 3 but only 11% from ePrU state 3. The probability of remaining in initial eGFR states 1, 2 and 3 was 85%, 11%, 3% and for ePrU was 62%, 29%, 4%. Male sex ($p = 0.04$) predicted improvement in eGFR states and older age ($p < 0.001$), race/ethnicity ($p < 0.001$), higher ePrU state ($p < 0.001$), higher renal biopsy chronicity score ($p = 0.013$) and baseline anticardiolipin antibodies ($p = 0.039$) predicted deterioration. For ePrU, race/ethnicity ($p = 0.009$), higher eGFR state ($p = 0.011$) and higher renal biopsy chronicity score ($p = 0.015$) predicted deterioration. Positive lupus anticoagulant ($p = 0.006$) and ISN/RPN class V nephritis ($p = 0.013$) were associated with lower improvement rates.

Conclusions Multistate modelling in patients with LN generates probability estimates of transitions between disease states that reflect improvement or deterioration in renal outcomes. This approach identifies predictors of change in renal status and can inform clinical trial design by identifying outcomes that new therapeutic interventions for LN should meet or exceed.

Acknowledgements Presented on behalf of the Systemic Lupus International Collaborating Clinics (SLICC)

CE-06 URINARY NEUTROPHIL GELATINASE – ASSOCIATED LIPOCALIN AND MONOCYTE CHEMOATTRACTANT PROTEIN-1 AS BIOMARKERS FOR LUPUS NEPHRITIS IN COLOMBIAN SLE PATIENTS

^{1,2}Jose A Gómez-Puerta*, ¹Blanca L Ortiz Reyes, ¹Tomás Urrego, ^{2,3}Adriana L Vanegas, ^{2,3}Carlos H Muñoz, ²Mauricio Restrepo, ²Luis A González, ^{1,2}Gloria Vásquez. ¹Grupo de Inmunología Celular e Inmunogenética, Universidad de Antioquia, Medellín; ²Sección de Reumatología, Universidad de Antioquia; ³Hospital Universitario de San Vicente Fundación, Medellín, Colombia

10.1136/lupus-2016-000179.85

Background Some previous studies in Caucasian, Asian, and African-American patients have shown that urine levels of Neutrophil Gelatinase-Associated Lipocalin (uNGAL) and Monocyte Chemoattractant Protein-1 (uMCP-1) were significantly greater in patients with LN. However, information in Mestizo and Afro-Latin American patients is very limited. Our aim was to evaluate diagnostic value of uNGAL and uMCP-1 as potential markers for the diagnosis of LN in Colombian SLE patients

Materials and methods We examined urinary levels of NGAL and MCP-1 in 93 consecutive SLE patients (ACR criteria 1982) from Hospital San Vicente Fundación, at Medellín, Colombia. uNGAL and uMCP-1 were measured by ELISA techniques (R&D system, Minneapolis, USA). In addition, serum Anti-C1q antibodies were measured (Inova, San Diego, USA). Several clinical and serological features were analysed as well as disease activity (SLE-DAI). Mann-Whitney tests were used to compare data and Spearman's rho for correlations. Additionally, ROC curves relating the specificity and sensitivity profiles of the 2 biomarkers were done.