

after being treated with rituximab (500mg d 0) plus bortezomib (2.5mg d 1, 4, 8, 11), the immune status of the patient was improved gradually and all indexes returned to normal ranges. Half a year later, rituximab (100mg d 0) plus bortezomib (2.5mg d 1, 4, 8, 11) was applied. Ultimately, the patient was in remission without recurrence until now. Additionally, we summarized the available literature and explained the possible mechanism. Rituximab combined with bortezomib was more effective for depleting B cells and plasmocytes, which was accompanied by decreases in antibodies. Finally, although there were no adverse effects in this patient, we pointed out that rituximab plus bortezomib might cause infection and peripheral neuropathy. Therefore, monitoring safety is required.

**Conclusions** Our case shows that rituximab combined with bortezomib is effective in treating SLE patients with refractory hemolytic anemia, but safety should be monitored.

LP-191

#### BASELINE PROFILE OF SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS ON TREATMENT WITH BELIMUMAB OF A SPANISH MULTICENTER COHORT

<sup>1,19</sup>Irene Altobás-González\*, <sup>1,19</sup>José María Pego-Reigosa, <sup>1,19</sup>Coral Mouriño-Rodríguez, <sup>1,19</sup>Norman Jiménez-Otero, <sup>2</sup>Andrea Hernández-Martín, <sup>3</sup>Judit Font-Urgelles, <sup>3</sup>Yvette Casafont-Sole, <sup>4</sup>José Andrés Roman-Ivorra, <sup>4</sup>Marta De La Rubia-Navarro, <sup>5</sup>María Galindo-Izquierdo, <sup>6</sup>Tarek Salman-Montes, <sup>7</sup>Javier Narváez, <sup>7</sup>Paola Vidal-Montal, <sup>8</sup>María Jesús García-Villanueva, <sup>8</sup>Sandra Garrote-Corral, <sup>8</sup>María Ángeles Blázquez-Cañamero, <sup>9</sup>Carlos Marras-Fernández, <sup>9</sup>María Piqueras-García, <sup>10</sup>Julia Martínez-Barrio, <sup>10</sup>Marina Sánchez-Lucas, <sup>11</sup>Josefina Cortés-Hernández, <sup>11</sup>Eleonora Penzo, <sup>12</sup>Jaime Calvo-Alen, <sup>12</sup>Juan Ramón De Dios Jiménez De Aberásturi, <sup>12</sup>Belén Álvarez-Rodríguez, <sup>12</sup>Margarida Vasques-Rocha, <sup>13</sup>Eva Tomero, <sup>14</sup>Raúl Menor-Almagro, <sup>14</sup>Myriam Gandía-Martínez, <sup>15</sup>José A Gómez-Puerta, <sup>16</sup>Consuelo Ramos-Giráldez, <sup>16</sup>Carmen Traperero-Pérez, <sup>17</sup>Elvira Díez-Álvarez, <sup>17</sup>Clara Moriano, <sup>18</sup>Alejandro Muñoz-Jiménez, <sup>2</sup>Íñigo Rúa-Figueroa. <sup>1</sup>Rheumatology, Complejo Hospitalario universitario de Vigo, Spain; <sup>2</sup>Rheumatology, Hospital Universitario De Gran Canaria Dr. Negrín, Spain; <sup>3</sup>Rheumatology, Hospital Universitario Germans Trias i Pujol, Spain; <sup>4</sup>Rheumatology, Hospital Universitario y Politécnico De La Fe, Spain; <sup>5</sup>Rheumatology, Hospital 12 De Octubre, Spain; <sup>6</sup>Rheumatology, Hospital Del Mar, Spain; <sup>7</sup>Rheumatology, Hospital Universitario De Bellvitge, Spain; <sup>8</sup>Rheumatology, Hospital Universitario Ramón y Cajal, Spain; <sup>9</sup>Rheumatology, Hospital Virgen Ae La Arrixaca De Murcia, Spain; <sup>10</sup>Rheumatology, Hospital General Universitario Gregorio Marañón, Spain; <sup>11</sup>Rheumatology, Hospital Universitario Valle d' Hebrón, Spain; <sup>12</sup>Rheumatology, Hospital Universitario Araba, Spain; <sup>13</sup>Rheumatology, Hospital Universitario De La Princesa, Spain; <sup>14</sup>Rheumatology, Hospital Universitario De Jerez, Spain; <sup>15</sup>Rheumatology, Hospital Clinic De Barcelona, Spain; <sup>16</sup>Rheumatology, Hospital Universitario Nuestra Señora de Valme, Spain; <sup>17</sup>Rheumatology, Hospital Universitario De León, Spain; <sup>18</sup>Rheumatology, Hospital universitario Virgen Del Rocío, Spain; <sup>19</sup>Rheumatology, IRIDIS Group, Spain

10.1136/lupus-2023-KCR.265

**Background** Baseline profile of systemic lupus erythematosus patients on treatment with Belimumab of a Spanish multicenter cohort.

**Methods** Multicenter retrospective and longitudinal cohort study. Data was collected at baseline, 6, 12 months and in the last visit available. Different periods (2010–2015/2016–2021) were compared.

**Results** 324 patients (91% female) were enrolled. Mean ( $\pm$ SD) age at diagnosis: 31.8 years ( $\pm$ 11.9); mean disease duration of 8.7 years ( $\pm$ 9.07). At baseline, mean SLEDAI-2K score was 10.4 ( $\pm$ 5.25); 152 (47.5%) patients had damage with a mean SDI score of 0.83 ( $\pm$ 1.2). 289 patients (89.2%) had received DMARDs before BLM: cDMARDs in 282 (87%) and bDMARDs in 74 patients (22.8%); 164 (51.9%) had received more than one cDMARDs, methotrexate being the most frequently used (44.4%) and Rituximab the most frequent

bDMARDs used (80%). Antimalarials were used in 83.2% and glucocorticoids (GC) in 91.2%, with a mean dose of 12.3 mg/day. 67.9% of patients were receiving more than 5 mg/day and 58.4% more than 7.5 mg/day of prednisone. BLM was used in monotherapy in 30.5% of subjects. Reasons of prescription: disease activity in 95% of patients and/or as a GC sparing agent in 59%. Only a few patients received BLM just for maintenance (4/322) or save GC (8/322). At baseline, 6 patients (1.9%) were in DORIS-21-remission and LLDAS. The main reasons of prescription for ongoing activity were arthritis (65.4%), cutaneous (40.7%) or both (81%). No differences were observed in prescription reasons between periods.

**Conclusions** Belimumab was mainly prescribed after the use of other DMARDs and more than 50% had received at least 2 DMARDs and were receiving GC at medium doses. Most patients received BLM due to active disease and/or as GC sparing agent. Activity in articular and cutaneous domains were the main reasons of indication. No changes in prescription habits were identified.

LP-195

#### DESIGN OF A PHASE 2 STUDY EVALUATING THE EFFICACY AND SAFETY OF NIPOCALIMAB IN ADULT PATIENTS WITH ACTIVE SYSTEMIC LUPUS ERYTHEMATOSUS

<sup>1</sup>Fang Liu-walsh, <sup>2</sup>Bart van Hartingsveldt, <sup>1</sup>Qing Zuraw, <sup>1</sup>Robert Hoffman, <sup>1</sup>Terence Rooney, <sup>1</sup>Sheng Gao, <sup>3</sup>Youngja Lee\*, <sup>1</sup>Robert Gordon, <sup>1</sup>Jocelyn Leu, <sup>1</sup>Cesar Calderon, <sup>4</sup>Federico Zazzetti, <sup>1</sup>Anne Stevens, <sup>1</sup>George Vratsanos. <sup>1</sup>Immunology, Janssen Research and Development, USA; <sup>2</sup>Immunology, Janssen Biologics Europe, Netherlands; <sup>3</sup>Immunology, Janssen Asia Pacific, Republic of Korea; <sup>4</sup>Immunology, Janssen-Cilag Argentina, Argentina

10.1136/lupus-2023-KCR.266

**Background** Systemic lupus erythematosus (SLE) is a chronic, complex autoimmune disease characterized by pathogenic auto-antibodies and tissue damage to multiple organ systems. Nipocalimab is a novel high affinity, fully human, aglycosylated, effectorless IgG1 monoclonal antibody that selectively blocks the neonatal Fc receptor (FcRn). Nipocalimab has demonstrated rapid and durable serum IgG and pathogenic autoantibody reductions (NCT02828046; NCT03896295). Here we describe the protocol of a Phase 2 study evaluating the efficacy and safety of nipocalimab in patients with active SLE (NCT04882878).

**Methods** Phase 2, multicenter, randomized, placebo-controlled, double-blind, parallel-group study enrolling adults with active, autoantibody-positive SLE with an inadequate response to one or more standard of care treatments. The study consists of a  $\leq$ 6-week screening period, a 52-week double-blind treatment period, and a 6-week follow-up period (figure 1). A target of approximately 225 participants will be enrolled. Participants will be randomized in a 1:1:1 ratio to receive nipocalimab dose 1, dose 2 or placebo intravenously every 2 weeks through Week 50.

**Results** The primary efficacy endpoint is the percentage of participants achieving an SLE Responder Index (SRI)-4 composite response at Week 24. Secondary efficacy endpoints assessed at Week 24 include the percentage of participants achieving:  $\geq$ 50% reduction in Cutaneous Lupus Erythematosus Disease Area and Severity Index Activity score (CLASI),  $\geq$ 50% reduction in active joints,  $\geq$ 4 points improvement in SLE Disease Activity Index 2000 (SLEDAI 2K), and British Isles Lupus Assessment Group Composite Lupus Assessment response (BICLA); time to first disease flare; and reduction