

for IgG in humans, we included transgenic mice expressing FcγRIIA to our *in vivo* mechanistic investigations. Platelet activation and mitochondrial release were monitored upon intravenous injection of synthetic ICs in wild type (WT) and transgenic (Tg) FcγRIIA mice.

Results We found that on activation, platelets relocate their mitochondria toward the cell membrane and then release respiratory-competent mitochondria into the extracellular milieu, both as free organelles and encapsulated within vesicles called microparticles (MP). Extracellular mitochondria were internalised by bystander leukocytes and, importantly, induced leukocytes rolling and adhesion to the blood vessel wall, suggesting that neutrophils and/or endothelial cells are activated by extracellular mitochondria. Activated platelets were more abundant in SLE patients than control subjects, and were associated with IgG. Extracellular mitochondria, both encapsulated in platelet MPs or naked, were also observed in blood circulation in SLE, and were frequently associated with IgG. Mechanistically, ICs present in blood induce profound cell activation, which is dependent on platelet FcγRIIA and its signalling cascade.

Conclusions Platelets represent an important source of mitochondria, which release in blood upon stimulation of FcγRIIA, might promote systemic inflammation in SLE. Whether the blockade of FcγRIIA might represent an attractive avenue in SLE research, and whether platelet activation markers and extracellular mitochondria might be utilised as potential biomarkers for the stratification of lupus patients needs to be further considered.

Acknowledgements This study was supported by the Canadian Institutes of Health Research (to PRF and EB). EB is recipient of an award from the Canadian Institutes of Health Research.

11-12 CSF-1 AND IL-34: DISTINCT POTENTIAL BIOMARKERS FOR LUPUS

¹Julia Menke, ¹Andreas Schwarting, ²Vicki Rubin Kelley*. ¹Department of Nephrology and Rheumatology, Johannes-Gutenberg University Mainz, German; ²Department of Medicine, Brigham and Women's Hospital, Boston, MA

10.1136/lupus-2016-000179.42

Background A noninvasive means to predict the onset and recurrence of lupus is needed to optimise and individualise treatment. Macrophages (M ϕ) are prominent in inflamed tissues targeted for destruction in SLE. We hypothesised that the principal molecules required for M ϕ survival and proliferation are biomarkers for SLE. CSF-1 and IL-34 are promising candidates as both (i) bind to cFMS expressed by M ϕ , thereby promoting M ϕ survival and proliferation and (ii) promote destructive inflammation. However, IL-34 and CSF-1 have differing functions, which may be related to IL-34, not CSF-1, binding to a second receptor and distinct spatial temporal expressions.

Materials and methods We analysed serum and urine CSF-1 and IL-34 levels in SLE patients with nephritis (LN), arthritis (LA), cutaneous and serositis compared with healthy controls in two large cohorts (ELISA). While serum and urine CSF-1 expression is elevated in each manifestation, CSF-1 is notably higher in LN. In contrast, serum IL-34 expression is dramatically higher in LA, not LN. Thus, we probed for CSF-1 and IL-34 expression in LN (kidney) and LA (synovium). Moreover, we longitudinally tracked serum CSF-1 and IL-34 prior to LN (biopsy proven), with disease activity including flares and during LA in comparison to disease activity

Results

- **LN.** CSF-1 and IL-34 are expressed in the same and different renal tubular epithelial cells in LN. Elevated serum or urine CSF-1, not IL-34, levels correlate with increasing intra-renal CSF-1 expression and histopathology index. Longitudinally tracking serum CSF-1, not IL-34, levels heralds the initial onset of nephritis and a rise in serum or urine CSF-1 predicts LN recurrences before clinical evidence of renal dysfunction and conventional serologic measures.
- **LA.** IL-34, not CSF-1, expression is higher in synovial fluid and synovium in LA compared to osteoarthritis and healthy controls and correlates with magnitude of intra-synovial leukocytes. Moreover, intra-synovial IL-34 expression is similar in LA and rheumatoid arthritis. Longitudinally monitoring serum IL-34, not CSF-1, levels track with clinical disease activity in LA and RA.

Conclusions Serial monitoring a rise in serum or urine CSF-1, not IL-34, in SLE reflects renal histopathology and clinical disease activity and the onset and recurrences of LN more accurately than conventional laboratory measures. While serial monitoring a rise in serum IL-34, not CSF-1, reflects clinical disease activity in LA. Thus, CSF-1 and IL-34 are inexpensive and accurate potential biomarkers for LN and LA, respectively.

11-13 SUPPRESSION OF SYSTEMIC AUTOIMMUNITY BY THE INNATE IMMUNE ADAPTOR STING

¹Shruti Sharma*, ²Allison M Campbell, ¹Jennie Chan, ¹Stefan A Schattgen, ¹Gregory M Orlowski, ³Ribhu Nayar, ¹Annie H Huyler, ⁴Kerstin Nündel, ⁵Chandra Mohan, ³Leslie J Berg, ⁶Mark J Shlomchik, ⁴Ann Marshak-Rothstein, ^{†1}Katherine A Fitzgerald. ¹Program in Innate Immunity, Division of Infectious Diseases, Department of Medicine, University of Massachusetts Medical School, Worcester, MA-01605; ²Department of Laboratory Medicine, Yale University School of Medicine, New Haven, CT 06519; ³Department of Pathology, University of Massachusetts Medical School, Worcester, MA - 01605; ⁴Division of Rheumatology, Department of Medicine, University of Massachusetts Medical School, Worcester, MA 01605; ⁵Department of Immunology and Medicine, University of Texas Southwestern Medical Centre, Dallas, TX 75390; ⁶Department of Immunology, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261; ⁷Centre of Molecular Inflammation Research, Norwegian University of Science and Technology, 7489 Trondheim, Norway
(* and † contributed equally)

10.1136/lupus-2016-000179.43

Background Cytosolic DNA-sensing pathways that signal via the adaptor Stimulator of Interferon Genes (STING) mediate immunity to pathogens and have also been known to promote autoimmune pathology in DNaseII/III-deficient mice. However, the role of these pathways in systemic models of autoimmunity is unexplored. We hypothesised that cytosolic DNA sensing pathways contribute to the pathogenesis of autoimmune disease. Surprisingly, we report here that STING *potently* suppresses inflammation in several models of systemic lupus erythematosus (SLE).

Materials and methods A controlled F2 intercross between heterozygote STING[±] lpr[±] littermates generated STING-deficient lupus-prone mice homozygous for deficiency in Fas as well as STING (STING/lpr, n ≥10) or wild type for STING (WT/lpr, n ≥10). Mice were analysed at 16 wk of age. A similar F2 cross was set up for IRF3^{-/-} and MRL/lpr mice as well as STING^{-/-} and C57Bl/6^{lpr/lpr} and analysed as above for STING/lpr mice (n ≥10 per group). C57Bl/6, cGAS^{-/-}, Unc93b^{3d/3d}, and STING^{-/-} mice were injected i.p. with TMPD and evaluated at day 14 and 6 months post injection.