Abstract 255 Table 2 Predictors of remission/LLDAS. Multivariable model.

Predictor	Hazard ratio (95% CI)	p value
Prednisone dose (higher dose before baseline)		
None	Ref.	
Low	1.745 (0.814 – 3.742)	0.1527
Medium	1.559 (1.081 – 2.247)	0.0174
High	0.891 (0.668 – 1.188)	0.4315
Very high	1.141 (0.802 – 1.623)	0.4622
Mucocutaneous involvement	0.652 (0.474 – 0.898)	0.0089
Renal involvement	0.686 (0.532 – 0.884)	0.0036
SLEDAI at baseline	0.979 (0.962 – 0.996)	0.0170
Prednisone dose: Low dose: < 7.5 mg/d, medium dose : 7.5-15 mg/d, high dose : 15-60 mg/d, very high dose : ≥ 60 mg/d		

associated with a reduced risk of remission and remission/LLDAS; lower socioeconomic status was associated with a reduced risk of remission. A medium prednisone dose was associated with an increased risk of remission/LLDAS.

256

THE EFFECT OF ADDING CURCUMIN ON VITAMIN D3 SUPPLEMENTATION ON THE DISEASE ACTIVITIES AND FATIGUE DEGREE IN SLE PATIENTS WITH HYPOVITAMIN D

¹CS Wahono*, ¹ZD Wahyuni, ¹CD Setyorini, ¹I Saveria, ²K Handono, ¹H Kalim. ¹Universitas Brawijaya/Saiful Anwar Hospital, Internal Medicine, Malang, Indonesia; ²Universitas Brawijaya/Saiful Anwar Hospital, Clinical Pathology, Malang, Indonesia

10.1136/lupus-2017-000215.256

Background and aims Vitamin D has important roles in the regulation of the immune system in Lupus. Seventy percent of lupus patients in Indonesia are experienced hypovitamin D. Curcumin is a natural VDR ligand and has synergic effect with vitamin D. This study was aimed to determine the effect of adding curcumin on vitamin D supplementation on the degree of disease activity and degree of fatigue, in SLE patients with hypovitamin D.

Methods This was a randomised controlled trial, double blind study. Forty SLE patients with hypovitamin D were studied, that randomized into two groups: 20 patients (supplementation group) received vitamin D (cholecalciferol 1200 IU daily) with curcumin 20 mg (three times daily); and 20 patients (placebo group) was given vitamin D (cholecalciferol 1200 IU daily) and placebo (3 times daily), for 3 months. Disease activity is determined by the SLEDAI scores and the degree of fatigue is determined by the FSS (Fatigue Severity Scale).

Results After supplementation for 3 months, this study showed that decreased of SLEDAI score in the supplementation group was greater than the placebo group. The decreased of FSS in the supplementation group was also greater than the placebo group.

Conclusions Adding curcumin on vitamin D supplementation, decreased SLEDAI scores and FSS greater than vitamin D supplementation plus placebo in SLE patients with hypovitamin D.

257

THE EFFECT OF ADDING CURCUMIN ON VITAMIN D3 SUPPLEMENTATION ON ANTI-DSDNA LEVELS AND PROTEINURIA, IN SLE PATIENTS WITH HYPOVITAMIN D

¹CS Wahono, ²H Susianti*, ¹ZD Wahyuni, ¹I Saveria, ¹CD Setyorini, ²K Handono, ¹H Kalim. ¹Universitas Brawijaya/Saiful Anwar Hospital, Internal Medicine, Malang, Indonesia; ²Universitas Brawijaya/Saiful Anwar Hospital, Clinical Pathology, Malang, Indonesia

10.1136/lupus-2017-000215.257

Background and aims Vitamin D has important roles in the regulation of the immune system in Lupus. Seventy percent of lupus patients in Indonesia are experienced hypovitamin D. Curcumin is a natural VDR ligand and has synergic effect with vitamin D. This study was aimed to determine the effect of adding curcumin on vitamin D supplementation on anti-dsDNA serum levels and proteinuria, in SLE patients with hypovitamin D.

Methods This was a randomised controlled trial, double blind study. Forty SLE patients with hypovitamin D were studied, that randomized into two groups: 20 patients (supplementation group) received vitamin D (cholecalciferol 1200 IU daily) with curcumin 20 mg (three times daily); and 20 patients (placebo group) was given vitamin D (cholecalciferol 1200 IU daily)

and placebo (3 times daily), for 3 months. Anti-ds DNA serum levels were measured by ELISA and urine protein were measured by urine albumin creatinine ratio (UACR).

Results After supplementation for 3 months, this study showed that decreased of anti-dsDNA serum levels in the supplementation group was significantly greater than in the placebo group. The decreased of UACR in the supplementation group was also significantly greater than the placebo group.

Conclusions Adding curcumin on vitamin D supplementation can decrease anti-dsDNA serum levels and proteinuria greater than vitamin D supplementation plus placebo in SLE patients with hypovitamin D.

258 THE EFFECT OF ADDING CURCUMIN ON VITAMIN D3 SUPPLEMENTATION ON CYTOKINES BALANCE, IN SLE PATIENTS WITH HYPOVITAMIN D

¹CS Wahono*, ¹I Saveria, ¹CD Setyorini, ¹ZD Wahyuni, ²K Handono, ¹H Kalim. ¹Universitas Brawijaya/Saiful Anwar Hospital, Internal Medicine, Malang, Indonesia; ²Universitas Brawijaya/Saiful Anwar Hospital, Clinical Pathology, Malang, Indonesia

10.1136/lupus-2017-000215.258

Background and aims Vitamin D has important roles in the regulation of the immune system in Lupus. Seventy percent of lupus patients in Indonesia are experienced hypovitamin D. Curcumin is a natural VDR ligand and has synergic effect with vitamin D. This study was aimed to determine the effect of adding curcumin on vitamin D supplementation on IFN- γ /IL-4 ratio and IL-17/TGF- β ratio, in SLE patients with hypovitamin D.

Methods This was a randomised controlled trial, double blind study. Forty SLE patients with hypovitamin D were studied, that randomised into two groups: 20 patients (supplementation group) received vitamin D (cholecalciferol 1200 IU daily) with curcumin 20 mg (three times daily); and 20 patients (placebo group) was given vitamin D (cholecalciferol 1200 IU daily) and placebo (3 times daily), for 3 months. Cytokines serum levels (IFN- γ , IL-4, IL-17, TGF- β), were measured by ELISA.

Results After supplementation for 3 months, this study showed that decreased of IFN- γ /IL-4 ratio in the supplementation group was significantly greater than in the placebo group. The decreased of IL-17/TGF- β ratio in the supplementation group was also significantly greater than the placebo group.

Conclusions Adding curcumin on vitamin D supplementation can decrease IFN- γ /IL-4 ratio and IL-17/TGF- β ratio than vitamin D supplementation plus placebo in SLE patients with hypovitamin D.

259 THE EFFECT OF VITAMIN D3 SUPPLEMENTATION ON THE ANTI-DSDNA LEVELS AND URINE PROTEIN IN SLE PATIENTS WITH HYPOVITAMIN D

¹H Susianti*, ²CS Wahono, ²ST Priyantoro, ¹K Handono, ²H Kalim. ¹Faculty of Medicine Brawijaya University/Dr. Saiful Anwar General Hospital, Clinical Pathology, Malang, Indonesia; ²Faculty of Medicine Brawijaya University/Dr. Saiful Anwar General Hospital, Internal Medicine, Malang, Indonesia

10.1136/lupus-2017-000215.259

Background and aims Vitamin D has important role in the regulation of the immune system in Lupus. 71% of lupus patients in Indonesia experienced hypovitamin D. This study was aimed to determine the effect of vitamin D

supplementation on the levels of anti ds dna and degree of urine protein in lupus patients with hypovitamin D.

Methods Thirty nine SLE patients with hypovitamin D were studied, that randomized into two groups: 20 patients was given vitamin D and 19 patients received placebo for 3 months. Anti-ds DNA levels were measured by ELISA and urine protein were measured by dipstick method.

Results Anti-dsDNA levels in the supplement group before and after giving vitamin D were 226.84 ± 82.11 vs 191 ± 72.55 ($p=0.00$), and the placebo group were 233.69 ± 66.52 vs 227.72 ± 61.21 ($p=0.077$). The degrees of urine protein in the supplement group before and after treatment were 24 vs 12 U/ml ($p=0.003$) and the placebo group were 16 vs 10 U/ml ($p=0.070$).

Conclusions Vitamin D supplementation plays a role on decreasing the levels of anti ds-DNA and urine protein in SLE patients with hypovitamin D.

260 THE EFFECT OF VITAMIN D3 SUPPLEMENTATION ON THE DISEASE ACTIVITIES AND DEGREE OF FATIGUE IN SLE PATIENTS WITH HYPOVITAMIN D

¹CS Wahono*, ²H Susianti, ¹A Rifai, ²K Handono, ¹H Kalim. ¹Faculty of Medicine Brawijaya University/Dr. Saiful Anwar General Hospital, Internal Medicine, Malang, Indonesia; ²Faculty of Medicine Brawijaya University/Dr. Saiful Anwar General Hospital, Clinical Pathology, Malang, Indonesia

10.1136/lupus-2017-000215.260

Background and aims Vitamin D has important roles in the regulation of the immune system in Lupus. 71% of lupus patients experienced hypovitamin D in Indonesia. This study was aimed to determine the effect of vitamin D supplementation on the degree of disease activity and degree of fatigue in SLE patients with hypovitamin D.

Methods Thirty nine SLE patients with hypovitamin D were studied, that randomised into two groups: 20 patients was given vitamin D and 19 patients received placebo for 3 months. Disease activity is determined by the SLEDAI scores and the degree of fatigue is determined by the FSS (Fatigue Severity Scale).

Results This study showed that supplementation of vitamin D 1200 IU per day increased 6.55 ± 1.27 ng/cc of 25 (OH) D. The decreased of SLEDAI scores in the supplementation group were greater than the placebo group (6.45 ± 3.07 vs 0.17 ± 1.63), $p=0.000$. The decreased of Fatigue Severity Scale in the supplementation group also greater than the placebo group (2.27 ± 0.73 vs 0.005 ± 0.62), $p=0.000$.

Conclusions Vitamin D supplementation plays a role on improving the activity of the disease (SLEDAI score) and the condition of fatigue (FSS) in SLE patients with hypovitamin D.

261 LONG-TERM EFFECTS OF OF HYDROXYCHLOROQUINE ON METABOLISM OF SERUM LIPIDS AND LEFT VENTRICULAR STRUCTURE AND FUCTION IN PATIENTS OF SYSTEMIC LUPUS ERYTHEMATOSUS

¹J Meng. ¹Beijing Chao-Yang Hospital- Capital Medical University, Department of Rheumatology and Immunology, Bei Jing, China

10.1136/lupus-2017-000215.261