in vitro with IL-3 for 6 or 24 hours. WB RNASeq analysis was also undertaken in n=31 SLE donors from the Monash Lupus Clinic and n=28 HDs.

Results Serum IL-3 levels correlated with serum IFN α ($r=0.612$, 95% CI 0.455–0.733, $p<0.001$). IL-3 stimulation *in vitro* altered 794 genes ($-1\geq\log_{2}FC\geq 1$, $FDR<0.05$). Thirty-five of these genes overlapped with differentially expressed genes between SLE and HD. These 35 genes were expressed in 28/31 SLE donors, revealing the presence of an 'IL-3 gene signature'. There was strong correlation between the IL-3 signature and an IFN signature determined by hierarchical clustering of the five hundred most variable genes in SLE donors ($r=0.939$, 95% CI 0.898–0.964, $p<0.0001$).

Conclusions We have previously reported a novel anti-IL-3R α mAb (CSL362/JNJ-473), which depletes pDCs and reduces IFN α production, as well as neutralising IL-3 signalling (Oon S, JCI Insight, 2016). An association between IL-3 and IFN α was found in this study, raising the possibility that CSL362 may be especially useful for lupus patients with a dual IL-3/IFN gene signature.

303 URINARY VCAM 1 AS A DISEASE ACTIVITY INDICATOR IN LUPUS NEPHRITIS

¹S Padiyar*, ¹J Mathew, ²TS Vijayakumar. ¹Christian Medical College, RHEUMATOLOGY, Vellore- Tamilnadu, India; ²Christian Medical College, NEPHROLOGY, Vellore- Tamilnadu, India

10.1136/lupus-2017-000215.303

Scatter plot showing VCAM values in cases (with active nephritis)

And controls (without nephritis)

Abstract 303 Figure 2

Abstract 303 Figure 1

Background and aims Currently we do not have a biomarker that can closely reflect the renal disease activity. So the aim of this study is to study the utility of urinary VCAM 1 (Vascular cell adhesion molecule 1) in lupus nephritis.

Methods It was a diagnostic case control study. The patients presenting to Rheumatology outpatient department were recruited. Patients were divided into 2 groups, SLE without active nephritis and SLE with active nephritis based on the renal SLEDAI. Urinary VCAM1 was tested in all patients using an early morning spot urine sample using ELISA. Renal biopsy was done in patients with active nephritis. VCAM1 levels were compared with the renal SLEDAI, renal biopsy disease activity (ISN/RPS) and standard of care markers. The results were analysed using SPSS software version 16. The validity and predictive value statistics was presented with 95 percent confidence interval.

Results Urinary VCAM 1 levels had significant correlation ($p=0.01$) with disease activity based on renal SLEDAI. However, the correlation between the biopsy findings and VCAM levels was not statistically significant. Class 4 and 5 lupus nephritis had higher VCAM level than the lower classes. A positive correlation ($r=0.38$) was found between VCAM 1 and double stranded DNA. There was a negative correlation between C3 value and VCAM ($r=-0.19$). The sensitivity and specificity of urinary VCAM 1 is 65.22% and 75% respectively. The cut off value of VCAM is 23.8 pg/mg of creatinine.

Conclusions Urinary VCAM 1 may not independently, but combined with other markers may be a promising biomarker for disease activity in lupus nephritis.

Abstract 303 Figure 3

304 TNF- α PROMOTER POLYMORPHISMS (G-238A AND G-308A) ARE ASSOCIATED WITH SUSCEPTIBILITY TO SYSTEMIC LUPUS ERYTHEMATOSUS: A STUDY IN P. FALCIPARUM ENDEMIC AREA

¹AK Panda*, ¹H MAHATO, ²R Tripathy, ³BK Das. ¹Central University of Jharkhand, Centre for Life Sciences, Ranchi, India; ²SCB Medical College, Department of Biochemistry, Cuttack, India; ³SCB Medical College, Department of Medicine, Cuttack, India

10.1136/lupus-2017-000215.304

Background and aims Tumour necrosis factor- α (TNF- α) is a proinflammatory cytokine associated with *P. falciparum* malaria and autoimmune disorders. Elevated plasma TNF- α has been linked to *P. falciparum* malarial severity and mortality. Higher levels of TNF- α has also been reported in systemic lupus erythematosus (SLE). Two functional common polymorphisms (G-238A and G-308A) at promoter region of TNF- α gene have been linked to SLE susceptibility in different population. In the present report, we conducted a case control study to investigate association of TNF- α (G-238A and G-308A) polymorphisms with susceptibility/resistance to SLE development in a *P. falciparum* malaria endemic cohort.

Methods A total of 204 female SLE patients and 224 age and sex matched healthy controls were enrolled in the study. TNF- α polymorphisms (G-238A and G-308A) were typed by polymerase chain reaction and restriction length polymorphism (PCR-RFLP). Plasma level of TNF- α was quantified by enzyme linked immunosorbent assay.

Results The prevalence of heterozygous mutants and minor alleles of TNF- α (G-238A and G-308A) polymorphisms were significantly higher in SLE patients compared to healthy controls. Furthermore, heterozygous (GA) and minor allele (A) of TNF- α (G-238A) polymorphism were associated with susceptibility to lupus nephritis. SLE patients displayed higher levels of plasma TNF- α compared to healthy controls. TNF- α (G-238A and G-308A) variants were associated with higher plasma TNF- α in both SLE patients and healthy control.

Conclusions The results of the present study demonstrate that TNF- α (G-238A and G-308A) variants are associated with higher plasma TNF- α level and increased susceptibility to development of SLE in malarial endemic areas.

305 ASSOCIATION OF C4A AND C4B GENOMIC COPY NUMBER VARIATIONS IN ADULT AND PAEDIATRIC SYSTEMIC LUPUS ERYTHEMATOSUS

¹P Das*, RW Minz¹, ¹B Saikia, ²A Sharma, ³S Singh. ¹Postgraduate Institute of Medical Education and Research – Chandigarh, Immunopathology, Chandigarh, India; ²Postgraduate Institute of Medical Education and Research – Chandigarh, Internal Medicine, Chandigarh, India; ³Postgraduate Institute of Medical Education and Research – Chandigarh, Advanced Paediatric Centre, Chandigarh, India

10.1136/lupus-2017-000215.305

Background and aims C4 complement gene has been observed to be a susceptibility gene for SLE. Lower C4 gene (C4A and C4B) copy number (CN) is a risk factor for SLE, where as higher C4 CN is a protective factor. We investigated the association of C4 gene copy number variation in a north Indian cohort of SLE patients

Methods We recruited 112 aSLE and 52 pSLE patients with 115 healthy adult (CA) and 60 healthy paediatric (CP) controls and compared for C4A and C4B CN by RT-PCR, serum C3, C4 by nephelometry and ANA autoantibodies by line blot assay

Results C4A low copy number was higher in pSLE (OR=1.82, p=0.67) and aSLE (OR=1.51, p=0.41) as compared to their respective controls, pSLE had higher C4A low copy number than the aSLE (OR=1.33, p=0.58), though they were not statistically significant. C4A and C4B CN negatively correlated with several ANA autoantibodies. The total C4 (C4A +C4B) CN negatively correlated with Ro52 (r=-0.29, p=0.03), dsDNA (r=-0.32, p=0.02), SSB (r=-0.33, p=0.01), nucleosome (r=-0.28, p=0.04) and histone (r=-0.34, p=0.01) in pSLE and with nucleosome (r=-0.20, p=0.03) in aSLE. The total C4 CN positively correlated with serum C4 level (r=0.26, p=0.007) in both groups of patients

Conclusions We demonstrate that low copy numbers of complement genes correlate with the propensity for increased antibody secretion in both aSLE and pSLE. Thus, more productions of autoantibodies cause large number of immune complex formation with defective clearance process, due to low serum C4 level and low gene copy number