

### 333 IN VIVO THERAPEUTIC SUCCESS OF MICRORNA-155 (MIR-155) ANTAGOMIR IN A MOUSE MODEL OF LUPUS ALVEOLAR HAEMORRHAGE

Y Wang\*, S Zhou, N Shen. *RenJi Hospital- School of Medicine- Shanghai JiaoTong University, Department of Rheumatology, Shanghai, China*

10.1136/lupus-2017-000215.333

**Background and aims** Diffuse alveolar haemorrhage (DAH) is a rare but life-threatening complication of systemic lupus erythematosus (SLE). Pristane-treated B6 mice develop severe DAH within 2 weeks of treatment. MicroRNA-155 (miR-155) is a pleiotropic microRNA that plays a crucial role in the regulation of immune responses. The purpose of this study was to examine the role of miR-155 in the development of DAH in pristane-induced lupus using miR-155-knockout (miR-155<sup>-/-</sup>) mice and miR-155 antagomir to silence miR-155.

**Methods** DAH was induced by an intraperitoneal injection of 0.5 mL of pristane. MiR-155 antagomir was intravenously administered to silence miR-155 expression. Lung tissues were collected for RNA extraction and were embedded in paraffin for sectioning. Gene expression profiling data were analysed using Ingenuity Pathway Analysis. Real time q-PCR was used for single validation. Luciferase reporter assay and RNA-Ago2 immunoprecipitation were performed for target validation.

**Results** MiR-155 expression was significantly increased in the development of DAH. Disease progression was reduced in miR-155<sup>-/-</sup> mice and by *in vivo* silencing of miR-155 using miR-155 antagomir. MiR-155 silencing dampened pristane-induced ectopic activation of multiple inflammatory pathways, and reduced the expression of pro-inflammatory cytokines. Several negative regulators of nuclear factor (NF)-κB signalling were inhibited by pristane, and were re-activated in miR-155<sup>-/-</sup> mice. In particular, the anti-inflammatory factor peroxisome proliferator-activated receptor-α was identified as a direct target of miR-155.

**Conclusions** MiR-155 promotes pristane-induced lung inflammation. MiR-155 contributes to ectopic activation of NF-κB signalling pathways by targeting multiple negative regulators. MiR-155 antagomir may be a promising therapeutic strategy for treating acute lung inflammation in lupus.

### 334 SERUM IL-18 AS BIOMARKER IN PREDICTING LONG-TERM RENAL OUTCOME AMONG PEDIATRIC-ONSET SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS

JL <sup>1</sup>KW Yeh\*, <sup>1</sup>CY Wu, <sup>2</sup>HY Yang, <sup>1</sup>J L. <sup>1</sup>*Chang Gung Memorial Hospital, Paediatrics, Taoyuan, Taiwan R.O.C.*; <sup>2</sup>*Chang Gung Memorial Hospital, Nephrology, Taoyuan, Taiwan R.O.C.*

10.1136/lupus-2017-000215.334

**Background and Aims** An urge of biomarker identification is needed to better monitor lupus nephritis (LN) disease activity, guide clinical treatment, and predict patient's long-term outcome. With the proinflammatory effect and its association with inflammasomes, the significance of interleukin-18 (IL-18) among pediatric-onset systemic lupus erythematosus (pSLE) patient.

**Methods** In a pSLE cohort of 96 patients with an average follow-up period of 10.39±3.31 years, clinical data and laboratory workups including serum IL-18 were collected at time of disease onset and 6 months after treatment despite their initial

renal status. Through Cox regression analysis, the parameters at baseline and at 6 months posttreatment were carefully analysed.

**Results** Average age of all cases was 12.74±3.01 years old and 65 of them underwent renal biopsy at the time of diagnosis. Nine (9.38%) progressed to end-stage renal disease (ESRD) and 2 (2.08%) died during follow-up. Through multivariate analysis, serum IL-18 level 6 months posttreatment was found to be the most unfavourable factor associating poor clinical outcome despite patient's initial renal status. The presentation of serum IL-18 in its correlation with SLE global disease activity as well as the presence and severity of LN were all significant ( $p < 0.001$ ,  $p = 0.03$ , and  $p = 0.02$ , respectively). The histological classification of LN was not associated with the level of IL-18 among the pSLE patients ( $p = 0.64$ ).

**Conclusions** The role of serum IL-18 as biomarker representing global disease activity and status of renal flares among pSLE population was shown for the first time. Additionally, we have identified IL-18 at 6 months posttreatment a novel marker for long-term renal outcome prediction.

### 335 MICRORNA-21 IS A CRITICAL REGULATOR OF AUTOIMMUNITY THROUGH PROMOTING EFFECTOR AND METABOLIC FUNCTION OF PATHOGENIC TH17 CELLS

X Yu\*, D Dai, N Shen. *RenJi Hospital- School of Medicine- Shanghai JiaoTong University, Department of Rheumatology, Shanghai, China*

10.1136/lupus-2017-000215.335

**Background and aims** Systemic lupus erythematosus is a prototypical autoimmune disease that causes mortality and morbidity worldwide. Recent studies suggest proinflammatory TH17 cells are key pathogenic factors that contribute to lupus nephritis. Our group previously demonstrate that microRNA-21 was highly upregulated in CD4<sup>+</sup> T cells from both lupus patients and lupus-prone mice. However, the role of microRNA-21 in pathogenic TH17 cells and they-mediated autoimmune diseases is still unclear. In this study, we systemically dissect the role of microRNA-21 in the differentiation and effector function of pathogenic TH17 cells.

**Methods** MicroRNA-21 knockout and conditional knockout mice were generated. EAE was induced to study the role of microRNA-21 in pathogenic TH17 cell-mediated autoimmune diseases. RNA-seq, RIP-seq and DAVID bioinformatic analysis were conducted to find key microRNA-21 regulated pathway and molecular targets in pathogenic TH17 cells. Metabolic assays were done to study the glycolytic activity of microRNA-21-deficient pathogenic TH17 cells.

**Results** In this study, we demonstrate that IL-6-STAT3 signalling induced microRNA-21 is essential for the late stage commitment and maintenance of pathogenic TH17 cells by targeting key regulators. MicroRNA-21-deficient TH17 cells express less pathogenic TH17 signature genes and show less glycolytic activity. Conditional deletion of microRNA-21 in CD4<sup>+</sup> T cells protects mice from EAE while loss of microRNA-21 expression by dendritic cells and myeloid cells do not.

**Conclusions** These findings suggest that microRNA-21 is a novel cell-intrinsic regulator of the commitment and metabolic function of pathogenic TH17 cells. It may be a potential therapeutic candidate with which to reprogram the immune system and help prevent and treat autoimmune diseases.