Abstract 419 Table 1 Patient Characteristics (at incident SLE hospitalisation) and study outcomes.

	SLE as reason for admission (n = 1516)	Matched Controls (n = 1320)	χ^2 or t-test
	MoCT or n(%)	MoCT or n(%)	(p-value)
Age	45.15 $\pm$ 19.44	50.07 $\pm$ 21.66	<0.001
Female	1172 (77.3%)	1031 (78.1%)	0.611
Indigenous	107 (7.1%)	21 (1.6%)	<0.001
Length of Stay (days)	5 (IQR 1, 11)	1 (IQR 1, 5)	<0.001
Privately Insured	176 (31.5%)	325 (52.3%)	<0.001
Diagnosed with Ischaemic Heart Disease	46 (3.0%)	28 (2.1%)	0.128
Diagnosed with a Cerebral Ischaemic Disease	16 (1.1%)	8 (0.6%)	0.193
Diagnosed with Hypertension	165 (10.9%)	71 (5.4%)	<0.001
Diagnosed with an Atherosclerotic Condition	54 (3.6%)	11 (0.8%)	<0.001
Diagnosed with a Kidney Disorder	308 (20.3%)	13 (1.0%)	<0.001
Diagnosed with a Thrombotic Disorder	10 (0.7%)	5 (0.4%)	0.304
Patient died during the hospital admission	37 (2.4%)	17 (1.3%)	<0.001
MoCT: Measure of Central Tendency, i.e. mean $\pm$ standard deviation or median (interquartile range)			

421

ASSESSING THE QUALITY, RELIABILITY AND READABILITY OF ONLINE HEALTH INFORMATION REGARDING SYSTEMIC LUPUS ERYTHEMATOSUS (SLE)

¹M Reynolds*, ¹A Hoi, ^{1,2}R Buchanan. ¹Austin Health, Rheumatology, Heidelberg, Australia; ²University of Melbourne, Department of Medicine, Melbourne, Australia

10.1136/lupus-2017-000215.421

Background and aims Systemic lupus erythematosus (SLE) is a complex multi-system autoimmune disorder. Patients frequently access the internet to increase their knowledge about the condition.

We assessed the quality, reliability and readability of online sites relating to SLE.

Methods The search-phrases 'systemic lupus erythematosus' was used with the three most commonly accessed internet search engines (Google, Bing and Yahoo) to identify websites. The first 25 hits (excluding duplicates and excluded websites) for each search were assessed for quality using the DISCERN

instrument (scores 15–80 points), reliability using the JAMA benchmarks (assessing authorship, attribution of references, currency (date of posted content) and disclosure of conflict of interest) and readability using the Gunning-FOG index (ideal score 7–8).

Results There was significant concordance between the hits returned from each search engine with 26 unique websites identified. The average DISCERN score was 47.7 (SD 13.2), and ranged from 19–75. Websites that appeared earlier in searches did not have higher DISCERN scores (Pearson correlation –0.16).

Currency was present in 50.0%, appropriate authorship in 46.2% and attribution of references and disclosure of interest in only 15.4% of websites.

The average readability of the websites was 9.3 (SD 3.4) using the FOG index.

Conclusions The overall quality of online health information relating to SLE is only fair.

Abstract 419 Figure 1 Kaplan-Meier survival analysis of all-cause mortality for SLE patients and age- and sex-matched controls (free of rheumatic disease) from index hospitalisation.

Reliability as measured by the JAMA benchmarks was average to poor.

The readability is higher than recommended for the general public, which may limit understanding.

This assessment highlights the need for clinicians to provide patients with alternative sources of high quality information regarding SLE.

422

ANTI-RNP/SM ANTIBODIES PLUS LUPUS ANTICOAGULANT AS RISK FACTOR FOR THROMBOSIS IN PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS

¹MC Zamora-Medina*, ¹A Hinojosa-Azaola, ²AG Vargas-Ruiz, ¹J Sanchez-Guerrero, ¹C Nunez-Alvarez, ¹J Romero-Diaz. ¹Instituto Nacional de Ciencias Medicas y Nutricion S.Z., Immunology and Rheumatology, Mexico city, Mexico; ²Instituto Nacional de Ciencias Medicas y Nutricion S.Z., Haematology, Mexico city, Mexico

10.1136/lupus-2017-000215.422

Background and aims Thrombosis still remains as main cause of morbidity and mortality in SLE patients. Results from a recent study identified a strong association of anti-RNP/Sm abs+LA with thrombosis. We aimed to validate this association

Abstract 419 Figure 2 Kaplan-Meier survival analysis of cancer outcomes for SLE patients and age- & sex-matched controls (free of rheumatic disease conditions) from incident hospitalisation.

Methods We studied 63 SLE patients (>4 criteria/ACR) with confirmed history of thrombosis after SLE diagnosis. As controls, 63 SLE patients without thrombosis, matched by age, gender and lupus duration were included. Disease characteristics, medication, traditional risk factors for thrombosis and thrombotic event information were retrieved from clinical files. A blood sample was drawn to determine anti-cardiolipin (IgG/IgM), antiβ2-GP1 (IgG/IgM), LA, anti-RNP/Sm and anti-Sm antibodies. Sensitivity, specificity, positive and negative predictive values (PPV, NPV) and Likelihood Ratios (LR) were calculated.

Results One hundred and twenty six SLE patients were studied. Cases and controls were similar in age, gender and disease duration (p=NS). There were no differences in the prevalence of traditional risk factor for thrombosis between cases and controls (p=NS). Among patients with thrombosis a higher frequency of anti-RNP/Sm (83% vs 62%, p<0.001), LA (62% vs 19%, p<0.001), aPL triple marker (17% vs 2%, p=0.04) and anti-RNP/Sm+AL combination (52% vs 14%, p<0.001) was observed. The combination of anti-RNP/Sm +AL showed a sensitivity 52%, specificity 86%, PPV 78% and NPV 65%, positive LR 3.67 (IC 95% 1.92–7.04) and negative LR 0.55 (IC 95% 0.42–0.74).

Conclusions This study confirmed that anti-RNP/Sm+LA association represents a risk factor for thrombosis in SLE patients.