

neutrophil elastase (NE), proteinase 3 (PR3), and cathepsin G (CG) and myeloperoxidase (MPO) expression by qRT-PCR. NE^{-/-}/PR3^{-/-}/CG^{-/-} triple knockout mice and *in vitro* depletion of neutrophils approaches were performed to determine the role of NSPs and neutrophils in estrogen-mediated inflammatory responses. The splenic neutrophil and NSPs expression in lupus-prone MRL-*lpr*, B6-*lpr* and NZB/W_{F1} and their respective controls were also analysed.

Results Although oestrogen reduced total splenocytes number, it markedly increased the splenic neutrophil numbers, NSPs and MPO expression in B6 mice (Figure 1). Splenic neutrophils, NSPs and MPO were also significantly increased in MRL-*lpr*, B6-*lpr* and NZB/W_{F1} mice (Figure 2). Despite of the critical role of NSPs and neutrophils in inflammation, depletion of NSPs *in vivo* did neither affect oestrogen's ability to increase in splenic neutrophils nor the induction of inflammatory mediators from *ex vivo* activated splenocytes, and depletion of splenic neutrophils *in vitro* had also no obvious effect on NSPs expression (due to the increase of NSPs in cells other than neutrophils) and LPS-induced IFN γ and MCP-1 (Figure 3).

Conclusions Overall, we demonstrated a remarkable commonality with regards to the increase of neutrophils and NSPs in the spleens of autoimmune-prone mice and estrogen-treated B6 mice.

113

EFFECT OF MICROPARTICLES DERIVED FROM PATIENTS WITH LUPUS ERYTHEMATOSUS SYSTEMIC (SLE) ON MODULATION OF MICRORNAS 146A AND 126, IN A MONOCYTE CELL LINE

¹G Vasquez, ¹LJ Carmona-Perez*, ²CH Muñoz-Vahos, ²AL Vanegas-García, ¹M Rojas. ¹Universidad de Antioquia, Grupo de Inmunología Celular e Inmunogenética GICIG. Instituto de Investigaciones Médicas- Facultad de Medicina-, Medellín, Colombia; ²Hospital Universitario San Vicente Fundación, Sección Reumatología, Medellín, Colombia

10.1136/lupus-2017-000215.113

Background and aims Important miRNAs are involved in the modulation of immune functions and can be found in the intra- and extracellular environments, in circulation attached to RNA-binding molecules or packed in form of microparticles (MP). Given this, MP could have an important role in inter-cellular communication, modulation and expression of miRNAs in their target cells.

To establish whether MP of SLE patients can modulate the expression of miRNAs (miRNA 126, miRNA 146a) and their target molecule (interferon response factor 5, IRF5) in the monocyte cell line U937.

Methods MP obtained from serum of SLE and other autoimmune diseases (OAD) patients, and healthy controls were used as stimulus in the cell line U937 to evaluate their effect over: 1) The expression of membrane markers through flow cytometry, 2) content of miRNAs 126 and 146a through PCR and 3) expression of their target molecule, IRF5 by Western Blot.

Results We observed, a decrease in HLA-DR, CD18, CD119 and an increase of IL-6 in U937 stimulated with MP from healthy controls, patients with active and inactive SLE, as well as patients with OAD. Additionally, a positive effect over the expression of miR126 and a negative effect over the expression of miR146a were observed. IRF5 as a target of miRNA146, did not change after MP treatment independent of MP origin.

Conclusions Our results suggested that MPs may have a regulatory effects, inducing a decreased expression of membrane molecules and miRNAs-146 levels, without effect in IRF5. In addition, MPs increased levels of cytokines and miRNA-126, latter is related with the demethylation.

114

N-ACETYL-L-CYSTEINE (NAC) CONTROLS OSTEOCLASTOGENESIS THROUGH REGULATING TH17 DIFFERENTIATION AND RANKL PRODUCTION IN RHEUMATOID ARTHRITIS

¹HR Kim*, ²BM Kim, ¹KA Lee, ¹SH Lee, ²KW Kim. ¹Konkuk University Hospital, Rheumatology, Seoul, Republic of Korea; ²Seoul St. Mary's Hospital- College of Medicine- The Catholic University, Convergent Research Consortium for Immunologic disease, Seoul, Republic of Korea

10.1136/lupus-2017-000215.114

Background and aims This study aimed to determine the regulatory role of N-Acetyl-l-cysteine (NAC), an antioxidant, in IL-17-induced osteoclast differentiation in rheumatoid arthritis (RA).

Methods After RA synovial fibroblasts were stimulated by IL-17, the expression and production of RANKL was determined by real-time PCR and ELISA. Human peripheral blood monocytes were cultured with M-CSF, IL-17, RANKL, and/or various concentrations of NAC, followed by counting of the cells for tartrate-resistant acid phosphatase activity to determine osteoclast formation. Osteoclastogenesis was also determined after cocultures of IL-17-stimulated RA synovial fibroblasts, Th17 cells and various concentrations of NAC with monocytes. After human peripheral CD4⁺ T cells were cultured with NAC under Th17 condition, IL-17, IFN- γ , IL-4, Foxp3, RANKL and IL-2 expression and production was determined by flow cytometry or ELISA.

Results When RA synovial fibroblasts were stimulated by IL-17, IL-17 stimulated the production of RANKL, and NAC reduced the IL-17-induced RANKL production in a dose-dependent manner. NAC decreased IL-17-activated phosphorylation of mTOR, JNK and I κ B. When human peripheral blood CD14⁺ monocytes were cultured with M-CSF and IL-17 or RANKL, osteoclasts were differentiated, and NAC reduced the osteoclastogenesis. After human peripheral CD4⁺ T cells were co-cultured with IL-17-pretreated RA synovial fibroblasts or Th17 cells, NAC reduced their osteoclastogenesis. Under Th17 polarising condition, NAC decreased Th17 cell differentiation and IL-17 and RANKL production.

Conclusions NAC inhibits the IL-17-induced RANKL production in RA synovial fibroblasts and IL-17-induced osteoclast differentiation. NAC also reduced Th17 polarisation. NAC could be a supplementary therapeutic option for inflammatory and bony destructive processes in RA.

115

QUANTITATIVE AND FUNCTIONAL EVALUATION OF PLASMA MICROPARTICLES IN SYSTEMIC LUPUS ERYTHEMATOSUS

^{1,2}A Lateef*, ¹L Shaffiee, ³GK Gill, ¹P Cheung, ⁴YC Lim. ¹National University Hospital, Medicine, Singapore, Singapore; ²Yong Loo Lin School of Medicine- National University of Singapore, Medicine, Singapore, Singapore; ³National University of Singapore, Physiology, Singapore, Singapore; ⁴National University of Singapore, Pathology, Singapore, Singapore

10.1136/lupus-2017-000215.115