Abstract 156 Table 1 Multivariate logistic regression analysis

Variables	OR	95% CI		P value
High Cyp61 level	7.822	2.224	41.138	0.002
Serositis	8.244	1.854	36.663	0.006
Anti-RNP positivity	3.902	1.067	14.267	0.040
Renal disorders	0.174	0.044	0.689	0.013
Anti-dsDNA positivity	0.171	0.046	0.636	0.008

Results Group A did not differ from control group. Group B had higher titers of anti-ABCA1 antibodies ($p=0.004$), with 4 patients showing positive anti-ABCA1 titers (11.4%). In group B, anti-ABCA1 antibodies titers tend to negatively correlate with HDL ($p=0.06$) and apolipoproteinA-I levels ($p=0.04$). In both groups, anti-HDL antibodies did not correlate with anti-ABCA1 titers.

Conclusions This is the first report of naturally occurring antibodies against ABCA1. These antibodies are increased in patients with SLE with at least 4 SLICC classification criteria. Future studies will determine their pathogenic role.

This work was supported by the Portuguese Foundation for Science and Technology via the post-doctoral SFRH/BPD/112411/2015 and Fundação Oriente.

158 CONTRIBUTORS TO DISEASE FLARES AMONG FILIPINO PATIENTS WITH SLE

ME Fernandez, MFJ Edar, S Navarra. *University of Santo Tomas Hospital, Internal Medicine-Rheumatology, Manila, Philippines*

10.1136/lupus-2017-000215.158

Background and aims SLE disease flares significantly contribute to morbidity and mortality in SLE. This study describes the characteristics and contributing factors to disease flares in a cohort of Filipino SLE patients.

Methods Included were SLE patients with disease flare seen consecutively from January 2012 to December 2015 at the Lupus Clinics of University of Santo Tomas (UST) Hospital, Manila, Philippines. Demographics, organ involvement and SLE flare severity by SELENA-SLEDAI Flare Index (SFI) were described. Contributing factors to flare were identified based on physician assessment notes and patient responses to the Life Experiences Survey (LES).

Results There were 100 disease flares in 73 patients (68 females). Mean age at SLE diagnosis was 25.15 ± 8.57 years (4-43), disease duration at each flare occurrence was 7.40 ± 4.42 (<1-21) years. Flares were severe in 60, mild to moderate in 40. Organ involvement included renal in 47 (32.41%), mucocutaneous 40 (27.59%), musculoskeletal 26 (17.93%), vasculitis 11 (7.59%), hematologic 11 (7.59%), neurologic 6 (4.14%), ocular (32.07%) and cardiopulmonary 1

(0.68%). Stress and infection in 35 (30.97%) each, were the leading contributors to flare. Others included non-compliance in 18 (15.93%), excess sun exposure 7 (6.20%), pregnancy 3 (2.66%), fetal loss 3 (2.66%), surgery and drugs during 1 (0.88%) flare occurrence each, and unknown in 10 (8.85%).

Conclusions Renal, mucocutaneous and musculoskeletal systems were most commonly involved in a disease flare. Stress and infection were the leading factors contributing to a flare. These findings underscore need for holistic management approach in SLE, integrating effective disease control with patient education

159 NEUTROPHIL-TO-LYMPHOCYTE RATIO AND SYSTEMIC LUPUS ERYTHEMATOSUS: A NEW PARAMETER FOR DISEASE ACTIVITY ASSESSMENT?

¹H Gunawan*, ¹A Awalia, ¹J Soero. *¹Airlangga University, Internal Medicine – Rheumatology Division, Surabaya, Indonesia*

10.1136/lupus-2017-000215.159

Background and aims Systemic Lupus Erythematosus (SLE) is a complex autoimmune disease characterised by enhanced apoptosis in conjunction with high levels of autoantibodies and deregulated cytokine productions. It contributes to immune dysfunction, tissue inflammation, and organ damage. Neutrophil to Lymphocyte ratio (NLR) is a promising marker in SLE, but its relationships with SLE's disease activity in Indonesian population has not been evaluated.

To investigate the correlation between NLR and SLE disease activity.

Methods A cross sectional study with consecutive sampling technique was conducted with 70 Indonesian SLE patients in Dr. Soetomo Hospital Surabaya at May 2016-July 2016. Peripheral blood count was evaluated during assessment with flowcytometry technique and disease activity was evaluated with Systemic Lupus Activity Measure (SLAM) score. Data analysis was done with Spearman's correlation rank test and ROC curve using SPSS v22.0.

Results There were 19 patients with SLAM score <7 (group 1) and 51 patients with SLAM score ≥ 7 (group 2). We found that lymphocyte count and NLR was statistically significant between groups [1.58 (0.66-2.73) vs 0.95 (0.14-3.49);