Abstract 159 Tabel 1 Basic characteristics of the study

Parameter	SLAM <7 (n=19)	SLAM ≥7 (n=51)	p
Sex			
Male	2	7	
Female	17	44	
Age (years)	35 (19-63)	30 (14-60)	0.22
Duration of disease (years)	2.5 (0.33-18)	1.33 (0.15-20)	0.16
Hemoglobin (g/dL)	12.6 (8.93-14.6)	9.50 (2.90-13.9)	0.15 [^]
Leukocytes (x10 ³ μ/L)	7.23 (3.73-16.38)	6.40 (1.02-17.04)	0.96
Neutrophils (x10 ³ μ/L)	4.76 (11.98)	4.92 (0.48-15.52)	0.41
Lymphocytes (x10 ³ μ/L)	1.58 (0.66-2.73)	0.95 (0.14-3.49)	0.01
Platelets (x10 ³ μ/L)	268 (175-370)	220 (4.68-687)	0.39
Random plasma glucose (mg/dL)	87 (73-212)	101 (62-232)	0.40
BUN [†] (mg/dL)	12 (7.0-23.3)	13 (4-107)	0.73 [^]
Creatinine serum (mg/dL)	0.91 (0.39-1.32)	0.85 (0.32-13.6)	0.08
Albumin (g/dL)	4.10 (2.0)	3.34 (1.64-4.35)	0.07 [^]
ESR ^{**} (mm/hour)	23.5 (11-82)	47 (9-141)	0.41
CRP ^{***} (mg/dL)	0.70 (0.1-3.3)	4.67 (0.1-194)	0.00
NLR	2.87 (0.14-9.29)	4.51 (0.57-25.85)	0.00
SLICC/ACR Index ^{****}	0 (0-2)	0 (0-6)	0.15
[†] : Blood Urea Nitrogen			
^{**} : Erythrocyte Sedimentation Rate			
^{***} : C-reactive protein			
^{****} : Systemic Lupus International Collaborating Clinics/American College of Rheumatology			
Damage Index			
[^] : Adjusted for Disease Activity			

Abstract 159 Tabel 2 Bivariate analysis between several parameters and SLAM score

Parameter	SLAM Score	p
NLR	r: 0.249 [*]	0.04
Age	r:-0.224	0.06
Duration of disease	r:-0.295 [*]	0.01
ESR	r:0.443 [*]	0.00
CRP	r:0.319	0.06
[*] : Significant correlation based on Spearman's Correlation Test (two tail hypothesis, p<0.05)		

p<0.01; 2.873 (0.14–9.29) vs 4.51 (0.57–25.85), p<0.01]. Positive correlation was observed between NLR and SLAM score (r=0.249, p<0.038) with Spearman's correlation test. Analysis with ROC curve revealed the cut-off value of NLR was 3.17 (AUC 0.684, p<0.02, 95% CI 0.55–0.813, sensitivity 69%, specificity 58%).

Conclusions NLR is correlated to disease activity and a promising tool for assessing disease activity in Indonesian SLE patients.

160

DYSLIPIDEMIA AND DISEASE ACTIVITY IN SYSTEMIC LUPUS ERYTHEMATOSUS: AN INDEPENDENT RISK FACTOR

H Gunawan*, A Awalia, J Soeroso. *Airlangga University, Internal Medicine Department – Rheumatology Division, Surabaya, Indonesia*

10.1136/lupus-2017-000215.160

Background and aims Systemic Lupus Erythematosus (SLE) patients have been associated with increased cardiovascular morbidity and mortality due to atherosclerosis. Dyslipidemia, a traditional atherosclerosis risk factor, has been reported as a long-term independent risk factor in SLE patients. Patients

Results There were 21 patients with SLAM score <7 (group 1) and 52 patients with SLAM score ≥ 7 (group 2). Dyslipidemia was more prominent in patients with 0–1 year's duration group (66.7%). SLAM Score was positively correlated with triglycerides ($r: 0.32, p < 0.00$) and negatively correlated with HDL ($r: -0.57, p < 0.00$). Analysis with multivariate backward linear regression revealed that HDL and total cholesterol were the independent risk factors influencing disease activity [$F(2,66)=19.5, R^2=0.35, p < 0.00$].

Parameter	SLAM <7 (n= 21)	SLAM ≥7 (n=52)	p [*]
Sex			
Male	2	7	
Female	19	45	
Age (years old)	35 (19-63)	30.5 (14-60)	0.25
Systole (mmHg)	120 (90-170)	120 (80-180)	0.84
Diastole (mmHg)	80 (58-100)	70 (50-100)	0.56
Hemoglobin (g/dL)	12.6 (8.93-14.6)	9.55 (2.9-13.9)	0.17 [*]
Leukocytes (x10 ³ μ/L)	7.23 (3.73-16.38)	6.40 (1.02-17.04)	0.96
Platelets (x10 ³ μ/L)	268 (122-370)	223.5 (4.68-687)	0.41
Random plasma glucose (mg/dL)	87 (73-212)	101(62-232)	0.46
BUN [†] (mg/dL)	12 (6-29)	13 (4-107)	0.50 [*]
Creatinine serum (mg/dL)	0.91 (0.39-1.32)	0.81 (0.32-13.6)	0.08
Albumin (g/dL)	4.10 (2.4-4.4)	3.33 (1.64-4.35)	0.08 [*]
Total Cholesterol (mg/dL)	174 (120-313)	196 (85-427)	0.40
Triglycerides (mg/dL)	126.5 (71-444)	163 (66-777)	0.07
HDL [*] Cholesterol (mg/dL)	53.5 (27-88)	37 (8-71)	0.00
LDL ^{**} Cholesterol (mg/dL)	95.5 (58-159)	108 (35-235)	0.31
ESR ^{††} (mm/hour)	25.5 (11-82)	47 (9-141)	0.41
CRP ^{†††} (mg/dL)	0.7 (0.1-3.3)	4.67 (0.1-194)	0.00
SLICC/ACR Index ^{††††}	0 (0-2)	0 (0-6)	0.15

[^]: Adjusted for Disease Activity

Normal P-P Plot of Regression Standardized Residual

Abstract 160 Figure 1 Normal p-p-plot of regression standardized residual with dependent variable

Conclusions: There is a positive correlation between dyslipidemia and SLE disease activity. Total cholesterol and HDL are independent predictor influencing disease activity. Therefore, we recommend lipid profile as a routine examination in SLE patients.

161

PULMONARY DISORDERS IN LUPUS PATIENT WITH NEPHRITIS

¹Q Guo*, ²Q Hui, ¹S Nan. ¹Renji Hospital- School of Medicine- Shanghai JiaoTong University, Shanghai Institute of Rheumatology, Shanghai, China; ²Renji Hospital- School of Medicine- Shanghai JiaoTong University, Respiratory, Shanghai, China

10.1136/lupus-2017-000215.161

Background and aims To analyse the association between renal histopathological features and chest computed tomography (CT) findings in lupus nephritis (LN) patients.

Methods We reviewed the medical records and chest thin-section CT findings of 152 patients with an established diagnosis of LN based on renal biopsy and 93 systemic lupus erythematosus (SLE) patients without LN between April 2009 and March 2012. The CT images were retrospectively evaluated by an experienced thoracic radiologist without knowledge of the patients' clinical information.

Results Lupus nephritis patients have a significantly higher incidence of lung/plural disease than those without LN. The patients in LN group were more prone to ground glass opacity, interlobular septal thickening, reticular opacities, pleural effusions, and consolidation on CT images than in non-LN group. Class I, III, and IV lupus nephritis were associated with traction bronchiectasis, ground glass opacity, and pleural effusions, respectively. The presence of cord on chest CT scans was significantly associated with renal interstitial lesion and interstitial inflammation/fibrosis. Ground glass opacity and reticular opacities on chest CT were related to renal hyaline thrombi. There was a significant association between pleural effusions and cellular/fibrous crescents, interstitial lesion, or interstitial inflammation/fibrosis. Hyaline thrombi in renal biopsy was an independent risk factor of the presence of ground glass opacity on CTs with logistic regression analysis.

Conclusions There are some relation between lung and renal disorders in lupus patient. LN patients were more likely to suffer from lung/pleural disease. The patients with hyaline thrombi in renal biopsy were more prone to have ground glass opacity on CTs.

Abstract 160 Table 2 Basic characteristics of the study based on duration illness

Parameter	0-1 Year (n=27)	1-5 years (n=29)	>5 years (n=17)	p
Dyslipidemia (n)	18 (66.7%)	17 (58.6%)	9 (52.9%)	0.54*
Total Cholesterol (mg/dL)	196.5 (104-427)	164 (85-309)	206 (120-346)	0.16 [^]
Triglycerides (mg/dL)	184 (70-777)	137.5 (66-453)	160 (66-444)	0.48 [^]
HDL Cholesterol	36 (8-76)	43 (19-70)	49 (20-88)	0.08 [^]
LDL Cholesterol	114 (36-199)	95.5 (35-204)	108 (64-235)	0.28 [^]
Use of corticosteroid (n)	23 (85.5%)	29 (100%)	16 (94.1%)	-
Corticosteroid dose (mg/day)	24 (4-50)	4 (2-48)	4 (2-16)	0.00 [^]
Use of Chloroquine (n)	14 (51.9%)	11 (37.9%)	5 (29.4%)	-
Use of Azathioprine (n)	9 (33.3%)	5 (17.2%)	4 (23.5%)	-
Azathioprine dose (mg/day)	100 (50-100)	100 (50-100)	100 (50-100)	0.58 [^]
Use of Cyclosporine (n)	4 (14.8%)	14 (48.3%)	7 (41.2%)	-
Cyclosporine dose (mg/day)	100 (50-200)	100 (50-200)	100 (50-100)	0.56 [^]
SLAM Score	14 (2-31)	7 (0-25)	9 (1-19)	0.02 [^]
SLICC/ACR Score	1 (0-6)	0 (0-2)	0 (0-5)	0.01 [^]

[^]: Statistically significant if p<0.05; two-tail hypothesis using Kruskal Wallis Test

*: Statistically significant if p<0.05; two-tail hypothesis using Chi-square Test