investigate and visualise characteristic marker prevalence and co-prevalence patterns.

Results Based on the individual marker pattern, patients can often be stratified belonging to different study subgroups. For example, for SLE we show that different reactivity groups exist including patients with different disease activity scores and organ damage patterns.

Conclusions We conclude that the approach of a comprehensive prevalence and signature analysis and a vivid data visualisation is useful for any multiplex omics assay.

PS1:21 URINARY MARKERS OF INFLAMMATION IN LUPUS NEPHRITIS PATIENTS

J Kosalka, B Jakiela, J Musial. Jagiellonian University Medical College, Department of Medicine, Krakow, Poland

10.1136/lupus-2018-abstract.69

Objectives Renal involvement is the most important manifestation of systemic lupus erythematosus, but assessing of inflammatory response in kidneys with non-invasive methods is still challenging. In this study we aimed to define markers of active lupus nephritis (LN) using urine immune profiling.

Methods Levels of cytokines (18-plex array) and mRNA expression (40 immune and glomerular injury genes) were measured in urine samples of LN patients with active disease (n=17), during remission (n=16), and in healthy subjects (n=19).

Results Urine levels of CCL2, CCL5, CXCL10 and IL-6 were elevated in active LN as compared to remission (best discrimination for CCL2), and correlated with LN activity. In the active disease, urinary cell transcriptome showed strong upregulation of proinflammatory cytokines (e.g. TNF, CCL2, CCL5, CXCL10), Th1 related genes (e.g. CD3G, CD4, TBX21, IFNG), and markers of glomerular damage (NPHS2 [podocin]). Active pattern of gene expression was also observed in 5 patients in remission, who had moderately increased urinary leukocyte count, two patients from this group (40%) developed renal exacerbation during following 3 months. Markers of Th17 immune axis (e.g. IL-17A) were not significantly increased in active LN.

Conclusions Active LN patients (also patients at risk of exacerbation) were characterised by marked increase of proinflammatory mediators in the urine. We identified CCL2 chemokine as the most promising marker for monitoring of disease flare.

PS1:22 USE OF INTERFERON ALPHA AND INTERLEUKIN-10 AS CLINICAL ACTIVITY BIOMARKERS IN SYSTEMIC LUPUS ERYTHEMATOUS PATIENTS

¹E Grau, ²M Fernandez Matilla, ¹CM Feced Olmos, ¹E Labrador Sanchez, ¹FM Ortiz Sanjuan, ²N Fernandez-Llanio Cornella, ¹I Chalmeta Verdejo, ¹K Arevalo Ruales, ¹R Negueroles Albuixech, ¹J Ivorra Cortes, ¹JJ Frago Gil, ¹I Martinez Cordellat, ¹R Gonzalez Mazario, ¹L Gonzalez Puig, ¹C Alcañiz Escandell, ¹C Najera Herranz, ¹I Canovas Olmos, ¹E Vicens Bernabeu, ¹JE Oller Rodriguez, ¹M de la Rubia Navarro, ²JA Castellano Cuesta, ³V Fornes Ferrer, ³D Hervas Marin, ¹JA Roman Ivorra. ¹Rheumatology Department. HUP La Fe, Valencia, Spain; ²Rheumatology Section. Hospital Arnau de Vilanova, Valencia, Spain; ³Biostatistics Unit. IIS La Fe, Valencia, Spain

10.1136/lupus-2018-abstract.70

Purpose To analyse the association among INF1A, IL10 and BLYS levels and clinical activity in SLE.

Methods A cross-sectional, observational study of 142 patients diagnosed of SLE according to SLICC 2012 criteria and 34 healthy controls was performed. In patients a complete blood-test was made, and clinical data by personal interview was collected. We analysed the serum concentration of IL10, BLYS and INF1A by colorimetric methods. SLE patients were dichotomized as high and low levels for each cytokine based on the cytokine level above 2 SD of the mean in healthy controls. Biostatistical analysis with R (3.3.2.) was performed.

Results In our SLE patients we observed higher values of IL10, BLYS and INF1A than healthy controls ($p<0.001$, $p=0.005$ and $p=0.043$ respectively), showing an average values in patients of 13.39 ± 27.73 pg/mL INF1A, 9.99 ± 15.84 pg/mL IL10 and 1811.31 ± 1757.81 pg/mL BLYS. The mean clinical activity measured by SLEDAI was 5.91 ± 5.06 .

Statistical analysis indicate that INF1A levels are correlated to IL10 levels ($p=0.001$) and BLYS levels ($p=0.034$). Due to this finding, we categorised SLE patients by low or high level of the three cytokines: 44 INF1A(-)IL10(-)BLYS(-); 61 INF1A(+)IL10(-)BLYS(-); 5 INF1A(+)IL10(-)BLYS(+); 18 INF1A(+)IL10(+)BLYS(-) and 14 INF1A(+)IL10(+)BLYS(+). There is a high association of increased IL10-INF1A levels and the increased of clinical activity measured by SLEDAI score ($p<0.0001$), and to a lesser extent with increased IL10-INF1A-BLYS levels. Patients with high IL10-INF1A and IL10-INF1A-BLYS showed a significant rise in C3-C4 consumption ($p<0.001$ and $p=0.001$ respectively) and high anti-dsDNA ($p=0.001$ and $p=0.002$ respectively). Patients with increased INF1A-BLYS exhibited high anti-dsDNA ($p=0.004$) and ENA positivity ($p<0.001$). In addition, patients with increased levels of IL10-INF1A-BLYS showed ANAs ($p<0.001$) and antiphospholipid autoantibody positivity ($p=0.004$).

Conclusions The 69% of our SLE patients displayed almost one cytokine increased, being the INF1A the cytokine that mainly is increased. However, increased IL10 levels, irrespective of whether there is also increased levels of BLYS and/or INF1A, is the cytokine which best fits to clinical activity in SLE.

Poster session 2: Autoantibodies, biomarkers and imaging (2), Environmental, epigenetics and genomics

PS2:23 ANTI-C1Q ANTIBODIES IN TURKISH SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS

S Tekeoglu, D Temiz Karadag, O Ozdemir Isik, A Yazici, A Cefle. Kocaeli University, Internal Medicine, Rheumatology Department, Kocaeli, Turkey

10.1136/lupus-2018-abstract.71

Purpose Anti-C1q has been shown to be associated with systemic lupus erythematosus (SLE) and disease activity of lupus nephritis in previous studies. We studied anti-C1q specificity for SLE versus rheumatic disease controls and healthy controls and the association with SLE manifestations in a single centre cross-sectional study.

Methods Demographics, disease information and blood samples were obtained during routine follow-up visits of patients attending Kocaeli University rheumatology outpatient clinic. There were 150 SLE patients (92% female, mean: 46 years). Control group had 85 rheumatoid arthritis patients, 16

Abstract PS2:23 Table 1 Association between clinics, serology and Anti-C1Q in SLE

Clinics & Serology	Anti-C1Q (+) (%)	Anti-C1Q (-) (%)	p value	Odds ratio (95% CI)	Adjusted p value for age
Photosensitivity	90.4	16.1	0.63	1.4 (0.3, 6.8)	0.01
Malar rash	56.8	56.6	0.70	0.7 (0.1, 3.3)	0.06
Discoid rash	0	100	0.99	-	0.40
Alopecia	0	100	0.58	1.6 (0.2, 8.9)	0.5
Oral ulcer	0	100	0.64	1.6 (0.1, 13.9)	0
Pleuritis	0	100	0.99	-	0.16
Pericarditis	0	100	0.99	-	0.12
Arthritis	97	8.2	0.15	2.9 (0.65, 13.6)	0.14
Renal disease	33.3	73.1	0.20	0.3 (0.06, 1.8)	0.21
Neurologic disease	0	100	0.99	-	0.46
Leukopenia	91.5	28.6	0.81	0.8 (0.1, 4.4)	0.005
Hemolytic anemia	0	100	0.99	-	0.52
Thrombocytopenia	0	100	0.87	-	0.2
Anti-dsDNA	89.1	13.8	0.67	0.6 (0.1, 3.7)	0.034
Anti-Smith	0	100	0.97	1.0 (0.1, 9.3)	0.73
Low complement	72.3	52.2	0.61	0.6 (0.1, 3.6)	0
Anti-phospholipid antibody	0	100	0.40	2.4 (0.2, 20.7)	0
Direct Coombs (+)	0	100	0.39	0.5 (0.1, 2.3)	0

patients with other diseases (Sjogrens's syndrome, systemic sclerosis, adult onset Still disease, psoriatic arthritis) and 49 healthy persons (88% female, mean: 45 years). Anti-C1q was measured by ELISA according to manufacturer's instructions.

Results In SLE group 72 patients had renal, 10 patients had neurologic disease, 25 patients had antiphospholipid syndrome (APS). Anti-dsDNA was positive in 92 patients, anti-Sm was positive in 18 patients, 83 patients had low complement levels. Prevalence of anti-C1q was 5% (8/150) in patients with SLE and 1% (2/150) in controls ($p=0.88$). There was no correlation between clinics (renal, neurologic, hematologic, mucocutaneous disease, arthritis, serositis, APS) or laboratory findings (Anti-dsDNA, anti-Sm, low complement levels, direct Coombs' test). Within anti-C1q positive group 6 patients had renal disease, 4 had SLEDAI scores 4 or more. None of them had nervous system disease. Six patients had leukopenia, 2 had thrombocytopenia, 5 had mucocutaneous disease. Laboratory findings were as follows; 6 patients had anti-dsDNA, 1 had anti-Sm antibody, 6 had low complement levels.

Conclusions Even though there were studies showing the relationship between anti-C1q antibodies and renal disease, it was shown that most of the patients with antibody positivity were Asian ethnicity and younger than 30 years of age. Furthermore anti-C1q antibody was related to disease activation and eliminated within 3 months of treatment. There were also different commercial products available with different cut-off levels. Meta-analysis have shown although anti-C1q antibodies

are associated with lupus nephritis, the post-test probabilities are not sufficient to provide certainty of the presence or absence of history of disease.

PS2:24 PLASMATIC AND URINARY ENDOTHELIAL MICROPARTICLES ARE INCREASED IN PATIENTS WITH LUPUS NEPHRITIS

F Miranda, C Barbati, S Truglia, FR Spinelli, F Ceccarelli, C Alessandri, G Valesini, F Conti. Dipartimento di Medicina Interna e Specialità Mediche, Reumatologia, Sapienza Università di Roma, Italy

10.1136/lupus-2018-abstract.72

Systemic lupus erythematosus(SLE) is a chronic autoimmune disease,characterised by alterations in both the innate and adaptive immune system ultimately leading to the loss of immunologic tolerance and occurrence of autoantibodies against nuclear material. Lupus nephritis is one of the most severe features of SLE determining an increase in morbidity e mortality rates.Renal biopsy still represent a fundamental diagnostic and prognostic tool for LN.Therefore, non-invasive surrogate biomarkers of active LN are urgently needed. Circulating, heterogeneous subcellular microparticles(MPs) are released from cells and platelets constitutively and upon cellular activation or apoptosis.Such MPs may reflect the state of