PS4:68

CAROTID AND FEMORAL ATHEROSCLEROSIS IN ANTIPHOSPHOLIPID SYNDROME: EQUIVALENT RISK WITH DIABETES MELLITUS IN A CASE-CONTROL STUDY

M Tektonidou, E Krawariti, G Konstantonis, N Tentolouris, PP Sfikakis. *National and Kapodistrian University of Athens – 1st Department of Propaedeutic and Internal Medicine, Athens, Greece*

10.1136/lupus-2018-abstract.114

Background Antiphospholipid syndrome (APS) may carry a worse prognosis for vascular complications when co-existing with subclinical atherosclerosis, however, the association between the two conditions remains ambiguous.

Methods We evaluated ultrasonographic markers of subclinical atherosclerosis in carotid and femoral arteries of 86 patients with thrombotic APS (43 primary APS (PAPS), 43 systemic lupus erythematosus-associated APS [SLE/APS]), 86 patients with diabetes mellitus (DM) and 86 healthy controls, individually matched for age and gender, and investigated their associations with traditional and disease-related factors in APS.

Results Carotid plaques were found in 28% of PAPS, 23% of SLE/APS, and 30% of DM patients versus 9% of controls ($p=0.006$). Femoral plaques were found in 33% of PAPS, 19% of SLE/APS, 20% of DM, and 9% of controls ($p=0.032$). Multivariate regression-derived relative risk estimates for atherosclerotic plaques in any location were 2.72 for PAPS, 2.63 for SLE/APS, and 1.98 for DM ($p=0.004$, 0.009, 0.032 respectively), after adjusting for age, gender, hypertension, dyslipidemia, smoking, BMI, and family history of coronary disease. Among patients with APS, atherosclerotic plaques were associated with the number of traditional CVD risk factors in both PAPS ($RR=2.75$, $p<0.001$) and SLE/APS ($RR=1.84$, $p<0.001$), and with IgG anti-beta2-glycoprotein I antibodies in SLE/APS.

Conclusions Patients with PAPS and SLE/APS have a nearly 2.5-fold risk of atherosclerotic plaques in carotid and femoral arteries compared to healthy controls, similar to DM patients. Atherosclerotic plaques are associated with the number of traditional risk factors in both APS and SLE/APS, and with IgG anti-beta2-glycoprotein I antibodies in SLE/APS.

Abstract PS4:68 Table 1 Baseline characteristics of study participants and vascular ultrasonographic findings by disease group

	PAPS n=43	APS, n=86 SLE/APS n=43	DM n=86	Controls n=86
Baseline characteristics				
Age (mean±SD)	46.4±12.6	46.4±10.9	46.7±12.0	46.4±12.1
Gender (% female)	63	84	73	73
Disease duration [median (Q1, Q3)]	7 (2, 15)	11 (4, 19)	14 (6, 21)	
Hypertension (%)	32	26	37	28
Dyslipidemia (%)	23	21	37	15
Smoking (% ever)	56	58	59	42
Smoking (pack years)	14±18	12±19	16±22	7±13
BMI (kg/m ²)	29.1±4.8	26.7±6.2	29.6±8.2	26.7±5.1
Family history of premature CAD (%)	7	7	14	12
Number of traditional CV risk factors except DM (%)				
- 0-1	23	49	40	53
- ≥2	77	51	60	46
Anti-cardiolipin IgG positivity (%)	67	79		
Anti-cardiolipin IgM positivity (%)	53	60		
Anti-β2-glycoprotein I IgG positivity (%)	56	47		
Anti-β2-glycoprotein I IgM positivity (%)	44	44		
Lupus anticoagulant positivity (%)	74	72		
Triple antiphospholipid antibody positivity (%)	44	51		
Statins current (%)	21	12	30	11
Antiplatelet agents, current (%)	33	47	14	0
Systemic anticoagulation, current (%)	91	79	0	2*
Hydroxychloroquine, current (%)	21	63	-	-
Corticosteroids, current (%)	12	65	-	-
Immunosuppressives, current (%)	14	47	-	-
Vascular Ultrasound Findings				
LCCA IMT (mean±SD)	0.669±0.119	0.627±0.147	0.704±0.177	0.655±0.139
LCCA IMT ≥ 0.9 mm (%)	5	8	31	23
Atherosclerotic plaques (%)				
- Carotid arteries	28	23	30	9
- Femoral arteries	33	19	20	9
- Any site	44	28	34	15

* Two of the control subjects were on anticoagulation for atrial fibrillation without CAD. APS: Antiphospholipid syndrome; DM: Diabetes Mellitus; PAPS: Primary APS; SLE/APS: Systemic Lupus Erythematosus with APS; Q1: 1st quartile; Q3: 3rd quartile; BMI: Body mass index; CAD: Coronary artery disease; CKD: Chronic kidney disease; LCCA IMT: Left Common Carotid Artery Intima-Media Thickness

Abstract PS4:68 Table 2 Relative risk of increased atherosclerotic plaque numbers conferred by participant disease group versus controls, after adjusting for the effect of traditional CV risk factors

	Relative risk*	95% Confidence Interval	p-value
Carotid and femoral plaques			
PAPS	2.72	1.37 – 5.40	0.004
SLE/APS	2.63	1.28 – 5.41	0.009
DM	1.98	1.06 – 3.67	0.032
Carotid plaques			
PAPS	2.36	1.01 – 5.52	0.047
SLE/APS	2.86	1.20 – 6.83	0.018
DM	2.87	1.36 – 6.07	0.006
Femoral plaques			
PAPS	2.44	1.14 – 5.20	0.021
SLE/APS	1.91	0.84 – 4.38	0.124
DM	1.08	0.49 – 2.37	0.844

* Estimates derived from multiple negative binomial regression controlling for the effect of age, gender, hypertension, body mass index, pack years of smoking, family history of premature coronary artery disease
PAPS: Primary Antiphospholipid Syndrome; SLE/APS: Systemic Lupus Erythematosus with Antiphospholipid Syndrome; DM: Diabetes Mellitus