Abstract PS4:76 Figure 1

PS4:77 TWO CASES OF FATAL CATASTROPHIC ANTIPHOSPHOLIPID SYNDROME

¹L Castelnovo, ²A Laria, ²A Lurati, ¹A Tamburello, ¹P Faggioli, ¹A Mazzone. ¹Asst Ovest Milanese, Legnano, Italy; ²Asst Ovest Milanese, Magenta, Italy

10.1136/lupus-2018-abstract.122

Introduction Catastrophic antiphospholipid syndrome (CAPS) is a rare and fatal condition secondary to antiphospholipid syndrome (APS) characterised by venous and/or arterial thromboembolism within a short period of time in the presence of positive AP antibodies. Thrombocytopenia is seen in >60% of cases. Here we report 2 cases of women with fatal CAPS. Case reports 1. A 55-old female affected by APS (excluded overlap SLE), treated initially with warfarin, due to previous DVT, then apixaban after a new EP episode, was admitted to our Unit for dyspnea: we found a mitral valve injury. After few days she developed a neurologic impairment with evidence of post-ischaemic alterations, likely embolics, and a sensory – motor neuropathy. We concluded for probable CAPS and started antibiotics plus high dose corticosteroids, unfractionated heparin and plasma exchange. However, we observed further worsening. The patient quickly succumbed to her illness. Case report 2. A 59-old female affected by SLE with CSN impairment and APS with thrombocytopenia and recurring DVT – EP episodes treated with low dose methylprednisolone, iloprost and heparin (warfarin suspension for severe thrombocytopenia impairment) was admitted to our Unit for appearance of diarrhoea with abdominal pain. Blood tests showed further worsening of platelet count. We started antibiotics with clinical improvement. Taking into account the patient history CT was performed, with evidence of massive arteriovenous and aortic arterial disease severe thrombotic

obstruction. Two days later she developed acute pain on the left foot with massive venous thrombosis of the left femoro-popliteo axis. We started unfractionated heparin, despite low platelet value, as rescue therapy. The patient progressively worsened with further progression of the arterial occlusions, until death, 3 days later.

Conclusion CAPS can lead to acute multiorgan failure and can be associated with infections or SLE. Early recognition is essential for effective life-saving treatment. Management of these patients is complex, especially in those with thrombocytopenia. Anticoagulation plus steroids plus plasma exchange should be the first line therapy but our two cases show that they are not always applicable and effective. New therapies, such as rituximab and eculizumab, may be options, but time is a fundamental variable.

PS4:78 WOMEN WITH SYSTEMIC LUPUS ERYTHEMATOSUS MORE OFTEN ACHIEVE PREGNANCY, AND HAVE A SHORTER TIME TO PREGNANCY COMPARED TO WOMEN WITH RHEUMATOID ARTHRITIS

^{1,2}C Gøtestam Skorpen, ³S Lydersen, ⁴I-M Gilboe, ⁵JF Skomsvoll, ^{6,7}KÅ Salvesen, ⁴Ø Palm, ⁵HSS Koksvis, ⁵B Jakobsen, ^{1,5}M Wallenius. ¹Dept. of Neuromedicine and Movement science, NTNU, Norwegian University of Science and Technology, Trondheim, Norway; ²Dept. of Rheumatology, Ålesund hospital, Ålesund, Norway; ³Regional Centre for Child and Youth Mental Health and Child Welfare, NTNU, Norwegian University of Science and Technology, Trondheim, Norway; ⁴Dept. of Rheumatology, Oslo University Hospital Rikshospitalet, Oslo, Norway; ⁵Norwegian National Advisory Unit on Pregnancy and Rheumatic Diseases, St Olavs hospital, Trondheim University Hospital, Trondheim, Norway; ⁶Dept. of laboratory medicine, Women and Children Health, NTNU, Norwegian University of Science and Technology, Trondheim, Norway; ⁷Dept. of Obstetrics and Gynaecology, St Olavs hospital, Trondheim University Hospital, Trondheim, Norway

10.1136/lupus-2018-abstract.123

Abstract PS4:78 Figure 1 Pregnancy outcome (a) and Kaplan-Meier plot (b) with TTP as dependent variable and achieved pregnancy as event, comparing woman with SLE or RA

Objectives To examine possible differences in the ability to get pregnant and time to pregnancy (TTP) in women with systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA).

Methods Data from RevNatus, a Norwegian nationwide observational register of women with rheumatic diseases planning pregnancy was used. We compared the number of women with SLE and RA achieving pregnancy, TTP and pregnancy outcome (live birth or pregnancy loss). Fifty-three women with SLE and 180 women with RA with a pregnancy wish had follow-up until pregnancy and known pregnancy outcome or at least one year if not achieving pregnancy. With cox regression we adjusted for maternal age, parity and use of disease-modifying antirheumatic drugs (DMARDs).

Results The two groups were similar concerning maternal age, parity, smoking, BMI, educational level, prior pregnancy loss and prior preeclampsia and/or preterm birth. Women with SLE had longer disease duration ($p=0.001$), more often active disease ($p=0.002$) and more often used a DMARD ($p<0.001$). Live birth occurred in 38 (71.7%) women with SLE and 104 (57.5%) women with RA, while pregnancy loss was experienced in 9 (17.0%) and 26 (14.4%), respectively. Six (11.3%) SLE-women and 50 (27.8%) RA-women did not achieve pregnancy during follow-up (Figure 1a). Women with SLE had almost doubled pregnancy rate compared to women with RA (pregnancy ratio 1.91, CI 1.27 to 2.88, $p=0.002$), and when succeeding they also had a substantially shorter mean TTP (3.9 vs 6.2 months, $p=0.017$) (Figure 1b). Sixty-five (65/180, 36.1%) RA-women and eight (8/53, 15.1%) SLE-women had a TTP (or censoring) exceeding one year ($p=0.006$), indicating subfertility.

Conclusions Women with SLE more often achieve pregnancy and have shorter time to pregnancy compared to women with RA.

PS4:79

LONG-TERM FOLLOW-UP OF 320 CHILDREN BORN TO MOTHERS WITH SYSTEMIC AUTOIMMUNE DISEASES: A MULTICENTRE SURVEY FROM 24 RHEUMATOLOGY CENTRES IN ITALY

¹MG Lazzaroni, ¹C Nalli, ¹L Andreoli, ¹C Carini, ¹F Dall'Ara, ²M Rodrigues, ³E Bartoloni Bocci, ³R Gerli, ⁴CB Chighizola, ⁴M Gerosa, ⁴PL Meroni, ⁴L Sinigaglia, ⁵P Conigliaro, ⁵R Perricone, ⁶A Corrado, ⁶F Cantatore, ⁷S D'Angelo, ⁸M Favaro, ⁸M Larosa, ⁸A Doria, ⁸A Ruffatti, ⁹E Generali, ⁹C Selmi, ¹⁰M Meroni, ¹⁰M Cutolo, ¹¹M Padovan, ¹¹M Govoni, ¹²G Pazzola, ¹²C Salvarani, ¹³S Peccatori, ¹⁴I Prevete, ¹⁴G Minisola, ¹⁴GD Sebastiani, ¹⁵A Brucato, ¹⁵V Ramoni, ¹⁶R Caporali, ¹⁶C Montecucco, ¹⁷C Tani, ¹⁷V Signorini, ¹⁷M Mosca, ¹⁸M Trevisani, ¹⁸N Malavolta, ¹⁹M Vadacca, ¹⁹A Afeltra, ²⁰E Vivaldelli, ²⁰A Maier, ²¹E Visalli, ²¹R Foti, ²²L Zuliani, ²²A Gabrielli, ²³C Campochiaro, ²³E Baldissera, ²³MG Sabbadini, ²⁴N Romeo, ¹A Tincani. ¹University and Spedali Civili di Brescia, Italy; ²Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal; ³Azienda Ospedaliera di Perugia, Italy; ⁴Istituto Ortopedico Gaetano Pini, Milano, Italy; ⁵Policlinico Tor Vergata, Roma, Italy; ⁶Ospedali Riuniti di Foggia, Foggia, Italy; ⁷Ospedale San Carlo, Potenza, Italy; ⁸University and Azienda Ospedaliera di Padova, Italy; ⁹Humanitas, Milano, Italy; ¹⁰University of Genova, IRCCS San Martino, Genova, Italy; ¹¹University and Azienda Ospedaliera di Ferrara, Italy; ¹²Azienda Ospedaliera ASMN, Istituto di Ricovero e Cura a Carattere Scientifico, Reggio-Emilia, Italy; ¹³Azienda Provinciale Servizi Sanitari, Trento, Italy; ¹⁴A.O. San Camillo, Roma, Italy; ¹⁵Ospedale Papa Giovanni XXXIII, Bergamo, Italy; ¹⁶University and Policlinico San Matteo di Pavia, Italy; ¹⁷University of Pisa, Italy; ¹⁸Policlinico S.Orsola-Malpighi, Bologna, Italy; ¹⁹University Campus Biomedico, Roma, Italy; ²⁰Ospedale di Bolzano, Italy; ²¹Policlinico Vittorio Emanuele-Ferraro-Bambino Gesù, Catania, Italy; ²²Ospedali Riuniti di Ancona, Italy; ²³Ospedale San Raffaele, Milano, Italy; ²⁴Ospedale S. Croce e Carle, Cuneo, Italy

10.1136/lupus-2018-abstract.124

Background Rheumatic Diseases (RD) frequently affect women during reproductive age, therefore counselling on family planning is crucial for their quality of life. Children's outcome is a major topic, but no large studies are available. This study aimed at assessing the long-term health conditions of children born to women with RD.

Methods 24 Italian Rheumatology Centres distributed the questionnaire (65 multiple-choice and 12 open-answer