Abstract CS-30 Table 1 Significant independent predictors of sleep disturbances

	Obstructive Sleep Apnea (OSA)* diagnosis	Restless Leg Syndrome (RLS)* symptoms	Sleep Problems Index I (SPI-I)†
Age, years	1.04 (1.01, 1.07)	(ns)	−0.2 (0.01)
White race	(ns)	(ns)	−6.4 (0.02)
Low education (<high school)	(ns)	(ns)	6.4 (0.004)
Smoking, ever	(ns)	(ns)	4.6 (0.03)
Obesity	4.6 (2.4, 8.8)	(ns)	(ns)
Asthma	2.5 (1.1, 5.7)	(ns)	(ns)
RDCI (0–9)	1.3 (1.1, 1.5)	1.3 (1.1, 1.5)	1.3 (0.02)
Pain rating (0–10)	(ns)	(ns)	1.5 (0.002)
Disease activity (SLAQ)	1.07 (1.01, 1.13)	1.1 (1.03, 1.13)	0.9 (<0.0001)

* Tabled values are odds ratio (95% CI) from multiple logistic regression analyses

† Tabled values are beta (p-value) from multiple linear regression analysis. Higher scores reflect greater sleep problems

All regression models include age, race, education, ever smoking, obesity (BMI ≥30), concurrent asthma, Rheumatic Disease Comorbidity Index (RDCI), prednisone dose, pain rating, disease damage (BILD), and disease activity (SLAQ)

Research in SLE has linked SDs to worse outcomes. Previous research in other conditions suggests that SDs might also be a cause of increased disease activity through heightened inflammation. Further research is needed to tease out disease-specific causes and effects of SD in SLE.

REFERENCES

- England BR. *Arthritis Care Res* 2015;6:865.
- Karlson EW. *Lupus* 2003;12:280.
- Yazdany J. *Arthritis Care Res* 2011;63:1170.

CS-31 SAFETY OF HYDROXYCHLOROQUINE WITHDRAWAL IN OLDER ADULTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS

¹Peter Izmirly*, ¹Nicole Bornkamp, ¹Anna Zeron, ¹Chung-E Tseng, ¹H Michael Belmont, ²Anca Askanase, ³Jane E Salmon, ³Michael D Lockshin, ¹Jill P Buyon. ¹New York University School of Medicine, USA; ²Columbia University College of Physicians and Surgeons, USA; ³Hospital for Special Surgery, USA

10.1136/lupus-2018-lsm.66

Background Although hydroxychloroquine (HCQ) is a mainstay of treatment for patients with Systemic Lupus Erythematosus (SLE), ocular toxicity can result from accumulated exposure. The introduction of highly sensitive tools has engendered even more concern. As the longevity of patients with SLE improves, additional data will help physicians accurately balance the risk of ocular toxicity and the risk of disease flare, especially in older patients who have stable/quiescent disease. Accordingly, this study was initiated to examine the safety of HCQ withdrawal in older SLE patients.

Methods Data were obtained by retrospective chart review at three lupus centers. Twenty-seven patients met the following inclusion criteria: ≥4 ACR criteria, disease duration ≥5 years, HCQ use of 200–400 mg per day ≥5 years, and discontinuation of hydroxychloroquine at ≥age 55 years. The comparator group comprised 39 age, gender and racial/ethnic matched patients who remained on HCQ. The primary outcome was a clinically meaningful flare within one year of HCQ withdrawal, defined as

moderate or severe, using a revised version of the SELENA-SLE-DAI Flare composite that separates mild from moderate flares, evaluates each organ system separately, and incorporates increases in corticosteroid dose and/or addition of immunosuppressive agents. Mild flares were considered secondary outcomes.

Results Demographics are provided in table 1. There was a trend toward longer disease duration in the HCQ withdrawal group but no difference in prevalence of prior lupus nephritis between the groups. The reasons for HCQ withdrawal were maculopathy (n=13), presumed/biopsy proven cardiomyopathy (n=2), patient request (n=4), and miscellaneous other reasons (n=8). There was no difference in the primary or secondary outcomes between the groups (table 1). Two patients had a moderate flare after discontinuing HCQ, of whom one had arthritis treated with methotrexate and one had thrombocytopenia (>30K) and proteinuria of 2 grams/d (baseline 700 mg). Three patients had severe flares while continuing HCQ, of whom two were hospitalized, one for seizures and one for pericarditis; the third had worsening nephritis (urinary protein >4 g/d, requiring treatment). Two patients had moderate flares while remaining on HCQ, one of whom had a rash and arthritis treated with tofacitinib and one a rash treated with prednisone.

Conclusions In this retrospective study of older patients with SLE on long-term HCQ, withdrawal did not increase the risk of moderate or severe flares. These data provide reassurance regarding the safety of withdrawing HCQ in stable older SLE patients.

Abstract CS-31 Table 1

	HCQ Withdrawal (n=27)	HCQ Continuation (n=39)	P value
Age	59.9	60.6	0.52
Gender (% Female)	92.6%	97.4%	0.56
Race/Ethnicity			0.78
White	33.3%	35.9%	
Black	29.6%	25.6%	
Hispanic	18.5%	20.5%	
Asian	18.5%	17.9%	
Duration of SLE (years) (n=61)	26.7	21.1	0.088
Duration of HCQ Use (years) (n=52)	19.3	17.6	
Prior Renal Disease (N,%)	12 (44.4%)	13 (33.3%)	0.44
Moderate/Severe Flares (N,%)	2 (7.4%)	5 (12.8)	0.69
Moderate Flare	2 (7.4%)	2 (5.1%)	
Severe Flare	0 (0%)	3 (7.7%)	
Mild Flare	5 (18.5%)	3 (7.7%)	0.26
All Flares	7 (25.9%)	8 (20.5%)	0.77

CS-32 SLE-YPLL (YEARS OF POTENTIAL LIFE LOST) AS A MEASURE OF RELATIVE BURDEN OF PREMATURE MORTALITY

Eric Y Yen, Ram R Singh*. University of California at Los Angeles (UCLA), Los Angeles, USA

10.1136/lupus-2018-lsm.67

Background Disease burden is the impact of a health problem on a given area, which can be used to prioritize actions in health, assess performance of healthcare and disease management, identify high-risk populations, and set research

priorities. Disease burden can be measured using a variety of indicators such as mortality, morbidity or financial cost. For SLE, despite a relatively good 10 year survival (95%, vs 97% in controls), its mortality rate remains high relative to general population mortality rate as shown in our analysis of the Centers for Disease Control (CDC)'s mortality database comprising of 62,843 SLE deaths. However, mortality rates may not adequately measure SLE burden, because among those who died, a fifth of patients died before reaching 40 years of age. Premature mortality, a measure of burden first proposed in 1940s to address the inadequacy of mortality in measuring the burden of tuberculosis, is an important way to quantify disease burden. In constructing a measure of premature death, an arbitrary limit to life is chosen, and the calculation of the difference between an age at death and this arbitrary limit is defined as the life lost as a result of that death. To assess the relative burden of SLE, we measured the years of potential life lost (YPLL) for SLE relative to CDC's top 15 leading causes of death.

Methods This is a population-based study. Death counts were obtained from the CDC-WONDER database. Number of deaths were tabulated for top 15 CDC's leading causes-of-death and SLE. To calculate YPLL, each decedent's age at death from a specific disease was subtracted from a predetermined age of 75 years. The years of potential life lost were then added together to yield the total YPLL.

Results 28 411 female SLE deaths were recorded during 2000–2015. The ranking of SLE deaths relative to the CDC's official leading-causes-of-death ranklist in females showed that SLE is within the top 15 leading causes-of-death in reproductive age women (15–44 years) and tenth among women ages 15–24 years. We calculated YPLL for SLE relative to the CDC's official 15 leading causes-of-death in women ages 15–44 and 15–24. YPLL for SLE ranked 14th in women ages 15–44, and 8th in women ages 15–24, above diabetes mellitus, HIV disease, septicemia, chronic lower respiratory disease, anemias, nephritis, and cerebrovascular disease.

Conclusions SLE is among the leading cause of YPLL in young women, underscoring SLE as an important public health issue.

CS-33 PERFORMANCE OF URINARY HER2 AS A BIOMARKER OF ACTIVE PROLIFERATIVE LUPUS NEPHRITIS

¹Patricia Costa-Reis, ²Rebecca E Sadun*, ¹Kelly Maurer, ²Laura E Schanberg, ¹Jon M Burnham, ³Emily von Scheven, ⁴Kathleen M O'Neil, ⁵Marisa S Klein-Gitelman, ⁶M Petri, ¹Kathleen E Sullivan. ¹Children's Hospital of Philadelphia, Philadelphia, PA; ²Duke University, Durham, NC; ³University of California, San Francisco, San Francisco, CA; ⁴Indiana University, Indianapolis, IN; ⁵Lurie Children's Hospital, Chicago, IL; ⁶Johns Hopkins University, Baltimore, MD

10.1136/lupus-2018-lsm.68

Background Human Epidermal Growth Factor Receptor 2 (HER2) expression is increased in the kidneys of patients with proliferative lupus nephritis (LN) as well as the kidneys of the mouse model NZM2410, though it is not increased in other glomerulonephritides. Preliminary data indicated that urinary HER2 (uHER2) may be elevated in adult patients with proliferative LN (class III and class IV). This study sought to establish whether uHER2 can serve as a biomarker to predict flare of proliferative lupus nephritis in patients with systemic lupus erythematosus (SLE).

Methods We performed a multicenter prospective study of children with LN as well as a single center prospective study

of adults from Johns Hopkins; children and adults without lupus were enrolled as controls. Urine samples were collected at scheduled visits over the course of 4 years. uHER2 was measured by a commercially available ELISA and compared to two other commercially assayable prospective urinary markers of LN, TNF-like weak inducer of apoptosis (TWEAK) and vascular cell adhesion molecule-1 (VCAM-1). Active lupus nephritis was defined as the presence of one or more of the following findings: granular or red blood cell casts, >5 red blood cells/high power field with the exclusion of other causes, or proteinuria of >0.5 grams/24 hours or recent increase of >0.5 grams/24 hours.

Results One hundred children with LN were enrolled from five academic centers across the United States alongside 91 age-matched controls (mean age 15 years). Children with LN had significantly higher levels of uHER2 compared with controls ($p < 0.01$); the difference was greater for children with active LN ($p < 0.005$). uHER2/creatinine correlated with uTWEAK/creatinine ($p < 0.0001$) and with uVCAM-1/Creatinine ($p < 0.05$); uHER2 was more sensitive for flare than TWEAK and more specific for flare than VCAM-1. The adult cohort consisted of 142 SLE patients and 23 healthy controls. uHER2/creatinine levels were increased in adult SLE patients when compared to controls ($p < 0.05$) with the difference also greatest in patients with active LN ($p < 0.005$).

Conclusion This multicenter prospective study establishes uHER2 as a promising new biomarker for determination of lupus nephritis flare in both children and adults. Combined with implications for HER2 as a driver of mesangial cell proliferation through microRNA downregulation, these results provide further impetus to explore already-available HER2-inhibiting drugs, including tyrosine kinases and certain monoclonal antibodies, for the treatment of LN, such as lapatinib (Tyverb) or trastuzumab (Herceptin).

CS-34 INCREASED PREVALENCE OF NOCTURNAL HYPERTENSION AND ATTENUATED BLOOD PRESSURE DIPPING IN PEDIATRIC SLE

¹Scott E Wenderfer*, ¹J Fallon Campbell, ¹Steffy Jose, ¹Mario Archila, ²Robyn Matloff, ²Kari Mynarski, ³D Jack Weaver, ⁴Sarah Hoffman, ⁵Jessica Nelson, ⁵Mahmoud Kallash, ⁶Hermine Brunner, ⁶Mark Mitsnefes. ¹Baylor College of Medicine, Houston, TX, USA; ²New York Medical College, Valhalla, NY, USA; ³Levine Children's Hospital of Carolinas, Charlotte, NC, USA; ⁴Virginia Commonwealth University, Richmond, VA, USA; ⁵The Ohio State University, Columbus, OH, USA; ⁶Cincinnati Children's Hospital, Cincinnati, OH, USA

10.1136/lupus-2018-lsm.69

Background The optimal goals for blood pressure management in SLE are unknown. Uncontrolled blood pressure is a known risk factor for cardiovascular disease. The increased cardiovascular risks associated with SLE are not fully explained by traditional risk factors, such as hypertension and dyslipidemia, at least when based on definitions established for individuals without SLE. One possibility is that the patterns of blood pressure dysregulation in SLE may differ from those in otherwise healthy individuals or patients with obesity/metabolic syndrome.

Methods The BPpSLE study is a cross-sectional multicenter analysis of blood pressure patterns in children with SLE. Inclusion criteria: ≥ 4 ACR classification criteria for SLE and age < 21 years. Exclusion criteria: intravenous steroid use within past month, glomerular filtration rate < 60 mL/1.73 m²/min, or kidney transplant. Interim analysis was performed on