macrophages transfected with hY3. IFIT1 MX1, and EIF2AK2 transcripts were significantly increased in the WISH cells treated with hY3 macrophage supernatants, but not macrophage supernatants alone (n=7, p=0.02).

Conclusion These data now provide a link between IFN and the inflammatory and possibly fibrosing component of CHB and position Siglec-1 positive macrophages as integral to the process.

Living with Lupus

LL-01 SOCIAL DETERMINANTS OF TREATMENT ADHERENCE AND DISEASE SEVERITY AMONG PEOPLE LIVING WITH LUPUS IN A SMALL ISLAND DEVELOPING STATE: A REPORT FROM ST. LUCIA

¹Amanda King*, ¹Cleopatra Altenor, ²Catherine Brown, ²Ian Hambleton. ¹Bay Medical Centre, St. Lucia, West Indies; ²The George Alleyne Chronic Disease Research Centre, Caribbean Institute for Health Research, The University of the West Indies, Barbados, West Indies

10.1136/lupus-2018-lsm.111

Background The occurrence of systemic lupus erythematosus (SLE) varies considerably worldwide, with documented high incidence rates among women of African descent. The clinical course is likely influenced by social determinants, including

socioeconomic position (SEP), yet findings remain inconsistent, with little information from the Caribbean diaspora. This study presents the epidemiology of SLE in St. Lucia for the first time, exploring the association of SEP and SLE medication adherence and disease severity.

Methods Data have been collected from the only specialist lupus clinic in St Lucia between 1995 and 2017. We explored the effect of selected markers of SEP on disease severity (yes/no), and treatment adherence (yes/no) using logistic regression, adjusting for the effects of age, sex and years since diagnosis at all times. We used education level (primary or secondary education, tertiary education) or patients eligible for treatment cost discount or exemption (yes/no) as indicators of SEP. We also explored the effect of enrolment in a self-help programme on both regression outcomes, and the effect of treatment adherence on disease severity.

Results 143 people with SLE have been registered at the clinic between 1995 and 2017. The mean age at diagnosis was 32 years (standard deviation 12 years), and 132 (92%) were female, for a female to male ratio of 12 to 1. Since 2010 (a period of full clinic operation) 66 women have been diagnosed with SLE, for a crude incidence rate of 9.3 per 1 00 000 person years (95% CI 7.2 to 11.8). Half (49%) had a severe clinical course, defined as having cerebritis, nephritis, or being on dialysis, and half (50%) were medication adherent at their last follow-up visit. Higher SEP was consistently associated with increased treatment adherence and decreased disease severity (treatment adherence odds ratios ranged from 2.4 to 3.4; disease severity odds ratios ranged from 1.0 to 3.5) (figure 1).

Conclusion In St Lucia, among a population of predominantly African descent, and using selected markers of SEP, patients of lower socioeconomic position have more severe disease and lower medication adherence than those of higher socioeconomic position.

Acknowledgements Christina Howitt, for assistance with data analysis.

LL-02 DEPRESSION IN PATIENTS WITH CHRONIC CUTANEOUS LUPUS ERYTHEMATOSUS: PREVALENCE AND ASSOCIATED FACTORS IN A PREDOMINANTLY AFRICAN AMERICAN COHORT FROM THE SOUTHEAST U.S

¹Jennifer Hong, ²Gaobin Bao, ³Tamara Haynes, ⁴Laura Aspey, ²S Sam Lim, ²Cristina Drenkard*. ¹Emory University School of Medicine, Atlanta, GA; ²Department of Medicine, Emory University School of Medicine, Division of Rheumatology, Atlanta, GA; ³Department of Medicine and Department of Psychiatry and the Behavioral Sciences, Emory University School of Medicine, Atlanta, GA; ⁴Department of Dermatology, Emory University School of Medicine, Atlanta, GA

10.1136/lupus-2018-lsm.112

Background Chronic dermatologic diseases in general and cutaneous involvement in patients with systemic lupus erythematosus (SLE) in particular have been linked to increased depression. However, little is known about the burden of depression and its risk factors in patients with primary chronic cutaneous lupus erythematosus (CCLE), the most common type of cutaneous lupus. Additionally, previous studies examining CCLE have included predominantly white patients, despite recent findings indicating that black individuals have higher susceptibility for this condition and experience earlier damage in the disease course. We aimed to examine the prevalence of depression in patients with primary CCLE in the Southeast

Abstract LL-01 Figure 1 Odds ratios with 95% confidence intervals for potential predictors of disease severity and treatment adherence