

# Plucked hair follicles from patients with chronic discoid lupus erythematosus show a disease-specific molecular signature

Mohammad Shalbf,<sup>1</sup> Adewonuola A Alase,<sup>1</sup> Anna Berekmeri,<sup>1,2</sup> Md Yuzaiful Md Yusof,<sup>1,2</sup> Jelena Pistolic,<sup>3</sup> Mark J Goodfield,<sup>4</sup> Sara Edward,<sup>5</sup> Natalia V Botchkareva,<sup>6</sup> Martin Stacey,<sup>7</sup> Edward M Vital,<sup>1,2</sup> Miriam Wittmann<sup>1,2</sup>

**To cite:** Shalbf M, Alase AA, Berekmeri A, *et al.* Plucked hair follicles from patients with chronic discoid lupus erythematosus show a disease-specific molecular signature. *Lupus Science & Medicine* 2019;**6**:e000328. doi:10.1136/lupus-2019-000328

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/lupus-2019-000328>).

Received 21 February 2019  
Revised 26 April 2019  
Accepted 17 June 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

**Correspondence to**  
Dr Miriam Wittmann; M. Wittmann@leeds.ac.uk

## ABSTRACT

**Objective** When faced with clinical symptoms of scarring alopecia—the standard diagnostic pathway involves a scalp biopsy which is an invasive and expensive procedure. This project aimed to assess if plucked hair follicles (HFs) containing living epithelial cells can offer a non-invasive approach to diagnosing inflammatory scalp lesions.

**Methods** Lesional and non-lesional HFs were extracted from the scalp of patients with chronic discoid lupus erythematosus (CDLE), psoriasis and healthy controls. RNA was isolated from plucked anagen HFs and microarray, as well as quantitative real-time PCR was performed.

**Results** Here, we report that gene expression analysis of only a small number of HF plucked from lesional areas of the scalp is sufficient to differentiate CDLE from psoriasis lesions or healthy HF. The expression profile from CDLE HFs coincides with published profiles of CDLE from skin biopsy. Genes that were highly expressed in lesional CDLE corresponded to well-known histopathological diagnostic features of CDLE and included those related to apoptotic cell death, the interferon signature, complement components and CD8<sup>+</sup> T-cell immune responses.

**Conclusions** We therefore propose that information obtained from this non-invasive approach are sufficient to diagnose scalp lupus erythematosus. Once validated in routine clinical settings and compared with other scarring alopecias, this rapid and non-invasive approach will have great potential for paving the way for future diagnosis of inflammatory scalp lesions.

## INTRODUCTION

Cutaneous lupus erythematosus (CLE) can present as part of the SLE spectrum or as a ‘skin only’ disease.<sup>1</sup> Up to 80% of all CLE cases seen in dermatology settings are diagnosed to be chronic discoid lupus erythematosus (CDLE).<sup>2</sup> In most patients with CDLE, lesions are predominantly in sun-exposed areas including face, neck, décolletage and, importantly, the scalp with often disfiguring lesions resulting in permanent scarring and irreversible hair loss.<sup>3–5</sup> CLE in visible parts

of the body is well recognised to significantly affect the quality of life of patients.<sup>6</sup>

Although CDLE can be diagnosed by experienced dermatologists based on morphological characteristics including erythema, pigmentary disturbances, telangiectasia and atrophy, a skin biopsy is usually required to confirm the diagnosis. Histopathological and immunohistological findings include interface dermatitis<sup>7</sup> with a dense, predominantly CD8<sup>+</sup> T-cell lymphocytic infiltrate particularly around the bulge region of the hair follicle (HF), basal cell/vacuolar degeneration representing apoptotic keratinocytes, basement membrane changes, infiltrating macrophages and plasmacytoid dendritic cells, pilosebaceous destruction and atrophic scarring.<sup>4 8–10</sup> Many of these features are also seen in lichen planopilaris (LPP), a chronic inflammatory skin disease which also leads to permanent scarring alopecia.<sup>11–14</sup> However, deposition of immunoglobulins and complement component 3 (C3) at the dermoepidermal junction as seen using direct immunofluorescence (DIF) is specific for CDLE but not LPP and thus a key differential diagnostic finding. Performing a scalp biopsy for histology and DIF is an invasive, thus often stressful procedure for the patient. It also requires availability of trained doctors and nurses, a special biopsy set-up with sterile equipment, patient’s consent regarding the side effects of local anaesthetic and invasive biopsy (including infection, scar, healing problems and bleeding) and usually a separate appointment, thus further time commitment for the patient. Furthermore, the biopsy needs to be processed, analysed and communicated by a dermatohistopathologist which takes up to 4 weeks in a standard dermatology setting. Thus, the overall

costs involved to receive a conventional biopsy histology report are several hundred pounds/Euro (depending on the health system used).

A molecular hallmark of lupus erythematosus lesions is the high expression of interferon (IFN)-stimulated genes (ISGs).<sup>9 15 16</sup> These include myxovirus protein A (*MxA*),<sup>17</sup> IFN inducible protein 6 (*IFI6*), *CXCL9*, *CXCL10*, 2,5-oligoadenylate synthase (*OAS*) including *OAS2* and *OASL*, IFN-induced helicase C domain-containing protein 1 (*IFIH1/MDA5*), bone marrow stromal antigen 2 (*BST2*) and guanylate-binding protein 1 (*GBPI*) in CLE lesions.<sup>8 18-21</sup> It is well known that IFNs induce expression of major histocompatibility complex (MHC) class I and MHC class I-pathway related molecules ( $\beta$ 2 microglobulin,  $\beta$ 2M) and this has also been reported for organ-cultured human scalp skin.<sup>22-24</sup> The proposed mechanism responsible for high expression of ISG includes the impaired clearance of apoptotic material, which results in secondary necrosis and release of immunostimulatory nucleic acids.<sup>25-27</sup> An upregulated IFN response goes along with antiproliferative and antineovascularisation properties and may be involved in the downregulation of tissue repair mechanisms observed in CDLE.<sup>9 25 28</sup> However, psoriasis which often presents with strong scalp involvement but never resulting in scars is also recognised to have some activation of the IFN pathway.<sup>29</sup> In this work, we aimed to explore the potential of plucked HF analysis to identify CDLE-specific changes. We mainly focused on differentially expressed genes comparing HFs from lesional versus non-lesional areas of the scalp. Psoriasis samples were also included in our analysis, as a disease comparator which shows significant scalp inflammation with leukocyte infiltration but presents with different clinical outcomes regarding atrophy and scarring.<sup>30</sup>

## MATERIALS AND METHODS

### Patients

All patients had confirmed psoriasis as diagnosed by consultant dermatologists or CDLE (biopsy proven) and had been suffering long term (eg, longer than 5 years). All patients showed clinically active scalp lesions. Most patients with CDLE were on hydroxychloroquine but were asked to discontinue the drug for 3–5 days prior sampling. All patients with CDLE were female (age range 20–75 years of age, mean 55), 80% of healthy controls were female (mean age 46) and for patients with psoriasis, 50% were male and 50% female (mean age 43 years). Patients had not used topical corticosteroids within 48 hours prior to sampling. All individuals provided informed written consent and this research was carried out in compliance with the Declaration of Helsinki. The patients' samples used for this study were collected under ethical approval, REC 10/H1306/88, National Research Ethics Committee Yorkshire and Humber–Leeds East. All experiments were performed in accordance with relevant guidelines and regulations.

### Plucked HF samples


Four to five HFs from five healthy individuals (male and female aged 35–55 years), seven patients with CDLE from both lesional and non-lesional areas and six patients with psoriasis from both lesional and non-lesional areas were plucked out using tweezers as described previously.<sup>31</sup> Only hairs with a full, visible HF were used (eg, telogen hairs or those with incompletely plucked follicle were excluded). Time to pluck four to five anagen hairs with suitable HFs usually takes an experienced person 5 min as up to 10 hairs have to be plucked to obtain this yield. In this study, we have restricted the area of 'lesional' sampling to scalp presenting with clinical signs of inflammation, such as erythema (including mild erythema), infiltration, hyperkeratosis/scaling. The hair shaft was cut-off and the white sheath (inner and outer root sheath) containing keratinocytes were used for this study (figure 3). Trimmed HFs were then immersed in Optimal Cutting Temperature (OCT) compound (Tissue-Tek; Sakura Finetek Europe, Alphen aan den Rijn, The Netherlands) in 1.5 mL tubes to preserve RNA quality before further processing. The samples were either stored in  $-80^{\circ}\text{C}$  or processed immediately for RNA extraction.

### Total RNA extraction from HFs

Plucked HFs were embedded longitudinally in OCT and 3–5  $\mu\text{m}$  thick frozen sections were cut using a Leica CM3050S cryostat (Leica Microsystems, Buckinghamshire, UK) and collected in RNase free universal tubes. This was followed by the addition of lysis buffer (Relia-Prep RNA Cell Miniprep System; Promega, Wisconsin, USA) to the cut HF sections which were stored on dry ice prior to RNA extraction, which was carried out according to the manufacturer's protocol. The quantity of the extracted RNA was measured using a NanoDrop 2000 (Thermo Fisher Scientific, Waltham, Massachusetts, USA), and the RNA quality was checked by an Agilent tape station Bioanalyser. RNA integrity number (RIN) number for RNA samples typically ranged between 7.5 and 10, with the majority of samples between 9 and 10, as determined at the Genomic Core Facility (GeneCore) in the European Molecular Biology Laboratory (EMBL, Heidelberg, Germany).

### Transcriptomic and gene microarray analyses

At least 100 ng of the extracted total RNA was used as an input for microarray analysis done for each individual. Samples were processed at the EMBL GeneCore Facility for Affymetrix Microarray analysis (Geo number GSE119207). Gene Spring analysis was performed; in addition, Affymetrix CHP data were analysed with Transcriptome Analysis Console 3.1 software using HuGene-2\_0-st-v1 library file from Affymetrix. A cut-off level of 1.5-fold change in the expression was used to analyse the differential gene expression when comparing lesional, non-lesional and healthy conditions. A selection of target genes identified by microarray was then validated via quantitative real-time PCR (qRT-PCR).



**Figure 3** Proposed pathogenically relevant genes overexpressed in lesional CDLE HF. Photographs show HF plucked from the scalp area and used for RNA extraction. A longitudinal HF section is shown in H&E stain highlighting outer root, inner root sheath and hair bulb. The green box shows differentially expressed genes in lesional CDLE HF samples versus non-lesional/healthy control samples. Dermatohistopathology features (right-hand pictures, CDLE biopsies from routine diagnosis, pictures provided by dermatohistopathology, Leeds) such as vacuolar degeneration (apoptosis—*XAF1*), lupus band (middle picture direct immunofluorescence; deposition of immunoglobulin and complement—*C3*),  $CD8^+$  T-cell immune responses ( $\beta 2M$ , *CXCL9*), atrophy (*BMP2*) along with IFN-stimulated genes (*MxA*, *IFI6*, *IFIH1*) correspond to highlighted genes in lesional CDLE HFs.  $\beta 2M$ ,  $\beta 2$  microglobulin; *BMP2*, bone morphogenetic protein 2; CDLE, chronic discoid lupus erythematosus; HF, hair follicle; *IFI6*, IFN inducible protein 6; IFN, interferon; *MxA*, myxovirus protein A; qRT-PCR, quantitative real-time PCR.

### Quantitative real-time PCR

The mRNA expression of the target genes performed for each individual sample in this study was determined with qRT-PCR using a QuantStudio5 Real-Time PCR system (Applied Biosystems, Foster City, California, USA) using Qiagen QuantiTect SYBR Green Master Mix (Qiagen, Manchester, UK). 50–100 ng of total RNA was reverse transcribed using a RevertAid First Strand cDNA Synthesis Kit (Thermo Scientific, Loughborough, UK) according to the manufacturer's instructions. *MxA*, *IFI6*, *IFIH1*, *CXCL9*, *CXCL10*, *BST2*, *OAS2*, *OASL*, *XAF1*, *BMP2*,  $\beta 2M$ , *C3* and *KDR* QuantiTect primer assays were purchased from Qiagen (Hilden, Germany), whereas *U6* snRNA<sup>32</sup> primer (forward—5'-CTCGCTTCGGCAG-CACA-3' and reverse—5'-AACGCTTCACGAATTTGC-3') was purchased from Sigma (Sigma-Aldrich, Poole, UK). The following parameters were used: initial heat activation, 95°C for 5 min; denaturation, 95°C for 10 s; combined annealing and elongation, 60°C for 30 s for a 40 cycle run. Data were analysed using the  $\Delta\Delta CT$  method. mRNA expression of each gene of interest was normalised to *U6* snRNA housekeeping gene.


Statistical analysis of qRT-PCR results were analysed with GraphPad Prism software, V.7.00 (GraphPad Software, La Jolla, San Diego, California, USA). Results for each group are depicted as mean $\pm$ SEM. Data were analysed by using one-way analysis of variance (ANOVA) followed by

the Tukey multiple comparison test to determine statistically significant differences between groups: \* $p < 0.05$ , \*\* $p < 0.01$ .

### RESULTS

#### Microarray analysis of plucked HFs reveals a strong IFN signature and differential expression of complement, apoptosis and MHC I related genes in CDLE lesions


When comparing matched lesional and non-lesional CDLE samples for each patient as well as lesional CDLE with healthy samples, analysis of Affymetrix CHP data with Transcriptome Analysis Console showed substantial upregulation of ISGs, in particular *Mx1*, *IFI6*, *BST2*, *OAS2* and *CXCL10*. This analysis also revealed significant upregulation between CDLE lesional samples compared with non-lesional or healthy samples of  $\beta 2M$ , complement factor 3 (*C3*) and of genes involved in apoptotic cell death (*XAF1*, *OAS2*) which are also known to be IFN inducible (figure 1). Gene Spring analysis of the same microarray dataset indicated that the most robustly differentially expressed genes ( $p < 0.005$ ) in lesional versus non-lesional CDLE included  $\beta 2M$ , *TLR3*, *C1R*, *TAP1*, *SAMHD1*, *IRF1*, *XAF1*, *GBP1*, *OAS2*, *C3*, *IFI27*, *STAT1*, *CXCL10* and *IFIH1/MDA5*. Lesional CDLE samples showed the same set of genes consistently differentially expressed ( $p < 0.005$  for  $\beta 2M$ , *TLR3*, *C1R*, *TAP1*, *SAMHD1*, *XAF1*, *GBP1*, *OAS2*, *C3*, *IFI27*, *STAT1* and *Mx1*), when compared with healthy controls (online supplementary tables 1 and 2).



**Figure 1** Microarray analysis of plucked HF from healthy, lesional and non-lesional CDLE. (A) Hierarchical clustering comparing gene expression in HF between healthy individuals and non-lesional CDLE and (B) between lesional and non-lesional CDLE. (C) Fold change of selected genes from microarray analysis comparing lesional with non-lesional CDLE, and (D) lesional CDLE with healthy. Genes which were also validated by qRT-PCR are indicated in blue. *CXCL9* was not expressed in healthy samples and fold change can therefore not be given. CDLE, chronic discoid lupus erythematosus; HF, hair follicle; qRT-PCR, quantitative real-time PCR.

Thus, with regard to differentially expressed genes associated with lupus pathology, such as ISGs, there is no significant overall difference between non-lesional and healthy samples as illustrated by the HeatMap overview (figure 1A) and figure 1B–D. However, subtle differences existed, such as lack of *CXCL9* expression in healthy but

not in non-lesional CDLE. Interestingly,  $\beta$ 2M was significantly upregulated in non-lesional CDLE ( $p=3.80 \times 10^{-4}$ ) compared with healthy samples in the Gene Spring analysis.



**Figure 2** Differential gene expression in HF from lesional and non-lesional scalp. Selected genes were validated by qRT-PCR and expression compared between lesional and non-lesional psoriasis, CDLE as well as healthy individuals. Genes selected include IFN-stimulated genes (*MxA*, *IFI6*, *CXCL9*, *IFIH1*), apoptosis-related genes (*XAF1*), MHC class I antigen presentation pathway ( $\beta 2M$ ), complement factors (*C3*) and tissue repair-related gene (*BMP2*). Gene expression was normalised to housekeeping gene *U6snRNA*; arbitrary units are given. Except for *BMP2*, the same level of significance was also found between lesional CDLE and both psoriasis samples (not depicted for reason of clarity).  $\beta 2M$ ,  $\beta 2$  microglobulin; *BMP2*, bone morphogenetic protein 2; CDLE, chronic discoid lupus erythematosus; HF, hair follicle; *IFI6*, IFN inducible protein 6; IFN, interferon; MHC, major histocompatibility complex; *MxA*, myxovirus protein A; qRT-PCR, quantitative real-time PCR.

### PCR analysis confirms that a set of genes related to IFN response, complement expression, apoptosis and MHC class I presentation characterises CDLE lesions

To confirm findings from the microarray analysis, qRT-PCR was performed on the same individual samples used in microarray analysis, as well as samples from patients with active scalp psoriasis. Expression of ISGs (*Mx1*, *IFI6*, *CXCL9*, *CXCL10*, *IFIH1/MDA5*), ISGs which are also apoptosis-related (*XAF1*),  $\beta 2M$  and *C3* were significantly higher in lesional CDLE as compared with lesional psoriasis or healthy samples (figure 2). Consistent with microarray data, no significant differences in ISGs expression between non-lesional CDLE and healthy samples were found. Interestingly, bone morphogenetic protein 2 (*BMP2*) was found downregulated in both lesional and non-lesional CDLE samples compared with psoriasis and/or healthy samples. Table 1 provides an overview regarding the fold changes of validated genes comparing CDLE with healthy samples and psoriasis with the same set of healthy samples. We failed to detect a marked IFN signature in psoriasis with the exception of slightly increased *Mx1* and *BST2*. However, Kinase insert domain receptor (*KDR*) which codes for vascular endothelial growth factor receptor 2 (*VEGFR2*) was markedly upregulated in psoriatic samples but not significantly regulated in CDLE samples.

### DISCUSSION

CDLE manifests with chronic inflammatory skin lesions which leads to permanent scarring, resulting in irreversible alopecia when affecting the scalp area.<sup>5 33</sup> Early lesions can be confused with other inflammatory or infectious (eg, fungal) skin reactions, more advanced

ones with scarring alopecias including primary scarring alopecias, such as LPP, alopecia mucinosa, folliculitis decalvans or secondary scarring alopecias such as morphea, cicatricial pemphigoid, neoplastic, traumatic

**Table 1** presents the fold change differences between genes validated by qRT-PCR when comparing average expression in lesional CDLE or psoriasis versus healthy control samples

Fold difference		
Gene symbol	Lesional CDLE versus healthy	Lesional psoriasis versus healthy
<i>Mx1</i>	266*	2.5
<i>IFI6</i>	156*	1.3
<i>CXCL9</i>	498*	1.7
<i>CXCL10</i>	40*	–
<i>BST2</i>	2742	5
<i>IFIH1</i>	8.5*	0.6
<i>OAS2</i>	–	–
<i>OASL</i>	–	–
<i>BMP2</i>	0.09	1.4
<i>XAF1</i>	101*	2.2
$\beta 2M$	13*	1
<i>C3</i>	12.6*	2.4
<i>KDR</i>	2.8	275

\*Indicates genes which were found significantly different using multiple comparison analysis. Fold changes for *OAS2* and *OASL* cannot be given as no expression was detected in healthy controls. CDLE, chronic discoid lupus erythematosus; OAS, 2,5-oligoadenylate synthase; qRT-PCR, quantitative real-time PCR.

or infection-associated granulomatous scarring alopecias. In clinical settings, the diagnosis can be difficult and histopathological confirmation is usually required.<sup>4 33 34</sup> In this study, we performed gene expression analysis on plucked HFs from patients with CDLE and psoriasis, as well as healthy individuals. This pilot study showed that disease-specific signatures can be obtained from plucked HFs and this offers a promising, non-invasive, easy to perform approach with advantages over the use of skin biopsies. However, for diagnostic purposes it would be highly desirable to measure the identified signature molecules on the protein levels as this would allow for a higher sample stability, lower costs and less labour and time-consuming analysis. Our results revealed that ISG are significantly upregulated in lesional CDLE HFs and the pattern of differentially regulated genes resembles those found in previous reports on lesional full skin biopsies from patients with CDLE compared with healthy individuals or patients with psoriasis.<sup>8 35–37</sup> Our finding of increased expression of the apoptosis-related genes, such as *OAS2*, *OASL*, *XAF1*<sup>38–40</sup> in lesional CDLE is also consistent with previously published reports using full skin biopsies.<sup>37 41</sup> Our analysis demonstrates a distinct molecular signature for lesional CDLE with strong upregulation of ISG, apoptosis, complement and MHC I related genes which are hallmarks of the immunopathology in CDLE.<sup>1 8 20 42–44</sup> CXCR3 ligands produced in response to epithelial IFN $\kappa$ /IFN $\lambda$ <sup>15 27 45</sup> attract CD8<sup>+</sup> cytotoxic T cells to the bulge area of the HFs leading to irreversible damage to the stem cell niche residing in this area, thus resulting in atrophic scarring.<sup>23 46–48</sup> Within the normal bulge area, MHC class I and  $\beta$ 2M are found downregulated and this stem cell niche is normally protected from inflammatory challenges, a phenomenon called ‘immune privilege (IP)’<sup>22 23 49</sup> In line with previous findings for scarring LPP,<sup>48</sup> our data also point to IP collapse via increased expression of MHC class I (online supplementary table) and  $\beta$ 2M in HF epithelium of lesional CDLE compared with non-lesional and healthy controls. Our microarray data show a slight tendency for downregulation of the key stem cell markers *K15*, *CD200*, *Lhx2* and *PHLDA1* residing within this area when comparing lesional CDLE with healthy/non-lesional CDLE (data not shown).

Regarding psoriasis, which can present with florid scalp inflammation, the HF IP is maintained and scarring hair loss is not seen although ISG have been found in psoriatic inflammation of the skin. This is in line with previous molecular analysis of scalp psoriasis highlighting it as an interfollicular disease.<sup>30</sup> Supporting those findings, our analysis found low expression of inflammatory molecules including ISGs in HFs from patients with psoriasis. This suggests that the HF may be actively protected from inflammatory attack. In the psoriasis samples, *KDR* expression was higher compared with lesional CDLE (table 1). *KDR* is one of the two VEGFRs. Angiogenesis is due to the action of vascular endothelial growth factor, which is a key molecule in psoriatic skin.<sup>50</sup> Reduced angiogenesis observed in CDLE may be due to high expression

of GBP-1,<sup>20</sup> which is well known for its action on new vessel formation. CDLE is characterised by atrophy and insufficient tissue repair response and the marked downregulation of *BMP2*, which was among the most downregulated genes found, is interesting in this context. BMPs are signalling molecules belonging to the transforming growth factor (TGF $\beta$ ) superfamily.<sup>51</sup> BMPs are implicated in a variety of pathophysiological processes in the skin including wound healing.<sup>51 52</sup> Thus, increased cellular senescence, reduced angiogenesis due to high expression of GBP-1,<sup>20</sup> as well as cytotoxic attack of the HF stem cell compartments and the impact of *BMP2* on TGF $\beta$  pathways are likely to contribute to atrophic scarring seen in patients with CDLE.<sup>53</sup> Once the HF is lost, consequences for tissue repair may deteriorate. Plikus *et al* reported that during the wound healing process, only those dermal cells which were adjacent to the regenerated HFs differentiated into lipid-laden newly formed adipocytes, but those in the hairless skin did not, suggesting that HFs are necessary to establish adipocyte precursors and normal wound repair.<sup>54</sup> Similar to CDLE, LPP leads to permanent scarring<sup>14</sup> and shares some molecular characteristics with CDLE,<sup>48</sup> hence the differential diagnosis can be difficult.<sup>13</sup> While LPP samples were not included in this pilot project, our qRT-PCR validation data showed a significant increase in C3 in lesional CDLE (figure 2), which is used together with immunoglobulin deposits at the basement membrane zone (dermoepidermal junction) to differentiate between patients with CDLE and LPP using DIF in skin biopsies.<sup>11–13</sup> Keratinocytes have been shown to synthesise C3 in response to cytokines, such as CCL2 and IFN $\gamma$ .<sup>55 56</sup> Interestingly, CCL2 was among the most upregulated genes in the microarray analysis (online supplementary data). Our results show only subtle differences between the non-lesional CDLE and healthy samples, suggesting non-lesional HFs of patients with CDLE are only mildly affected by the disease. Patients with CDLE recruited to this study followed measures of ultraviolet (UV) light protection. We have not yet analysed sun-exposed HFs from patients with LE, which could well show subclinical inflammation.

In summary, this pilot project shows great potential regarding the diagnostic value of analysed plucked HFs. Once validated and optimised for protein detection, this could offer a significant advantage in clinical settings where the diagnosis of inflammatory lesions in HF-rich areas, such as the scalp, is often difficult and delayed due to the need of diagnostic biopsies and histopathological assessment. With the results obtained so far, we propose a diagnostic panel (figure 3), using plucked HFs followed by analysis of selected molecules, which should allow distinguishing CDLE from other pathologies. However, this requires validation in a prospective clinical study and comparison with standard dermatohistology diagnosis. Using this approach has the potential to save cost and avoid invasive biopsies, but would also offer significant advantages for research into the pathophysiology of scarring alopecias by allowing repeated sampling in the

context of environmental challenges (eg, UV) or therapeutic interventions.

#### Author affiliations

<sup>1</sup>Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK

<sup>2</sup>National Institute for Health Research (NIHR) Leeds Biomedical Research Centre (BRC), Leeds Teaching Hospitals, Leeds, UK

<sup>3</sup>Genomics Core Facility, European Molecular Biology Laboratory, Heidelberg, Germany

<sup>4</sup>Department of Dermatology, Leeds Teaching Hospitals, Leeds, UK

<sup>5</sup>Department of Histopathology, Leeds Teaching Hospitals NHS Trust, Leeds, UK

<sup>6</sup>Centre for Skin Sciences, Faculty of Life Sciences, University of Bradford, Bradford, UK

<sup>7</sup>Faculty of Biological Sciences, School of Molecular and Cellular Biology, University of Leeds, Leeds, UK

**Contributors** Study design: MW, MJG, EMV; Clinical contribution: AB, MW, MJG, EMV, MYMY; Laboratory-based experiments: MSh, AAA, JP; Data analysis: MSh, MW, MST, EMV; Contribution to critical discussion of results: MST, NVB, MJG, SE, EMV, MW; Manuscript writing and revision: MSh, AAA, SE, MJG, MYMY, AB, NVB, MST, EMV, MW.

**Funding** This project was mainly supported by a grant from Lupus UK with additional funding by Medical Research Council grant MR/M01942X/1. MYMY is funded as an National Institute for Health Research (NIHR) Doctoral Research Fellow and EMV is funded as an NIHR clinician scientist. This research is also supported by the NIHR Leeds Biomedical Research Centre.

**Disclaimer** The views expressed are those of the author(s) and not necessarily those of the National Health Service, the NIHR or the Department of Health.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available in a public, open access repository. All data relevant to the study are included in the article or uploaded as supplementary information.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

- Robinson ES, Werth VP. The role of cytokines in the pathogenesis of cutaneous lupus erythematosus. *Cytokine* 2015;73:326–34.
- Grönhagen CM, Fored CM, Granath F, et al. Cutaneous lupus erythematosus and the association with systemic lupus erythematosus: a population-based cohort of 1088 patients in Sweden. *Br J Dermatol* 2011;164:1335–41.
- Al-Refu K, Goodfield M. Scar classification in cutaneous lupus erythematosus: morphological description. *Br J Dermatol* 2009;161:1052–8.
- Chung HJ, Goldberg LJ. Histologic features of chronic cutaneous lupus erythematosus of the scalp using horizontal sectioning: Emphasis on follicular findings. *J Am Acad Dermatol* 2017;77:349–55.
- Wilson CL, Burge SM, Dean D, et al. Scarring alopecia in discoid lupus erythematosus. *Br J Dermatol* 1992;126:307–14.
- Verma SM, Okawa J, Probert KJ, et al. The impact of skin damage due to cutaneous lupus on quality of life. *Br J Dermatol* 2014;170:315–21.
- Wenzel J, Tütting T. An IFN-associated cytotoxic cellular immune response against viral, self-, or tumor antigens is a common pathogenetic feature in "interface dermatitis". *J Invest Dermatol* 2008;128:2392–402.
- Jabbari A, Suárez-Fariñas M, Fuentes-Duculan J, et al. Dominant Th1 and minimal Th17 skewing in discoid lupus revealed by transcriptomic comparison with psoriasis. *J Invest Dermatol* 2014;134:87–95.
- Wittmann M, Goodfield M. Cytokines in cutaneous lupus erythematosus. *Expert Rev Dermatol* 2011;6:381–94.
- Werth VP. Cutaneous lupus: insights into pathogenesis and disease classification. *Bull NYU Hosp Jt Dis* 2007;65:200–4.
- Burnham TK, Neblett TR, Fine G. The Application of the Fluorescent Antibody Technic to the Investigation of Lupus Erythematosus and Various Dermatoses. *J Invest Dermatol* 1963;41:451–6.
- Isfer RS, Sanches Júnior JA, Festa Neto C, et al. Direct immunofluorescence in lupus erythematosus (LE). *Sao Paulo Med J* 1996;114:1141–7.
- Trachsler S, Trueb RM. Value of direct immunofluorescence for differential diagnosis of cicatricial alopecia. *Dermatology* 2005;211:98–102.
- Bolduc C, Sperling LC, Shapiro J. Primary cicatricial alopecia: lymphocytic primary cicatricial alopecias, including chronic cutaneous lupus erythematosus, lichen planopilaris, frontal fibrosing alopecia, and Graham-Little syndrome. *J Am Acad Dermatol* 2016;75:1081–99.
- Zahn S, Rehkämper C, Kümmerer BM, et al. Evidence for a pathophysiological role of keratinocyte-derived type III interferon (IFN $\lambda$ ) in cutaneous lupus erythematosus. *J Invest Dermatol* 2011;131:133–40.
- Md Yusof MY, Psarras A, El-Sherbiny YM, et al. Prediction of autoimmune connective tissue disease in an at-risk cohort: prognostic value of a novel two-score system for interferon status. *Ann Rheum Dis* 2018;77:annrheumdis-2018-213386.
- Wenzel J, Uerlich M, Wörrenkämper E, et al. Scarring skin lesions of discoid lupus erythematosus are characterized by high numbers of skin-homing cytotoxic lymphocytes associated with strong expression of the type I interferon-induced protein MxA. *Br J Dermatol* 2005;153:1011–5.
- Meller S, Winterberg F, Gilliet M, et al. Ultraviolet radiation-induced injury, chemokines, and leukocyte recruitment: An amplification cycle triggering cutaneous lupus erythematosus. *Arthritis Rheum* 2005;52:1504–16.
- Alase AA, El-Sherbiny YM, Vital EM, et al. IFN $\lambda$  Stimulates MxA production in human dermal fibroblasts via a MAPK-Dependent STAT1-Independent Mechanism. *J Invest Dermatol* 2015;135:2935–43.
- Naschberger E, Wenzel J, Kretz CC, et al. Increased expression of guanylate binding protein-1 in lesional skin of patients with cutaneous lupus erythematosus. *Exp Dermatol* 2011;20:102–6.
- Dang W, Xu L, Yin Y, et al. IRF-1, RIG-I and MDA5 display potent antiviral activities against norovirus coordinately induced by different types of interferons. *Antiviral Res* 2018;155:48–59.
- Paus R, Nickoloff BJ, Ito T. A 'hairy' privilege. *Trends Immunol* 2005;26:32–40.
- Meyer KC, Klatte JE, Dinh HV, et al. Evidence that the bulge region is a site of relative immune privilege in human hair follicles. *Br J Dermatol* 2008;159:1077–85.
- Pi L-Q, Jin X-H, Hwang ST, et al. Effects of calcitonin gene-related peptide on the immune privilege of human hair follicles. *Neuropeptides* 2013;47:51–7.
- Mahajan A, Herrmann M, Muñoz LE. Clearance deficiency and cell death pathways: a model for the pathogenesis of SLE. *Front Immunol* 2016;7:35.
- Gaipi US, Voll RE, Sheriff A, et al. Impaired clearance of dying cells in systemic lupus erythematosus. *Autoimmun Rev* 2005;4:189–94.
- Scholtissek B, Zahn S, Maier J, et al. Immunostimulatory endogenous nucleic acids drive the lesional inflammation in cutaneous lupus erythematosus. *J Invest Dermatol* 2017;137:1484–92.
- Nyberg F, Fransson J, Stephansson E. Proliferation and effects of UVA irradiation in cultured fibroblasts from lesions in cutaneous lupus erythematosus. *Exp Dermatol* 2000;9:53–7.
- Conrad C, Gilliet M. Type I IFNs at the interface between cutaneous immunity and epidermal remodeling. *J Invest Dermatol* 2012;132:1759–62.
- Ruano J, Suárez-Fariñas M, Shemer A, et al. Molecular and cellular profiling of scalp psoriasis reveals differences and similarities compared to skin psoriasis. *PLoS One* 2016;11:e0148450.
- Gho CG, Braun JEF, Tilli CMLJ, et al. Human follicular stem cells: their presence in plucked hair and follicular cell culture. *Br J Dermatol* 2004;150:860–8.
- Galiveti CR, Rozhdestvensky TS, Brosius J, et al. Application of housekeeping npcRNAs for quantitative expression analysis of human transcriptome by real-time PCR. *RNA* 2010;16:450–61.
- Fabbri P, Amato L, Chiarini C, et al. Scarring alopecia in discoid lupus erythematosus: a clinical, histopathologic and immunopathologic study. *Lupus* 2004;13:455–62.

34. Annessi G, Lombardo G, Gobello T, *et al.* A clinicopathologic study of scarring alopecia due to lichen planus: comparison with scarring alopecia in discoid lupus erythematosus and pseudopelade. *Am J Dermatopathol* 1999;21:324–31.
35. Wenzel J, Wörenkämper E, Freutel S, *et al.* Enhanced type I interferon signalling promotes Th1-biased inflammation in cutaneous lupus erythematosus. *J Pathol* 2005;205:435–42.
36. Dey-Rao R, Sinha AA. Genome-wide transcriptional profiling of chronic cutaneous lupus erythematosus (CCLE) peripheral blood identifies systemic alterations relevant to the skin manifestation. *Genomics* 2015;105:90–100.
37. Dey-Rao R, Smith JR, Chow S, *et al.* Differential gene expression analysis in CCLE lesions provides new insights regarding the genetics basis of skin vs. systemic disease. *Genomics* 2014;104:144–55.
38. Choi UY, Kang J-S, Hwang YS, *et al.* Oligoadenylate synthase-like (OASL) proteins: dual functions and associations with diseases. *Exp Mol Med* 2015;47:e144.
39. Lee M-G, Han J, Jeong S-I, *et al.* XAF1 directs apoptotic switch of p53 signaling through activation of HIPK2 and ZNF313. *Proc Natl Acad Sci U S A* 2014;111:15532–7.
40. Liston P, Fong WG, Kelly NL, *et al.* Identification of XAF1 as an antagonist of XIAP anti-Caspase activity. *Nat Cell Biol* 2001;3:128–33.
41. Toberer F, Sykora J, Göttel D, *et al.* Apoptotic signal molecules in skin biopsies of cutaneous lupus erythematosus: analysis using tissue microarray. *Exp Dermatol* 2013;22:656–9.
42. Braunstein I, Klein R, Okawa J, *et al.* The interferon-regulated gene signature is elevated in subacute cutaneous lupus erythematosus and discoid lupus erythematosus and correlates with the cutaneous lupus area and severity index score. *Br J Dermatol* 2012;166:971–5.
43. Wenzel J, Tüting T. Identification of type I interferon-associated inflammation in the pathogenesis of cutaneous lupus erythematosus opens up options for novel therapeutic approaches. *Exp Dermatol* 2007;16:454–63.
44. Zahn S, Graef M, Patsinakidis N, *et al.* Ultraviolet light protection by a sunscreen prevents interferon-driven skin inflammation in cutaneous lupus erythematosus. *Exp Dermatol* 2014;23:516–8.
45. Sarkar MK, Hile GA, Tsoi LC, *et al.* Photosensitivity and type I IFN responses in cutaneous lupus are driven by epidermal-derived interferon kappa. *Ann Rheum Dis* 2018;77:1653–64.
46. Ito M, Cotsarelis G. Is the hair follicle necessary for normal wound healing? *J Invest Dermatol* 2008;128:1059–61.
47. Al-Refu K, Edward S, Ingham E, *et al.* Expression of hair follicle stem cells detected by cytokeratin 15 stain: implications for pathogenesis of the scarring process in cutaneous lupus erythematosus. *Br J Dermatol* 2009;160:1188–96.
48. Harries MJ, Meyer K, Chaudhry I, *et al.* Lichen planopilaris is characterized by immune privilege collapse of the hair follicle's epithelial stem cell niche. *J Pathol* 2013;231:236–47.
49. Ito T, Ito N, Bettermann A, *et al.* Collapse and restoration of MHC class-I-dependent immune privilege: exploiting the human hair follicle as a model. *Am J Pathol* 2004;164:623–34.
50. Detmar M, Brown LF, Claffey KP, *et al.* Overexpression of vascular permeability factor/vascular endothelial growth factor and its receptors in psoriasis. *J Exp Med* 1994;180:1141–6.
51. Botchkarev VA. Bone morphogenetic proteins and their antagonists in skin and hair follicle biology. *J Invest Dermatol* 2003;120:36–47.
52. Moura J, da Silva L, Cruz MT, *et al.* Molecular and cellular mechanisms of bone morphogenetic proteins and activins in the skin: potential benefits for wound healing. *Arch Dermatol Res* 2013;305:557–69.
53. Korntner S, Lehner C, Gehwolf R, *et al.* Limiting angiogenesis to modulate scar formation. *Adv Drug Deliv Rev* 2018. doi:10.1016/j.addr.2018.02.010. [Epub ahead of print: 03 Mar 2018].
54. Plikus MV, Guerrero-Juarez CF, Ito M, *et al.* Regeneration of fat cells from myofibroblasts during wound healing. *Science* 2017;355:748–52.
55. Terui T, Ishii K, Ozawa M, *et al.* C3 production of cultured human epidermal keratinocytes is enhanced by IFN $\gamma$  and TNF $\alpha$  through different pathways. *J Invest Dermatol* 1997;108:62–7.
56. Purwar R, Wittmann M, Zwirner J, *et al.* Induction of C3 and CCL2 by C3a in keratinocytes: a novel autocrine amplification loop of inflammatory skin reactions. *J Immunol* 2006;177:4444–50.