Abstract 255 Table 1 Histological characteristics of patients

RPS/ISN Classification	Number of patients
Class II	1
Class III	4
Class IV	11
Class V	6
Class III+IV	6
Class IV+V	2

treated with high dose steroids. The other medications, the patients were previously treated with, were Cyclophosphamide (n=25), Azathioprine (n=7), Mycophenolate mofetil (n=17). Also one patient was treated with plasmapheresis due to thrombotic thrombocytopenic purpura. Two patients were lost in follow-up. Twelve patients (38%) are still on rituximab, in which 5 of them (42%) still have active LN. Treatment of 11 out of 18 patients was terminated owing to remission of disease. All of the patients whom rituximab was terminated in disease remission did not flare, except one. Moreover, in 7 patients, treatment was terminated because 3 had inadequate response, 2 had severe infection (cellulitis, pneumonia), 1 had Rituximab-induced serum sickness, 1 had will of pregnancy. The mean level of proteinuria decreased from 3440 ± 2476 mg/day to 927 ± 895 mg/day ($p < 0.0001$). Additionally, there was a reduce in steroid dose from $23,75 \pm 19,6$ mg/day to $9,02 \pm 10,59$ mg/day ($p < 0.0001$). None of the patients had a new organ involvement during therapy.

Conclusions In our single center, most of the patients were class IV nephritis whom we observed the efficacy of rituximab. Generally, the side effects were acceptable. Rituximab can be a steroid sparing agent in these patients.

Funding Source(s): None

256

EXPLORING THE PERCEIVED IMPACT OF THE CHRONIC DISEASE SELF-MANAGEMENT PROGRAM ON SELF-MANAGEMENT BEHAVIORS AMONG AFRICAN AMERICAN WOMEN WITH LUPUS: A QUALITATIVE STUDY

¹Abena Twumasi, ¹Anna Shao, ²Charmayne Dunlop-Thomas, ¹Hannah Cooper, ²Cristina Drenkard. ¹Emory Rollins School of Public Health; ²Emory University

10.1136/lupus-2019-lsm.256

Background Effective self-management is critical to improve outcomes in individuals with systemic lupus erythematosus (SLE). African Americans with SLE face formidable barriers to accessing self-management education. We qualitatively explored the perceived processes through which the Chronic Disease Self-Management Program (CDSMP), an evidence-based, worldwide disseminated educational intervention for people with chronic conditions, affects self-management behaviors among African American women with SLE.

Methods Participants were recruited from the WELL (Women Empowered to Live with Lupus) Study, a behavioral trial of the effectiveness of the CDSMP on African American women with SLE. The CDSMP is a community-based, small-group program designed to enhance self-efficacy and self-management skills in people with chronic conditions. The CDSMP

addresses self-management challenges that are universal across chronic conditions (e.g. pain and fatigue management, healthy eating, physical activity, communication with providers). Using a longitudinal pre-/post-intervention design, we conducted one-on-one, semi-structured interviews among a purposive sample of WELL participants. Pre-intervention interviews explored self-management behaviors at baseline; post-intervention interviews focused on changes in these behaviors after the CDSMP. Transcripts were analyzed using thematic methods.

Results Twenty-four and 23 women participated in the pre- and post-intervention interviews, respectively. Mean age and disease duration were 48.6 (SD=13.5) and 14.1 years (SD=8.1), respectively. Most women (91.7%) were employed and 1/3 reported an annual income of \$10,000-\$19 000. The sample was balanced with regard to education, depression, and SLE activity.

Participants perceived the CDSMP to be a valuable resource that helped them to improve fundamental self-management behaviors, including exercise, relaxation, diet, and medication-taking. With few exceptions, womens reported changes in self-management behaviors were not dependent on age, education, SLE activity or depression status. Womens most widespread perceived effects of the CDSMP were on relaxation and exercise. Strategies that generated improvements in those behaviors included goal setting, action planning, encouragement to pursue low-impact physical activity, and introduction of mindfulness techniques to better manage SLE symptoms.

Conclusions African American women living with SLE, a population vulnerable to high rates of SLE morbidity and mortality, perceived benefits from CDSMP participation. The programs potential to catalyze improvements in self-management behaviors was independent of womens demographic or disease characteristics. Because the CDSMP is widely disseminated

Funding Source(s): Research reported in this publication was supported by the National Institute On Minority Health And Health Disparities of the National Institutes of Health under Award Number R01MD010455. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

257

RESIDENTIAL EXPOSURES ARE ASSOCIATED WITH INCREASED ODDS OF SLE DIAGNOSIS

¹Jing Li*, ²Laura Trupin, ³Milena A Gianfrancesco, ³Cristina Lanata, ²Patricia Katz, ²Maria C Dall'era, ²Jinoos Yazdany. ¹University of California, San Francisco; ²UC San Francisco; ³Division of Rheumatology, Department of Medicine, University of California, San Francisco

10.1136/lupus-2019-lsm.257

Background Environmental exposures, such as pesticides, silica, and asbestos have been associated with risk of SLE. In this study, we assessed the association between different residential exposures and SLE diagnosis in a diverse cohort of lupus participants and healthy controls.

Methods Data were from the California Lupus Epidemiology Study (CLUES), a racially/ethnically diverse cohort of SLE patients, and healthy controls with no history of autoimmune systematic diseases who resided in the San Francisco Bay Area. SLE diagnoses were confirmed through medical records and rheumatologist clinical examination. All participants completed a structured interview that included demographics (age, sex, race/ethnicity and education) and residential exposures (frequent exterminator use, other home insecticide use, kerosene