adjusting for age, sex, disease duration, total cumulative corticosteroid dose, and top 10 PCs as covariates. In further analysis of SLE patients with AVN compared with SLE patients without AVN treated by a high corticosteroid [cumulative dose >20 g], the AVN-risk loci were highly expressed in fibroblasts ($p=8.78 \times 10^5$) and musculoskeletal system ($p=1.13 \times 10^4$) using DEPICT.

Conclusions These findings provide genetic evidence that pathway of fibroblasts are relevant to osteonecrosis. Individual genetic risk for the development of AVN should be assessed prior to treatment for SLE.

Funding Source(s): None

136 INFLUENCE OF GENETIC RISK VARIANTS ON THE CLINICAL SUBPHENOTYPES OF SYSTEMIC LUPUS ERYTHEMATOSUS IN A KOREAN COHORT

¹Hyuk-Hee Kwon, ²Eunji Ha, ¹So-Young Bang*, ¹Hye-Soon Lee, ²Kwangwoo Kim, ¹Sang-Cheol Bae. ¹Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases; ²Department of Biology, Kyung Hee University, Republic of Korea

10.1136/lupus-2019-lsm.136

Background Systemic lupus erythematosus (SLE) is a multifactorial autoimmune disease with diverse heterogeneous phenotypes. Impact of complex genetic variants on the clinical manifestations of SLE was not fully understood. The aim of this study is to identify the cumulative effect of SLE-risk variants on clinical manifestations and autoantibody profiles in patients with SLE.

Methods A total of 781 Korean patients from the Hanyang BAE lupus cohort were genotyped by Immunochip and genome-wide association study (GWAS) using Illumina HumanOmni1-Quad array. Weighted genetic risk score (GRS) were calculated from 45 well-validated non-HLA SNPs and HLA SLE-risk loci. Individual GRS was tested for associations with the clinical subphenotypes based on 1997 ACR criteria of SLE and the development of autoantibody by using a linear regression or a logistic regression adjusting for the top 10 PCs as covariates, respectively.

Results We identified weighted GRS calculated from non-HLA and HLA SLE-risk loci with significant associations in various SLE subphenotypes defined by the ACR criteria (mean number 5.74) among 11 criteria. Individuals weighted GRS showed significantly positive correlation with the number of ACR criteria in a linear regression model (coefficient=0.128, $p=9.00 \times 10^3$). Consistently, a significant positive correlation with the number of ACR in both non-HLA GRS (coefficient=0.110, $p=0.027$) and HLA GRS (coefficient=0.06, $p=0.021$) was observed, respectively.

In a clinical subphenotype analysis, weighted GRS from non-HLA and HLA risk loci were significantly related to malar rash (OR 1.23, $p=2.68 \times 10^3$), renal disorder (OR 1.15, $p=4.41 \times 10^2$), and thrombocytopenia (OR 1.21, $p=7.55 \times 10^3$) using a multivariable logistic regression. Weighted GRS were strongly associated with production of anti-DNA antibody (OR 1.38, $p=2.18 \times 10^3$).

Conclusions In conclusion, a high cumulative SLE-risk effect promoted diverse clinical manifestations and influenced the production of auto-antibodies including anti-DNA, that could provide clues to identifying distinctive mechanisms in SLE.

Funding Source(s): None

137 MYOCARDITIS IN PEDIATRIC LUPUS: A CLINICAL CONUNDRUM

¹Nameirakpam Johnson*, ²Deepti Suri, ³Anju Gupta, ³Amit Rawat, ²Surjit Singh. ¹Dept. of Pediatrics, Allergy- Immunology Unit, Postgraduate Institute of Medical Education and Research; ²Postgraduate Institute of Medical Education and Research, Chandigarh, India; ³PGIMER Chandigarh

10.1136/lupus-2019-lsm.137

Background Myocarditis in pediatric lupus is uncommon but carry a significant morbidity and mortality.

Methods We reviewed our cohort of children with lupus registered between January 1993 to November 2018. Of the 140 lupus patients that were diagnosed during this time, 4 had myocarditis.

Abstract 137 Table 1 Laboratory Investigations

Investigation	Case 1	Case 2	Case 3	Case 4
Haemoglobin (g/L)	94 g/L	92 g/L	80 g/L	69 g/L
White cell counts	$6.7 \times 10^9/L$	$7.2 \times 10^9/L$	$6.5 \times 10^9/L$	$6.4 \times 10^9/L$
Lymphocyte count	$1.15 \times 10^9/L$	$1.91 \times 10^9/L$	$1.65 \times 10^9/L$	$1.25 \times 10^9/L$
Platelets	$130 \times 10^9/L$	$150 \times 10^9/L$	$220 \times 10^9/L$	$90 \times 10^9/L$
Urine routine	No RBC,	No RBC,	No RBC,	Plenty RBCs, 3+albumin
Urine protein (mg/m ² /hour)	No albumin	No albumin	No albumin	82 mg/m ² /hour
C3 (Normal 50–150 mg/dL)	27 mg/dl	28 mg/dL	31 mg/dl	21 mg/dl
C4 (Normal 20–50 mg/dL)	4 mg/dl	3 mg/dL	10 mg/dl	3 mg/dl
ANA	4+diffuse	3+diffuse	3+diffuse	4+diffuse
Anti dsDNA (N:<25 IU/mL)	59	583	112	346
Antiphospholipid antibodies:	Positive	Negative	Negative	Negative
a) Lupus anticoagulant	Negative	Negative	Negative	Negative
b) Anticardiolipin antibody (IgG and IgM) c) Anti B2	Negative	Negative	Negative	Negative
Glycoprotein –1 antibody (IgG and IgM)				
2Dimensional Echocardiography	Mild pericardial effusion, dilated left ventricle, ejection fraction (EF) 30%	Mild pericardial effusion, EF 25%	Mild pericardial effusion, moderate mitral regurgitation, mild tricuspid regurgitation, global hypokinesia, EF 56%	Moderate mitral regurgitation, mild tricuspid regurgitation, global hypokinesia, EF 18%