Towards a practical management of fatigue in systemic lupus erythematosus

Philippe Mertz ^{1,2} Aurélien Schlencker,^{1,2} Matthias Schneider,³ Pierre-Edouard Gavand,^{2,4} Thierry Martin,^{2,4} Laurent Arnaud ^{1,2}

To cite: Mertz P, Schlencker A, Schneider M, *et al.* Towards a practical management of fatigue in systemic lupus erythematosus. *Lupus Science & Medicine* 2020;7:e000441. doi:10.1136/lupus-2020-000441

Received 25 August 2020
 Revised 22 October 2020
 Accepted 30 October 2020

ABSTRACT

Significant fatigue is reported by two-thirds of patients with SLE and severe fatigue by one-third. The assessment and treatment of fatigue remains a major challenge in SLE, especially in patients with no disease activity. Here, we suggest a practical algorithm for the management of fatigue in SLE. First, common but non-SLE-related causes of fatigue should be ruled out based on medical history, clinical and laboratory examinations. Then, presence of SLE-related disease activity or organ damage should be assessed. In patients with active disease, remission is the most appropriate therapeutic target while symptomatic support is needed in case of damage. Both anxiety and depression are major independent predictors of fatigue in SLE and require dedicated assessment and care with psychological counselling and pharmacological intervention if needed. This practical algorithm will help in improving the management of one the most common and complex patient complaints in SLE.

SLE is a chronic autoimmune disease which may cause a broad spectrum of clinical manifestations as well as subjective symptoms^{1 2} such as fatigue. In SLE, fatigue is reported by 67% to 90% of patients^{1 3 4} and is rated as severe in up to one-third of patients using validated fatigue instruments, as shown in the recent multicentre FATILUP study.¹ Also, fatigue is often reported as the most debilitating symptom of the disease by patients^{5 6} and leads to both altered health-related quality of life^{4 7–9} and significant work disability with tremendous indirect costs.³ The rational assessment and treatment of fatigue remains a major challenge in SLE,⁵ especially in patients without active disease. Noteworthy, fatigue is a highly multifactorial concept¹⁰ which, in the context of SLE, may be due either to lupus-related or non-lupus-related general causes. Importantly, those causes can also be intricate with significant psychobehavioural determinants. Here, we suggest a practical step-by-step algorithm for the general assessment and management of fatigue in SLE (figure 1).

STEP 1: SEARCH FOR A GENERAL (NON-LUPUS-RELATED) CAUSE OF FATIGUE

In the primary care setting, a medical or psychiatric diagnosis can be found in at least two-thirds of patients presenting with acute fatigue. The most frequent causes of fatigue are summarised in table 1. It is crucial to understand the patient's perspective, the detailed history of fatigue and the impact on the patient's mood as well as on daily activities such as work, household chores, physical activity and leisure. Box 1 summarises the main questions to be asked when confronted with a patient with fatigue. A review of current medications is suggested, as some drugs may induce fatigue (eg, antihypertensive drugs such as beta blockers or sedatives). Lifestyle assessment is also essential: a significant association between smoking and fatigue has been reported in SLE.^{11–14} Also, obesity has been associated with increased fatigue both in the general population and SLE.¹⁵

A full clinical examination of the heart, lung, thyroid and nervous system¹⁶ is crucial. For women, a gynaecological examination is also recommended, especially in case of anaemia.^{17 18} Laboratory tests should rule out most common causes of fatigue (box 1) such as inflammation and infection, anaemia, renal or hepatic failure, viral hepatitis or HIV infection, and major endocrine or metabolic complications such as abnormal calcemia, hypothyroidism, diabetes or adrenal insufficiency (especially in patients who recently stopped glucocorticoids). Cancer screening should be updated, according to current guidelines, and indirect signs of malignancy (anorexia, weight loss and lymphadenopathy) should be carefully searched for. Although controversial,¹⁹ vitamin D deficiency has been associated with fatigue in SLE.²⁰ In an observational study of 80 patients with SLE, vitamin D supplementation improved fatigue in participants.²¹ Also, vitamin D supplementation was associated with a decrease

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Rheumatology, Hôpitaux universitaires de Strasbourg, Strasbourg, France

²Centre National de Référence des Maladies Systémiques et Auto-immunes Rares Est Sud-Ouest (RESO), Strasbourg, France, France

³Policlinic for Rheumatology & Hiller Research Centre for Rheumatology, Heinrich-Heine-University, Düsseldorf, Germany

⁴Service d'immunologie clinique, Hôpitaux universitaires de Strasbourg, Strasbourg, France

Correspondence to

Pr Laurent Arnaud; laurent.arnaud@chru-strasbourg.fr

Figure 1 Practical algorithm for the management of fatigue in patients with SLE.

in fatigue scores in a randomised controlled trial in juvenile-onset SLE.²²

STEP 2: OBJECTIVE ASSESSMENT OF FATIGUE USING VALIDATED PROS

Because fatigue is a highly subjective symptom, the standardised assessment of fatigue using validated patient-reported outcomes (PROs) is an important step. The use of validated PROs also allows for an individual follow-up of fatigue intensity and symptoms over time, and may help in underlining the benefit of a therapeutic intervention at the patient level. It is also a way to show that the physician is genuinely interested in understanding and treating the cause of fatigue, which is important from the patient's perspective, and helps in establishing a trusting physician–patient relationship. Among a total of 16 different fatigue PROs which have been used in SLE, the FSS (Fatigue Severity Scale) and the short-FSS are the most used ones, but the MFI (Multi-dimensional Fatigue Inventory) and the Fatigue-VAS have also been used, although less commonly¹⁶ (see table 2). The FACIT-Fatigue score is

Box 1 Checklist for the initial assessment of fatigue in SLE

- 1. Characterisation of fatigue:**
 - Onset: acute or insidious
 - Evolution: recent (<1 month), persistent (<6 months) or chronic (>6 months)
 - Presence of fatigue-free periods
 - Is fatigue ameliorated by rest?
 - Physical and mental impact of fatigue
- 2. Assessment of lifestyle** (work, restrictive diet or obesity, physical activity), **habitus** (smoking, drinking) and **sleep quality**
- 3. Presence of concurrent manifestations**
 - Lupus flare?
 - Any other (non-SLE-related) disorder?
- 4. List of medications**
- 5. List of previous or current associated medical conditions**

commonly used in clinical trials but has been infrequently used in routine clinical practice.¹

STEP 3: IDENTIFYING A LUPUS-RELATED CAUSE OF FATIGUE

Fatigue can be a manifestation of active SLE but can also be related to organ damage (figure 2). The relationship between disease activity and fatigue remains controversial in SLE and has been shown to be less strong than with other factors such as anxiety or depression.^{1 23 24} Its association with serological markers (C3, anti-dsDNA antibodies) is also controversial.¹¹

Fatigue has been associated with several SLE-specific organ manifestations:

- Several studies found an association between neurological involvement,^{11 25} including white matter hyperintensities,²⁶ and fatigue.

Table 1 Frequent causes of fatigue and suggested first-line assessment

Common causes of fatigue		First-line assessment
Drug-induced fatigue	Steroids, anti-arrhythmic, anti-hypertensive, benzodiazepines or other antidepressant and sedative agents, anti-histaminic, diuretics	Medication review
Pregnancy		β-HCG
Anaemia	Iron or vitamin deficiencies	C-reactive protein, haemoglobin, Coombs test, ferritin, B ₉ and B ₁₂ vitamins
Metabolic disorders	Hypothyroidism, adrenal insufficiency, panhypopituitarism	TSH, fasting glucose...
Vitamin D insufficiency		Vitamin D levels
Infection	Chronic bacterial (mycobacterial) or viral (HIV, hepatitis B and C, EBV) infections	C reactive protein Hepatitis B/C, EBV and HIV serology Quantiferon
Organ failure	Cardiac insufficiency Respiratory insufficiency Liver insufficiency Kidney insufficiency	BNP, chest X-ray, echocardiography creatinine, full hepatic tests
Sleep disorders	Obstructive sleep apnoea syndrome Jet lag syndrome in frequent travellers	Polysomnography

b-HCG, b-human chorionic gonadotropin hormone; BNP, brain natriuretic peptide; EBV, Epstein-Barr virus; TSH, thyroid-stimulating hormone.

Table 2 Most frequently used fatigue patient-reported outcomes (PROs) in SLE

Fatigue PROs	Description
Fatigue Severity Scale (FSS)	9-item scale covering the general aspects of fatigue Originally derived for people with multiple sclerosis and SLE
Multi-dimensional Fatigue Inventory (MFI)	20-item scale divided into five domains: general fatigue, physical fatigue, mental fatigue, reduced motivation and reduced activity The threshold for significant fatigue depends on age and gender
Visual analogue scale to evaluate fatigue severity (VAS-F)	The scale consists of 18 items related to the subjective experience of fatigue, using fatigue and energy subscales
Functional Assessment of Chronic Illness Therapy–Fatigue (FACIT-Fatigue)	13-item self-reported questionnaire assessing aspects of physical and mental fatigue, and their effects on function and daily living

- Renal failure can be an important cause of fatigue. It is crucial to assess whether it is related to active kidney disease or to chronic lesions (damage).^{27 28}
- Cardiac failure is an obvious cause of fatigue.
- Hepatic failure and cirrhosis can (rarely) be due to lupus hepatitis, and overlap with autoimmune hepatitis or less frequently sclerosing cholangitis.^{29 30}

Other clinical manifestations of SLE have been associated with fatigue: in the FATILUP study,¹ we found arthritis and oral ulcers to be individual SLE Disease Activity Index score components associated with severe fatigue. This may underline a more specific role for painful disease manifestations in SLE.^{11 23}

Finally, the prevalence of fibromyalgia is estimated to range between 6.2% and 22% of patients with SLE^{31 32} and has been strongly associated with fatigue.³³ Sjögren's syndrome should also be thought of in patients with sicca syndrome and has been associated with significant fatigue across several studies.^{34 35}

STEP 4: LOOKING FOR INTRICATE PSYCHOLOGICAL DETERMINANTS

Emotional and functional well-beings as well as abnormal illness-related behaviours strongly correlate with depression and fatigue in SLE.^{1 36 37} Pain, stress and depression have been shown to be the most important predictors of fatigue in patients with SLE³⁸ and their intensity and

consequences should be assessed, with the help of a psychologist or psychiatrist if needed. Mood disorders are reported in up to 13% of patients with SLE and attributed to SLE in about 40% of cases.³⁹ Sleep disorders are also common in the general population as well as in SLE, and have been associated with fatigue^{40–42} and depression.⁴¹

TOWARDS A PRACTICAL MANAGEMENT OF FATIGUE IN PATIENTS WITH SLE

In patients with significant disease activity, the main therapeutic target is remission (or alternatively low-disease activity) and reaching these goals can be sufficient to improve fatigue.^{43 44} However, a common situation is the presence of significant fatigue contrasting with the absence of disease activity or any underlying organic cause. In these patients, immunosuppressive treatment escalation is not indicated and other non-pharmacological interventions such as psychological and behavioural assessment or physical activity workshops should be favoured.⁴⁵ Importantly, lack of optimal physical activity as well as sedentary behaviour have been associated with fatigue in SLE.^{44 46 47}

Physical exercise is recommended for the management of pain and fatigue in patients with inflammatory arthritis in the last European League Against Rheumatism recommendations.⁴⁸ Physical activity has been shown to improve fatigue in patients with SLE,^{44 49} with more time spent in moderate or high physical activity associated with less fatigue.⁵⁰ Patients with SLE with otherwise unexplained fatigue should undergo dedicated psychological assessment, and behavioural issues should be specifically taken care of using appropriate psychological counselling^{51–55} and pharmacological intervention, when needed. The exact benefit of antidepressants on fatigue is difficult to assess in SLE because there is no specific trial, but there is no reason to believe that those treatments would not be appropriate, keeping in mind the potential interaction with hydroxychloroquine, which may lead to QT prolongation. Tobacco smoking cessation should be encouraged as it significantly reduces therapeutic efficacy of many drugs and could promote flares.¹⁴ Last but not least, hydroxychloroquine observance should be evaluated and if needed, non-scheduled hydroxychloroquine serum

Figure 2 Main determinants of fatigue in patients with SLE.

concentrations should be measured to verify therapeutic adherence and adjust daily posology.^{56 57}

CONCLUSION

Significant fatigue is reported by two-thirds of patients with SLE and severe fatigue by one-third.¹ As in the general population, general non-SLE-related causes of fatigue should be ruled out. Then, it is important to assess whether fatigue may be related to disease activity or damage. In the former situation, disease remission is the most appropriate therapeutic target, with an emphasis on painful manifestations. Importantly, both anxiety and depression are major independent predictors of fatigue in several studies. These manifestations should be thoroughly assessed and taken care of using appropriate psychological counselling and pharmacological intervention, when needed. We believe this practical algorithm will help in improving the management of one of the most common and complex patient complaints in SLE.

Twitter Laurent Arnaud @Lupusreference

Contributors All authors have equally contributed to this work.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No additional data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Philippe Mertz <http://orcid.org/0000-0002-9781-7388>

Laurent Arnaud <http://orcid.org/0000-0002-8077-8394>

REFERENCES

- Arnaud L, Gavand PE, Voll R, *et al*. Predictors of fatigue and severe fatigue in a large international cohort of patients with systemic lupus erythematosus and a systematic review of the literature. *Rheumatology* 2019;58:987–96.
- Cleanthous S, Tyagi M, Isenberg DA, *et al*. What do we know about self-reported fatigue in systemic lupus erythematosus? *Lupus* 2012;21:465–76.
- Baker K, Pope J. Employment and work disability in systemic lupus erythematosus: a systematic review. *Rheumatology* 2009;48:281–4.
- Fonseca R, Bernardes M, Terroso G, *et al*. Silent burdens in disease: fatigue and depression in SLE. *Autoimmune Dis* 2014;2014:1–9.
- Felten R, Sagez F, Gavand P-E, *et al*. 10 most important contemporary challenges in the management of SLE. *Lupus Sci Med* 2019;6:e000303.
- Schneider M. Pitfalls in lupus. *Autoimmun Rev* 2016;15:1089–93.
- Devilliers H, Amoura Z, Besancenot J-F, *et al*. Responsiveness of the 36-item short form health survey and the lupus quality of life questionnaire in SLE. *Rheumatology* 2015;54:940–9.
- Jolly M, Annareddy N, Arnaud L, *et al*. Changes in quality of life in relation to disease activity in systemic lupus erythematosus: post-hoc analysis of the BLISS-52 trial. *Lupus* 2019;28:1628–39.
- Sutanto B, Singh-Grewal D, McNeil HP, *et al*. Experiences and perspectives of adults living with systemic lupus erythematosus: thematic synthesis of qualitative studies. *Arthritis Care Res* 2013;65:1752–65.
- Nikolaus S, Bode C, Taal E, *et al*. Which dimensions of fatigue should be measured in patients with rheumatoid arthritis? A Delphi study. *Musculoskeletal Care* 2012;10:13–17.
- Burgos PI, Alarcón GS, McGwin G, *et al*. Disease activity and damage are not associated with increased levels of fatigue in systemic lupus erythematosus patients from a multiethnic cohort: LXVII. *Arthritis Rheum* 2009;61:1179–86.
- Yilmaz-Oner S, Ilhan B, Can M, *et al*. Fatigue in systemic lupus erythematosus: association with disease activity, quality of life and psychosocial factors. *Z Rheumatol* 2017;76:913–9.
- Pettersson S, Boström C, Eriksson K, *et al*. Lifestyle habits and fatigue among people with systemic lupus erythematosus and matched population controls. *Lupus* 2015;24:955–65.
- Parisis D, Bernier C, Chasset F, *et al*. Impact of tobacco smoking upon disease risk, activity and therapeutic response in systemic lupus erythematosus: a systematic review and meta-analysis. *Autoimmun Rev* 2019;18:102393.
- Patterson SL, Schmajuk G, Jafri K, *et al*. Obesity is independently associated with worse patient-reported outcomes in women with systemic lupus erythematosus. *Arthritis Care Res* 2019;71:126–33.
- Pettersson S, Lundberg IE, Liang MH, *et al*. Determination of the minimal clinically important difference for seven measures of fatigue in Swedish patients with systemic lupus erythematosus. *Scand J Rheumatol* 2015;44:206–10.
- Zard E, Arnaud L, Mathian A, *et al*. Increased risk of high grade cervical squamous intraepithelial lesions in systemic lupus erythematosus: a meta-analysis of the literature. *Autoimmun Rev* 2014;13:730–5.
- Wincup C, Parnell C, Cleanthous S, *et al*. Red cell distribution width correlates with fatigue levels in a diverse group of patients with systemic lupus erythematosus irrespective of anaemia status. *Clin Exp Rheumatol* 2019;37:852–4.
- Ruiz-Irastorza G, Egurbide MV, Olivares N, *et al*. Vitamin D deficiency in systemic lupus erythematosus: prevalence, predictors and clinical consequences. *Rheumatology* 2008;47:920–3.
- Stockton KA, Kandiah DA, Paratz JD, *et al*. Fatigue, muscle strength and vitamin D status in women with systemic lupus erythematosus compared with healthy controls. *Lupus* 2012;21:271–8.
- Ruiz-Irastorza G, Gordo S, Olivares N, *et al*. Changes in vitamin D levels in patients with systemic lupus erythematosus: effects on fatigue, disease activity, and damage. *Arthritis Care Res* 2010;62:1160–5.
- Lima GL, Paupitz J, Aikawa NE, *et al*. Vitamin D supplementation in adolescents and young adults with juvenile systemic lupus erythematosus for improvement in disease activity and fatigue scores: a randomized, double-blind, placebo-controlled trial. *Arthritis Care Res* 2016;68:91–8.
- Moldovan I, Cooray D, Carr F, *et al*. Pain and depression predict self-reported fatigue/energy in lupus. *Lupus* 2013;22:684–9.
- Da Costa D, Dritsa M, Bernatsky S, *et al*. Dimensions of fatigue in systemic lupus erythematosus: relationship to disease status and behavioral and psychosocial factors. *J Rheumatol* 2006;33:1282–8.
- Zonana-Nacach A, Roseman JM, McGwin G, *et al*. Systemic lupus erythematosus in three ethnic groups. VI: Factors associated with fatigue within 5 years of criteria diagnosis. *Lupus* 2000;9:101–9.
- Harboe E, Greve OJ, Beyer M, *et al*. Fatigue is associated with cerebral white matter hyperintensities in patients with systemic lupus erythematosus. *J Neurol Neurosurg Psychiatry* 2008;79:199–201.
- Gregg LP, Jain N, Carmody T, *et al*. Fatigue in nondialysis chronic kidney disease: correlates and association with kidney outcomes. *Am J Nephrol* 2019;50:37–47.
- Artom M, Moss-Morris R, Caskey F, *et al*. Fatigue in advanced kidney disease. *Kidney Int* 2014;86:497–505.
- Adiga A, Nugent K. Lupus hepatitis and autoimmune hepatitis (lupoid hepatitis). *Am J Med Sci* 2017;353:329–35.
- Kadokawa Y, Omagari K, Matsuo I, *et al*. Primary sclerosing cholangitis associated with lupus nephritis: a rare association. *Dig Dis Sci* 2003;48:911–4.
- Torrente-Segarra V, Salman-Monte TC, Rúa-Figueroa Íñigo, *et al*. Fibromyalgia prevalence and related factors in a large registry of patients with systemic lupus erythematosus. *Clin Exp Rheumatol* 2016;34:S40–7.
- Wolfe F, Petri M, Alarcón GS, *et al*. Fibromyalgia, systemic lupus erythematosus (SLE), and evaluation of SLE activity. *J Rheumatol* 2009;36:82–8.
- Macfarlane GJ, Kronisch C, Dean LE, *et al*. EULAR revised recommendations for the management of fibromyalgia. *Ann Rheum Dis* 2017;76:318–28.

- 34 Gilboe IM, Kvien TK, Uhlig T, *et al.* Sicca symptoms and secondary Sjögren's syndrome in systemic lupus erythematosus: comparison with rheumatoid arthritis and correlation with disease variables. *Ann Rheum Dis* 2001;60:1103–9.
- 35 Carsons SE, Vivino FB, Parke A, *et al.* Treatment guidelines for rheumatologic manifestations of Sjögren's syndrome: use of biologic agents, management of fatigue, and inflammatory musculoskeletal pain. *Arthritis Care Res* 2017;69:517–27.
- 36 Branco JC, Rodrigues AM, Gouveia N, *et al.* Prevalence of rheumatic and musculoskeletal diseases and their impact on health-related quality of life, physical function and mental health in Portugal: results from EpiReumaPt – a national health survey. *RMD Open* 2016;2:e000166.
- 37 Choi ST, Kang JI, Park I-H, *et al.* Subscale analysis of quality of life in patients with systemic lupus erythematosus: association with depression, fatigue, disease activity and damage. *Clin Exp Rheumatol* 2012;30:665–72.
- 38 Azizoddin DR, Gandhi N, Weinberg S, *et al.* Fatigue in systemic lupus: the role of disease activity and its correlates. *Lupus* 2019;28:163–73.
- 39 Hanly JG, Su L, Urowitz MB, *et al.* Mood disorders in systemic lupus erythematosus: results from an international inception cohort study. *Arthritis Rheumatol* 2015;67:1837–47.
- 40 Da Costa D, Bernatsky S, Dritsa M, *et al.* Determinants of sleep quality in women with systemic lupus erythematosus. *Arthritis Rheum* 2005;53:272–8.
- 41 Iaboni A, Ibanez D, Gladman DD, *et al.* Fatigue in systemic lupus erythematosus: contributions of disordered sleep, sleepiness, and depression. *J Rheumatol* 2006;33:2453–7.
- 42 McKinley PS, Ouellette SC, Winkel GH. The contributions of disease activity, sleep patterns, and depression to fatigue in systemic lupus erythematosus. A proposed model. *Arthritis Rheum* 1995;38:826–34.
- 43 van Vollenhoven R, Voskuyl A, Bertsias G, *et al.* A framework for remission in SLE: consensus findings from a large international task force on definitions of remission in SLE (DORIS). *Ann Rheum Dis* 2017;76:554–61.
- 44 Tench CM, McCarthy J, McCurdie I, *et al.* Fatigue in systemic lupus erythematosus: a randomized controlled trial of exercise. *Rheumatology* 2003;42:1050–4.
- 45 Arnaud L, Mertz P, Amoura Z, *et al.* Patterns of fatigue and association with disease activity and clinical manifestations in systemic lupus erythematosus. *Rheumatology* 2020. doi:10.1093/rheumatology/keaa671. [Epub ahead of print: 11 Nov 2020].
- 46 Keyser RE, Rus V, Cade WT, *et al.* Evidence for aerobic insufficiency in women with systemic lupus erythematosus. *Arthritis Rheum* 2003;49:16–22.
- 47 Margiotta DPE, Basta F, Dolcini G, *et al.* Physical activity and sedentary behavior in patients with systemic lupus erythematosus. *PLoS One* 2018;13:e0193728.
- 48 Geenen R, Overman CL, Christensen R, *et al.* EULAR recommendations for the health professional's approach to pain management in inflammatory arthritis and osteoarthritis. *Ann Rheum Dis* 2018;77:797–807.
- 49 Balsamo S, Santos-Neto LD. Fatigue in systemic lupus erythematosus: an association with reduced physical fitness. *Autoimmun Rev* 2011;10:514–8.
- 50 Mahieu MA, Ahn GE, Chmiel JS, *et al.* Fatigue, patient reported outcomes, and objective measurement of physical activity in systemic lupus erythematosus. *Lupus* 2016;25:1190–9.
- 51 Yuen HK, Cunningham MA. Optimal management of fatigue in patients with systemic lupus erythematosus: a systematic review. *Ther Clin Risk Manag* 2014;10:775–86.
- 52 Fangtham M, Kasturi S, Bannuru RR, *et al.* Non-pharmacologic therapies for systemic lupus erythematosus. *Lupus* 2019;28:703–12.
- 53 Bisung E, Elliott SJ, Clarke AE. Non-pharmacological interventions for enhancing the working life of patients with lupus: a systematic review. *Lupus* 2018;27:1755–6.
- 54 Conceição CTM, Meinão IM, Bombana JA, *et al.* Psychoanalytic psychotherapy improves quality of life, depression, anxiety and coping in patients with systemic lupus erythematosus: a controlled randomized clinical trial. *Adv Rheumatol* 2019;59:4.
- 55 Neill J, Belan I, Ried K. Effectiveness of non-pharmacological interventions for fatigue in adults with multiple sclerosis, rheumatoid arthritis, or systemic lupus erythematosus: a systematic review. *J Adv Nurs* 2006;56:617–35.
- 56 Arnaud L, Zahr N, Costedoat-Chalumeau N, *et al.* The importance of assessing medication exposure to the definition of refractory disease in systemic lupus erythematosus. *Autoimmun Rev* 2011;10:674–8.
- 57 Costedoat-Chalumeau N, Amoura Z, Hulot J-S, *et al.* Very low blood hydroxychloroquine concentration as an objective marker of poor adherence to treatment of systemic lupus erythematosus. *Ann Rheum Dis* 2007;66:821–4.