OR=9.3 for adults with DScores of 2–3 and 4–6 [compared with DS 0–1], respectively). Odds of CS (vs FS) were also higher with higher DScores (OR=6.9 and OR=7.6 for adults with a DS of 2–3 and 4–6 [compared with DS 0–1], respectively).

Conclusion Smoking is highly prevalent in patients with pCCLE. DScores were positively associated with CS and inversely associated with FS. Smoking cessation is particularly important for adults with pCCLE, and such efforts should target individuals from the most disadvantaged sociodemographic groups.

Acknowledgements The GOAL Cohort is supported by the Centers for Disease Control and Prevention (CDC) Grant 1U01DP005119. The content of this research is solely the responsibility of the authors and does not necessarily represent the official views of the CDC. The authors have no conflicts of interest to declare.

P58 SYSTEMIC LUPUS ERYTHEMATOSUS IN NATIVE SUB-SAHARAN AFRICANS: A SYSTEMATIC REVIEW AND META-ANALYSIS

¹Mickael Essouma, ¹Jan René Nkeke, ²Francky Teddy A Endomba, ^{3,4}Jean Joel Bigna, ^{1,5}Madeleine Singwe-Ngandeu, ⁶Eric Hachulla. ¹Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, Yaoundé, Cameroon; ²University of Bourgogne, Dijon, France; ³Centre Pasteur of Cameroon, Yaoundé, Cameroon; ⁴Faculty of Medicine, University of Paris Sud XI, Le Kremlin-Bicêtre, France; ⁵Yaoundé Central Hospital, Yaoundé, Cameroon; ⁶Claude Huriez Hospital, Lille University, Lille, France

10.1136/lupus-2020-eurolupus.105

Background This systematic review of literature and meta-analysis aimed to determine the prevalence, phenotype and treatment of systemic lupus erythematosus (SLE) in Native sub-Saharan Africans.

Methods PubMed, EMBASE, Web of Science, African Journals Online, and Global Index Medicus as well as references of retrieved papers were searched to select studies addressing SLE in Native sub-Saharan Africans and published during January 1, 2008– October 7, 2018. Results were pooled through narrative review and random-effects model. Heterogeneity (I^2) was assessed via the χ^2 test. Pooled estimates are expressed

with 95% confidence intervals. This study is registered with PROSPERO: registration number CRD42019139226.

Results Fifteen hospital-based studies were included out of 1502 records. The pooled prevalence of SLE was 1.7% (0.8–2.9). The mean age at diagnosis ranged from 28.8 to 39.2 years. The female proportion was 88%–100%. Rheumatological (5.1%–99.9%), dermatological (4.3%–100%) and hematological (1.4–86.9%) manifestations were the commonest. Patients had a high seroprevalence for anti-ribonucleoprotein 57.9% (36.4–77.9), anti-Smith 53.5% (40.4–66.2), anti-Sjogren syndrome antigen A 45.6% (19.2–73.4) and anti-Sjogren syndrome antigen B 33.7% (13.6–57.6) autoantibodies. The most used treatments were corticosteroids 99% (94.9–100) and antimalarials 62.8% (23.3–94.1). The pooled mortality rate was 10.3% (3.3–20.6); mainly due to infections, kidney and neurological involvement.

Conclusions Over the last 30 years, SLE was not rare among Native sub-Saharan Africans and its featured characteristics were earlier onset, female predominance, and high seropositivity for extractable nuclear antigen autoantibodies. The standard treatments were corticosteroids and antimalarials. The mortality rate was high. Population prevalence and incidence as well as full description of SLE characteristics in Native sub-Saharan Africans are needed.

P59 PREDICTORS OF RENAL SURVIVAL IN A COHORT OF PATIENTS WITH LUPUS NEPHRITIS WITH MORE THAN 30 YEARS OF FOLLOW-UP

¹Filipa Farinha, ²Ruth J Pepper, ³Daniel G Oliveira, ¹Thomas McDonnell, ¹David A Isenberg, ¹Anisur Rahman. ¹Centre for Rheumatology, University College London, London; ²Centre for Nephrology, University College London-Royal Free Campus, London, UK; ³Internal Medicine Dept., Centro Hospitalar e Universitário do Porto, Porto, Portugal

10.1136/lupus-2020-eurolupus.106

Background Despite the improvement in survival of patients with lupus nephritis (LN) over the last decades, LN is associated with progression to end-stage renal disease (ESRD) in a significant proportion of patients. We aimed to investigate the factors influencing renal survival in patients with LN.

Abstract P59 Figure 1 Kaplan-meier curves showing cumulative renal survival for patients with LN