P112 LUPUS NEPHRITIS IN MALES: CLINICAL AND HISTOLOGICAL PROFILE, DISEASE COURSE AND MANAGEMENT – RESULTS FROM A NEPHROLOGY CLINIC IN SRI LANKA

Duminda Basnayake, Abdul Wazil, Nishantha Nanayakkara, Mohommed Shafraz. Nephrology and Transplant Unit, Teaching Hospital Kandy, Kandy, Sri Lanka

10.1136/lupus-2020-eurolupus.156

Background Lupus Nephritis (LN) in males is poorly understood due to the low prevalence in men. Our aim was to identify the clinical and histological profile on initial presentation and to report the experience of disease course and management.

Methods Seventeen biopsy proven LN males from nephrology clinic, Teaching Hospital Kandy were retrospectively analyzed.

Results The median age at the time of renal biopsy was 25.7 years (range, 14–57 years). Commonest initial presentation was nephrotic syndrome in 10 of 17 patients followed by rapidly progressive glomerulonephritis (RPGN) in 4 and nephritic syndrome in 3.

Anemia was noted in 11 patients. Thrombocytopenia was seen among 4 patients. ANA report was positive in 13 and negative in 4 patients. Diffuse LN was the commonest histological pattern accounting to 10 cases followed by Focal LN in 4, membranous LN in 2 and mesangial proliferative LN in 1 patient.

After remission induction treatment, 10 patients achieved complete renal remission. Five patients achieved partial remission while two patients did not achieve remission. Induction was done with intravenous (IV) methyl prednisolone (0.5/1.0g) daily 3 doses and IV cyclophosphamide (750 mg/m² body surface) 2 doses two weeks' apart and 4 doses monthly. Maintenance treatment was with oral prednisolone and mycophenolate mofetil.

Median follow-up of the cohort was 52 months (2–89 months). During the follow-up repeat biopsy was performed in 4 patients. The changes observed were, class shift from class III to IV in 2 patients and from class V to class IV in one patient. Only one patient required renal replacement therapy during the follow up period. Six patients developed chronic kidney disease (CKD) after a median of 33 months follow-up.

Conclusion Commonest presentation of LN in males is nephrotic syndrome, and type IV LN is the most frequent histological form. Pattern of initial presentation is seemingly associated with the renal outcome. Better delineation needs further studies with larger cohorts.

P113 ONLINE CASE-BASED EDUCATION SUBSTANTIALLY IMPROVED RHEUMATOLOGISTS' CONFIDENCE AND COMPETENCE IN MANAGING SLE AND SELECTING APPROPRIATE TREATMENTS

¹Elaine Bell, ²Robert McCarthy, ³Ronald van Vollenhoven. ¹Articulate Medical Communications Ltd, Hitchin, UK; ²Medscape LLC, New York, USA; ³Amsterdam Rheumatology and Immunology Center ARC, Amsterdam, The Netherlands

10.1136/lupus-2020-eurolupus.157

Background/Purpose The prognosis for patients with systemic lupus erythematosus (SLE) has improved in recent years, but many patients fail to achieve optimal disease control. We assessed whether an online case-based CME activity could

improve rheumatologists' confidence in managing SLE and in selecting appropriate treatments for their patients.

Methods Rheumatologists participated in a segmented online multi-media activity consisting of videos portraying realistic physician–patient interaction followed by a test question and expert commentary. Educational effect was assessed using a repeated-pair design, pre-/post-assessment. A Chi-square test of independence determined if a statistically significant improvement (5% significance level, $P < 0.05$) existed in the number of correct responses between the pretest and posttest scores. Cramer's V estimated the effect size of the education. The activity launched 26 March 2019 with data collection through 29 April 2019.

Results

- 35% of rheumatologists' (n=46) answered all 3 questions correctly at baseline rising to 85% post assessment ($P < .001$)
- Significant improvement in knowledge of the proportion of SLE patients testing positive for anti-nuclear antibodies (72% at baseline, 93% post assessment; $P < .01$), and significant improvement in treatment choice for a patient who experienced worsening symptoms following initial success with prednisone, hydroxychloroquine and methotrexate (59% at baseline, 91% post assessment; $P < .001$)
- Extensive educational impact (Cramer's V =0.277) on knowledge and competence overall with 30% of rheumatologists reporting greater confidence in their ability to treat SLE symptoms
- 38% of participants reported an intent to make changes in their practice based on what they learned; of those who intended to make a change, 92% were highly committed to implementing these changes

Conclusions Participation in this segmented online multi-media activity was highly effective in improving rheumatologists' confidence and competence in treatment selection and management of SLE. The extensive impact of this education should translate into improvements in clinical practice.

Acknowledgement Supported by GlaxoSmithKline

P114 EVALUATION OF NEUROPSYCHIATRIC INVOLVEMENT AND PSYCHOMETRIC STATUS IN PEDIATRIC SYSTEMIC LUPUS ERYTHEMATOSUS (SLE)

Emil Aliyev, Yelda Bilginer. Pediatric Rheumatology Dept., Hacettepe University Faculty of Medicine, Turkey

10.1136/lupus-2020-eurolupus.158

Background Systemic Lupus Erythematosus (SLE) is an autoimmune disease characterized by multiple organ involvement. Neuropsychiatric involvement is seen in 20–27% of patients. Approximately 1/4 of diagnosed SLE patients develop neuropsychiatric SLE 2 years after diagnosis.

Methods The aim of this study was to determine the risk groups before developing neurological, psychiatric and cognitive involvement and symptoms in SLE patients and to take measures and to develop cheaper and more applicable algorithms.

Results Ninety eight children and adolescents were included in the study. 61 (62.2%) of the patients were female and 37 (37.8%) were male. Research group included 34 (34.7%) SLE patients, 31 (31.6%) JIA patients, and 33 (33.7%) healthy children and adolescents. The median age of the patients was 17 (12–21) years, and the median follow-up period was 4