activity of BLyS, effectively and safely treats SLE, but data on treatment cessation are lacking. Therefore, we investigated belimumab-free remission in SLE patients.

Methods SLE patients receiving belimumab in our institute (1/1/2013–5/31/2019) were retrospectively identified using electronic health records. Eligibility criteria were receiving belimumab for >180 days and discontinuation for any reason. BILAG category A or B in at least one organ system defined disease flares. Follow-up monitoring after 52 weeks post-treatment identified relapse-free and relapse patients.

Results 31 patients received belimumab. While 14 patients discontinued, eight were included. Four patients relapsed within 52 weeks. Relapse-free patients received significantly less steroid at discontinuation (prednisolone equivalent, median 3.0 mg/day [IQR 2.75–3.19] vs. 9.5 mg/day [IQR 7.25–13.25], $p=0.02$), and significantly more of them achieved PSL dosage of <5 mg/day on discontinuation day than relapse patients. ($p=0.03$) At belimumab discontinuation, relapse-free patients tended to have higher C3 (median 91.0 mg/dL [IQR 78.75–102.25] vs. 56.0 mg/dL [IQR 39.75–73.00], $p=0.15$) and C4 levels (median 22.0 mg/dL [IQR 19.00–26.00] vs. 11.0 mg/dL [IQR 6.00–16.00], $p=0.08$) and less anti-DNA antibody (median 5.2 IU/mL [IQR 3.75–7.83] vs. 48.0 IU/mL [IQR 11.50–137.25], $p=0.08$) than relapse patients, but differences were not significant.

Conclusion Belimumab discontinuation after >180 days is recommended for 50% of SLE patients. Steroid dosage (prednisolone equivalent <5 mg/day) might be a prognostic marker for belimumab-free remission.

Funding The authors received no financial support for the research, authorship, and/or publication of this article.

P121

IDENTIFICATION AND ACTIVE MANAGEMENT OF SECONDARY ADRENAL INSUFFICIENCY IN SLE REDUCES RISK OF RELAPSE

¹Benjamin Rhodes, ²Haroon Ahmad, ³Sarah Logan, ¹Elizabeth Rankin, ³Peter Hewins. ¹Rheumatology Dept., Queen Elizabeth Hospital, Birmingham; ²University of Birmingham Medical School, Birmingham; ³Nephrology Dept., Queen Elizabeth Hospital, Birmingham, UK

10.1136/lupus-2020-eurolupus.165

Background Steroids are an important component of Lupus treatment, but long-term use is associated with an accumulation of damage, emphasising the importance of optimising steroid withdrawal protocols. The factors influencing successful steroid withdrawal in SLE are poorly understood. Secondary adrenal insufficiency is common following therapeutic steroid usage, but the prevalence in steroid-treated lupus patients is undocumented. We documented rates of adrenal insufficiency and evaluated whether actively managing steroid withdrawal influenced the risk of disease relapse.

Methods 137 SLE patients who had been on Prednisolone for more than a year before reducing the dose below 5 mg/day for the first time were retrospectively reviewed, documenting the clinician decision to screening for adrenal insufficiency, the frequency of adrenal insufficiency, the method of steroid reduction (stopping vs. tapering regimens), clinical, pharmacological and immunological variables. Parameters influencing the risk of relapse from the point of dropping below 5 mg/day of Prednisolone were evaluated by Cox' regression.

Results 65 patients were screened for adrenal insufficiency. 38% failed initial screening and 12% showed no signs of adrenal recovery after more than a year. 46.7% returned to higher dose steroids, with 34% having a true disease relapse (BILAG A or B flare). In a multivariate model the decision to screen for adrenal insufficiency was the variable most strongly associated with risk of disease relapse (BILAG A/B flare) (marginal Hazard Ratio 4.33, $P=0.0001$), followed by complement C4 at baseline and South Asian ancestry. Concomitant medications or the method of steroid reduction (stopping vs. tapering) did not influence relapse. A full analysis of factors associated with failed adrenal screening and risk of disease relapse will be presented.

Conclusions Adrenal insufficiency in common in patients with SLE and unless screening takes place this may go undetected. Detection and active management of adrenal insufficiency is associated with a significant and meaningful reduction in disease relapse.

P122

WHAT DO PATIENTS WITH LUPUS AND SJÖGREN'S SYNDROME KNOW ABOUT CARDIOVASCULAR RISK?

Elvira Chocano Navarro, George Robinson, Kirsty Waddington, Thomas McDonnell, Chris Wincup, Lucia Martin-Gutierrez, Annalisa Maggio, Eve McLoughlin, Lizzy Rosser, David Isenberg, Anastasia Kalea, Coziana Ciurtin, Ines Pineda-Torra, Elizabeth Jury. Dept. of Medicine, University College London, London, UK

10.1136/lupus-2020-eurolupus.166

Background Cardiovascular risk (CVR) is the leading cause of mortality in patients with lupus. Understanding increased CVR in autoimmune rheumatic diseases could improve management of risk in patients. This event aimed to explore patient opinions about CVR and potential CVR treatment options.

Methods We hosted a patient event promoted through social media, relevant charities, hospitals and research groups. 13 patients with lupus and/or Sjögren's syndrome attended and were asked about CVR using a questionnaire (14 questions) and round table discussion with patients, researchers, clinicians and dietitians.

Results Sixty percent of patients were aware of the increased CVR associated with autoimmune rheumatic disease and 60% stated that their doctor had spoken to them about this risk. 73% thought that it was important for them to be aware of this increased CVR.

When asked about medication to reduce CVR, no patients wanted to take a statin (lipid lowering drug), however, 70% of patients would take statins if advised to do so by their doctor. Conversely, respondents were more positive about using diet or taking a dietary supplement to reduce CVR; 71% would change their diet and 57% would take a supplement either on their own accord or on advice from health professionals. Some patients had already made changes to their diet to reduce their CVR, including reducing fat and increasing fruit and vegetable consumption. All attendees were prepared to participate in a clinical study using diet modification strategies, having vascular scans to assess atherosclerosis and provide blood samples for CVR research in lupus/Sjögren's syndrome.

Conclusion This multidisciplinary event successfully gathered patient information regarding CVR. The opinions and comments provided evidence that patients support further research in cardiovascular studies and a preference to changing their diet or take a supplement, whilst avoiding medication, to reduce their CVR.