recent *in vivo* pathogenesis studies in mice, have shown that intestinal colonization by RG strains isolated from Lupus patients in flare, but not RG strains from healthy adults, induce zonulin-dependent increases in intestinal permeability, RG translocation to mesenteric lymph nodes as well as serum IgG anti-RG antibody and anti-native DNA autoantibody responses.

Conclusions As many Lupus patients are known to suffer relapsing illness despite appropriate treatment, we speculate that gut blooms of pathogenic bacteria that impair gut barrier function and stoke systemic inflammation directly contribute to immunopathogenesis. We propose that future therapeutic interventions designed to promote a durable remission consider the potential necessity to target both the immunologic abnormalities of the disease, as well as re-establish stability within the gut microbiota community.

Acknowledgments This work was supported in part by National Institutes of Health Grants; R01-AR42455, P50 AR070591, the Lupus Research Alliance, and the Judith and Stewart Colton Autoimmunity Center (GJS). 16S rRNA amplicon sequence determinations and analysis were supported by the P. Robert Majumder Charitable Trust.

512

THE RISING INCIDENCE, PREVALENCE AND MORTALITY GAP OF LUPUS NEPHRITIS: A POPULATION-BASED STUDY OVER FOUR DECADES

^{1,2}Mehmet Hocaoglu, ¹Maria O Valenzuela-Almada, ¹Jesse Y Dabit, ¹Shirley-Ann Osei-Onomah, ^{1,3}Baptiste Chevet, ⁴Rachel E Giblon, ^{1,4}Cynthia S Crowson, ^{1,5}Ali Duarte-Garcia*. ¹Division of Rheumatology, Mayo Clinic, Rochester, MN, USA; ²Department of Medicine, University of Maryland Medical Center, Midtown Campus, Baltimore, MD, USA; ³Division of Rheumatology, Brest Teaching Hospital, Brest, France; LBAI, UMR1227, Univ Brest, Inserm, CHU de Brest, Brest, France; ⁴Department of Quantitative Health Sciences, Mayo Clinic, Rochester, MN, USA; ⁵Division of Nephrology and Hypertension, Mayo Clinic, Rochester, MN, USA

10.1136/lupus-2021-lupus21century.30

Background There is paucity of population-based studies that investigate the epidemiology of lupus nephritis (LN) and its long-term outcomes.

Methods We used the Rochester Epidemiology Project (REP) a record-linkage system that includes all residents from Olmsted County, Minnesota since 1966 with additional counties since 2010. A cohort of LN patients between 1976 and 2018 in an eight-county region was identified based on (1) biopsy proven LN in the presence of a positive ANA or ds-DNA antibody, or (2) meeting ACR/EULAR 2019 criteria and having documented proteinuria (500 mg in 24hrs urine collection or protein/creatinine ratio > 0.5) that did not have a better

explanation than SLE. Age- and sex-specific incidence rates and point prevalence for four decades, adjusted to 2000 US total population, were reported. Standardized mortality ratio (SMR), survival rates, and time to end-stage renal disease (ESRD) were estimated using Kaplan-Meier methods. Poisson regression models were used to look at trends in relative survival over time.

Results There were 72 incident LN from 1976-2018. Mean age was 38.4 years (SD 16.24), 76% were female, and 69% non-Hispanic White. The overall LN incidence between 1976 and 2018 was estimated at 1 per 100,000 (95%CI 0.8-1.3). The incidence was highest in 30-39 age group. The overall incidence of LN increased from 0.7 to 1.3 with sex specific incidence in males quadrupling from 0.2 to 0.8 per 100,000 between 1976-1988 and 2000-2018 periods (table 1). LN prevalence increased from 16.8 per 100,000 in 1985 to 21.2 in 2015. LN had SMR of 6.33 (95%CI 3.81-9.89) compared to the general population. Survival was 50% at 20 years, and among survivors 40% had ESRD. There was a widening mortality gap with an increase in the SMR of 7% per year (1.07; 95%CI 1.03-1.10, p<0.001) or almost a doubling of the SMR every 10 years (1.90 per 10 years; 95%CI 1.33-2.70).

Conclusions The incidence and prevalence of LN has increased in the last 40 years. Long-term outcomes in patients with LN remain poor. Patients with LN had a mortality rate of more than 50% at 20 years and of those surviving, 40% had ESRD. The widening mortality gap we observed with near doubling of SMR each decade shows that the mortality reduction in LN patients is not keeping up with the mortality improvements of the general population despite changes in clinical practice over the decades.

513

RESPONSE GENE TO COMPLEMENT -32 FACILITATES LOCAL RECRUITMENT OF IL-17- PRODUCING CELLS IN IMMUNE COMPLEX MEDIATED GLOMERULONEPHRITIS THROUGH THE CCR6/CCL20 AXIS

^{1,2}Violeta Rus*, ¹Alexandru Tatmir, ^{1,2}Vinh Nguyen, ¹John Papadimitriou, ¹Sergei Atamas, ^{1,2}Irina G Luzina, ³Tudor Badea, ⁴Sonia Vlaicu, ^{1,2}Horea Rus. ¹University of Maryland School of Medicine, Baltimore, MD, USA; ²VA Maryland Health Care System, Baltimore, MD, USA; ³NIH, NEI, Bethesda, MD, USA; ⁴University of Medicine and Pharmacy 'Iuliu Hatieganu', Cluj-Napoca, Romania

10.1136/lupus-2021-lupus21century.31

Background Response Gene to Complement (RGC)-32 is a cell cycle regulator induced by complement activation, growth factors and cytokines. RGC-32 mediates TGF- β dependent profibrotic pathways, while in immune cells promotes the

Abstract 512 Table 1 Age-adjusted incidence (1976-2018) and prevalence (1985,1995,2005,2015) rates of Lupus Nephritis (LN), 8-county region in Southeast Minnesota.[†]

	LN incidence rates/100,000 (95% CI) *					LN Prevalence rates/100,000 (95% CI) *				
	1976-1989	1990-1999	2000-2009	2010-2018	1976-2018	01/01/1985	01/01/1995	01/01/2005	01/01/2015	
Females	1.2 (0.4, 2.0)	1.3 (0.4, 2.2)	1.7 (0.7, 2.7)	1.7 (1.0, 2.4)	1.5 (1.1, 1.9)	25.9 (11.4, 40.5)	30.7 (16.7, 44.7)	36.5 (22.4, 50.7)	33.4 (24.7, 42.1)	
Males	0.2 (0.0, 0.6)	0.0 (0.0, 0.0)	0.6 (0.0, 1.2)	0.8 (0.4, 1.3)	0.5 (0.3, 0.8)	7.1 (0.0, 15.1)	6 (0.0, 13)	1.4 (0.0, 4.3)	8.6 (4.1, 13)	
Overall	0.7 (0.2, 1.2)	0.6 (0.2, 1.1)	1.2 (0.6, 1.8)	1.3 (0.9, 1.7)	1.0 (0.8, 1.3)	16.8 (8.3, 25.3)	18.5 (10.7, 26.4)	19.4 (12.1, 26.8)	21.2 (16.3, 26.2)	

*Rates for females and males were age-adjusted to the 2000 US Total population. Overall rates were age- and sex-adjusted to the 2000 US Total population.

[†]8-county region from Minnesota (Olmsted, Mower, Freeborn, Waseca, Steele, Dodge, Wabasha, Goodhue) Incidence rates were calculated using the number of incident cases as the numerator and population counts for residents from the Rochester Epidemiology Project (REP) census as the denominator. Incident and prevalent cases were from Olmsted County region between 1976 and 2010 and the 8 counties from 2010-2018.