Conclusions Our cohort showed high rate of patient satisfaction with telemedicine healthcare. However, the relatively low healthcare provider satisfaction rate raises concern as to whether telemedicine constitutes a satisfactory alternative to conventional in-person care. Additional researches are required to investigate the feasibility of telemedicine in long-term disease activity evaluation and patient outcome measurement.

1106

PREDICTING ADVERSE PREGNANCY OUTCOMES IN WOMEN WITH SYSTEMIC LUPUS ERYTHEMATOSUS: A COMPARISON OF MACHINE LEARNING METHODS

¹Melissa Fazzari, ²Marta Guerra, ^{2,3}Jane Salmon, ¹Mimi Kim*. ¹Albert Einstein College of Medicine, New York, USA; ²Hospital for Special Surgery, New York, USA; ³Weill Cornell Medical College, New York, USA

10.1136/lupus-2021-lupus21century.49

Background Nearly 20% of pregnancies in patients with Systemic lupus erythematosus (SLE) result in an adverse pregnancy outcome (APO); early identification of women with SLE who are at high risk of APO is vital. We previously derived a risk model for APO using logistic regression and data from the PROMISSE Study, a large multi-center, multi-ethnic/racial study of APO in women with mild/moderate SLE and/or aPL. While this highly interpretable regression model showed promising predictive performance, we sought to determine if novel and increasingly popular machine learning (ML) approaches would enhance APO risk prediction using all available predictors and potential complex relationships such as interactions or higher order terms. We compared logistic regression modeling to LASSO, a regression approach that handles high-dimensionality and correlated predictors through shrinkage of estimated coefficients, as well as several 'black box' ML algorithms. ML techniques are well-suited to high-dimensional data, require no variable selection, and unlike regression-based approaches are able to explore complex relationships without explicit input by the user.

Methods We used the original PROMISSE data (41 predictor variables from 385 subjects) with APO (71/385, 18.4%) defined as preterm delivery due to placental insufficiency or preeclampsia, fetal or neonatal death, or fetal growth restriction. Logistic regression with stepwise selection (LR-S) was compared to LASSO, random forest (RF), neural network (NN) with 2 hidden neurons, support vector machines with RBF kernel (SVM_{RBF}), and gradient boosting (GB). To summarize discrimination we present the area under the receiver operating curve (AUC), along with sensitivity (Sn) and specificity (Sp) at an optimal cut-point.

Results Regression based classifiers confirmed the predictors of APO identified in our previously reported model: non-white race, use of anti-hypertensive medication, low platelets, SLE disease activity, lupus anticoagulant (LAC) +, and high diastolic blood pressure (DBP). RF additionally revealed two novel interaction variables that increased APO risk: LAC+ with anti-β2GPI IgM, high DBP with low C3. LR-S and LASSO were observed to have similar overall discrimination (AUC=0.75 vs. 0.77, table 1) but LASSO had higher sensitivity (Sn=0.71 vs. 0.65). ML classifiers RF and SVM_{RBF} had similar good performance (AUC=0.77-0.78), while NN and GB were inferior.

Conclusions Several popular ML algorithms did not provide meaningful improvements in the prediction of APO. The strong relative performance of regression-based models with

Abstract 1106 Table 1 Summary of 5x10 fold cross-validation results

Model	AUC	Sensitivity	Specificity
LR-S	0.75	0.65	0.78
LASSO	0.77	0.71	0.75
NN	0.71	0.70	0.61
RF	0.77	0.77	0.78
GB	0.72	0.72	0.72
SVM-RBF	0.78	0.78	0.73

this large and well-characterized clinical data set is notable as these models are highly interpretable, well-understood, and generally require fewer variables to generate a risk prediction. It is unlikely that complex ML algorithms with existing variables will yield superior APO predictions; new clinical and laboratory markers may improve predictions in the future.

Acknowledgments This work was supported by NIH grant R21 AR076612

1107

ECONOMIC EVALUATION OF HYDROXYCHLOROQUINE USE IN AN INTERNATIONAL INCEPTION COHORT

¹Megan RW Barber, ²Yvan St Pierre, ³John G Hanly, ⁴Murray B Urowitz, ⁵Caroline Gordon, ⁶Sang-Cheol Bae, ⁷Juanita Romero-Díaz, ⁴Jorge Sanchez-Guerrero, ⁸Sasha Bernatsky, ⁹Daniel J Wallace, ¹⁰David A Isenberg, ¹⁰Anisur Rahman, ¹¹Joan T Merrill, ¹²Paul R Fortin, ⁴Dafna D Gladman, ¹³Ian N Bruce, ¹⁴Michelle Petri, ¹⁵Ellen M Ginzler, ¹⁶Mary Anne Dooley, ¹⁷Rosalind Ramsey-Goldman, ¹⁸Susan Manzi, ¹⁹Andreas Jönsen, ²⁰Graciela S Alarcón, ²¹Ronald F Van Vollenhoven, ²²Cynthia Aranow, ²²Meggan Mackay, ²³Guillermo Ruiz-Irastorza, ²⁴Sam Lim, ²⁵Murat Inanc, ²⁶Kenneth C Kalunian, ²⁷Søren Jacobsen, ²⁸Christine A Peschken, ²⁹Diane L Kamen, ³⁰Anca Askanase, ¹Ann E Clarke*. ¹University of Calgary, Alberta, Canada; ²Research Institute of the McGill University Health Center, Montreal, Canada; ³Queen Elizabeth II Health Sciences Centre and Dalhousie University, Halifax, Nova Scotia, Canada; ⁴Centre for Prognosis Studies in the Rheumatic Diseases, Toronto Western Hospital and University of Toronto, Toronto, Ontario, Canada; ⁵Rheumatology Research Group, Institute of Inflammation and Ageing, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK; ⁶Hanyang University Hospital for Rheumatic Diseases, Seoul, Korea; ⁷Instituto Nacional de Ciencias Médicas y Nutrición, Mexico City, Mexico; ⁸McGill University Health Centre, Montreal, Canada; ⁹Cedars-Sinai/David Geffen School of Medicine at the University of California, Los Angeles, USA; ¹⁰University College London, London, UK; ¹¹Department of Clinical Pharmacology, Oklahoma Medical Research Foundation, Oklahoma City, OK, USA; ¹²CHU de Québec – Université Laval, Québec City, Canada; ¹³Arthritis Research UK Epidemiology Unit, Institute of Inflammation and Repair, Manchester Academic Health Sciences Centre, the University of Manchester, and NIHR Manchester Musculoskeletal Biomedical Research Unit, Central Manchester University Hospitals National Health Service Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK; ¹⁴Johns Hopkins University School of Medicine, Baltimore, Maryland, USA; ¹⁵State University of New York Downstate Medical Center, Brooklyn, New York, USA; ¹⁶Thurston Arthritis Research Center, University of North Carolina, Chapel Hill, NC, USA; ¹⁷Northwestern University and Feinberg School of Medicine, Chicago, Illinois, USA; ¹⁸University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA; ¹⁹Lund University, Lund, Sweden; ²⁰University of Alabama at Birmingham, USA; ²¹University of Amsterdam, Rheumatology and Immunology Center, Amsterdam, Noord-Holland, NL; ²²Feinstein Institute for Medical Research, Manhasset, New York, USA; ²³BioCruces Health Research Institute, Hospital Universitario Cruces, University of the Basque Country, Barakaldo, Spain; ²⁴Emory University School of Medicine, Atlanta, Georgia, USA; ²⁵Istanbul University, Istanbul, Turkey; ²⁶University of California Los Angeles School of Medicine, La Jolla, California, USA; ²⁷Copenhagen Lupus and Vasculitis Clinic, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark; ²⁸University of Manitoba, Winnipeg, Manitoba, Canada; ²⁹Medical University of South Carolina, Charleston, USA; ³⁰Hospital for Joint Diseases, New York University Seligman Center for Advanced Therapeutics, New York, New York, USA

10.1136/lupus-2021-lupus21century.50