physicians (dermatology or rheumatology). Quota sampling was used to recruit patients from racially and ethnically diverse backgrounds and physicians with varied medical practices. Patient and physician participants provided informed consent. Semi-structured interviews explored patients' experiences with the healthcare system and physicians' challenges when treating patients with lupus.

Results Interviews were conducted with 33 patients and 20 physicians. Age, region, education, employment status, and type/severity of lupus varied among patients, as did physicians' work experience and practice settings (table 1). Patients reported challenges at each step of their journey. Prior to diagnosis, patients entered the healthcare system often lacking awareness of lupus disease and reported feeling initially ignored, dismissed, or misdiagnosed, which delayed treatment and triggered mistrust of the healthcare system. Lack of trust was cited by patients as a reason to discontinue treatment, despite persistent symptoms. Some patients reported perceived challenges with access to branded medications due to insurance coverage or delays in prior authorization and were concerned about losing their job and insurance. Patients communicated that treatment access might be improved by establishing support systems for reliable disease information (advocacy groups, community support), increasing cultural sensitivity in physician practices, and enabling an efficient and transparent treatment initiation process. From the physician perspective, barriers to treatment access included delayed diagnoses, patient concerns over side effects and the need for life-long treatment, communication challenges with patients with limited English proficiency (LEP), and medication costs. Physicians suggested addressing perceived barriers by providing better patient information resources, having a live translator present during visits for patients with LEP, and navigating accessibility to medications through patient assistance programs to ensure access to branded therapies when needed.

Conclusions Improving access to quality care for underrepresented patients is essential to reduce health disparities across racial/ethnic groups and build trust in the healthcare system. Increasing disease awareness among these diverse groups and physicians, establishing patient support networks, and improving patient-physician communication should be critical components of health equity efforts.

REFERENCES

1. Arora. *Rheum Dis Clin N Am*. 2020;46:623-638.
2. Maningding. *Arthritis Care Res*. 2020;72(5):622-629.

1112

IMPACT STUDY: CAN WE CONDUCT A TRIAL WITH A BIOLOGIC TO PREVENT PREECLAMPSIA IN WOMEN WITH ANTIPHOSPHOLIPID SYNDROME?

¹Jane E Salmon*, ¹Marta Guerra, ²Mimi Kim, ³D Ware Branch. ¹Hospital for Special Surgery, USA; ²Albert Einstein College of Medicine, USA; ³University of Utah, USA

10.1136/lupus-2021-lupus21century.55

Background Pregnant women with antiphospholipid antibodies and/or lupus have higher rates of adverse pregnancy outcomes (APOs), such as fetal loss and preterm birth due to severe preeclampsia (PE) or placental insufficiency (PI). The presence of lupus anticoagulant (LAC) is the strongest predictor of an APO. At present, there is no effective treatment for women with these high-risk pregnancies, but in an animal model that mimics this human condition we found that TNF- α was a

critical downstream effector of abnormal placental development and fetal damage, and that TNF- α blockade normalized placentation and spiral artery remodeling, and rescued pregnancies. We sought to determine whether TNF- α blockade during pregnancy, added to a regimen of heparin and low dose aspirin, reduces the rate of APOs in women with clinical APS and LAC.

Methods The IMPACT Study (IMProve Pregnancy in APS with Certolizumab Therapy) is an open label single-stage Phase II trial to evaluate the effect of certolizumab, a TNF- α inhibitor that does not cross the placenta and has been shown to be well tolerated in pregnancy, to reduce the risk of adverse outcomes in this population. Patients with APS and LAC are referred to IMPACT by their physicians (often before a planned pregnancy), consented and screened remotely by a study investigator, and medication is sent to the patient. They are treated with certolizumab from gestational week 8 through 28. Investigators contact patients every 2 weeks and receive medical reports and research blood samples monthly. Without the intervention being studied, 44% of these pregnancies were predicted to have serious complications, including severe preterm preeclampsia, growth restricted fetuses, and/or fetal death. Assuming a target 50% reduction in APO rate with the intervention, forty-five evaluable pregnancies are required for 90% power to prove that the intervention reduces the rate of adverse outcomes.

Results Since May 2017, we have enrolled 33 patients from nine states. Characteristics: 61% had previous PE or PI <34 weeks requiring delivery, 70% had previous fetal death >10 weeks, 12% had neonatal death due to complications of prematurity; 42% venous thrombosis, 24% Stroke/TIA, 24% SLE. There have been no study drop-outs.

Conclusion We are successfully using a 'rare disease study' approach to conduct the first trial of a biologic therapy to prevent pregnancy complications women with APS and LAC. For rare conditions such as this, one must target assorted specialists (maternal-fetal medicine, hematology, rheumatology) for recruitment. Identifying patients preconceptionally is ideal, because improving placental vascularization requires enrollment early in pregnancy, and screening evaluations take time. Despite regulatory and logistic complexities and the small number of patients meeting inclusion criteria, their enthusiasm and that of their physicians has allowed us to move forward, albeit slowly.

Acknowledgements NIAMS, Lupus Research institute, Lupus Foundation of America.

Trial Registrations ClinicalTrials.gov NCT03152058.

1113

PATIENT AND PHYSICIAN PERSPECTIVES OF LUPUS FLARE

¹Jennifer L Rogers*, ¹Megan EB Clowse, ²Kevin McKenna, ²Summer Starling, ²Teresa Swezey, ²Nneka Molokwu, ²Amy Corneli, ³David S Pisetsky¹Kai Sun, ¹Lisa G Criscione-Schreiber, ¹Rebecca E Sadun, ¹Mithunan Maheswaranathan, ¹Jayanth Doss, ¹Amanda M Eudy. ¹Department of Medicine, Duke University Medical Center, USA; ²Department of Population Health Sciences, Duke University, USA; ³Durham VA Medical Center, Duke University Medical Center, USA

10.1136/lupus-2021-lupus21century.56

Background Lupus flares can be associated with increased damage, poor outcomes and decreased health-related quality of life. Patients and providers may differ about the nature of a flare, however, complicating communication and