differences in disease development. Female CD103DC have decreased ability to convert conventional CD4 cells to Foxp3⁺ Tregs due to defects in retinoic acid (RA) production; female macrophages have decreased ability to mediate efferocytosis and defects in genes associated with PPAR γ and LXR signaling. Male cecal transplants can restore function of both CD103DC and macrophages in female recipients. Metabolomics analysis of feces revealed that males have higher phytanic acid (PA), an RXR/PPAR γ /LXR agonist. Treatment of female mice *in vivo* with PA delays disease to some extent and restores both CD103DC and macrophage function. PA also restores CD103DC and macrophage function when added to female cells directly *in vitro*. PA enhances CD103DC function by restoring RA production via the RXR pathway and macrophage efferocytosis via the PPAR γ and LXR pathways in female mice.

Conclusions We have identified specific gut microbiota taxa and metabolite candidates from lupus-resistant male mice that have immunomodulatory activity and the potential to mitigate disease in lupus-prone female mice. Disease mitigation may be mediated through an increase in CD103DC function and macrophages efferocytosis.

Acknowledgments Funded by ALR and NIH-R01AR067188. 16S rRNA gene sequencing was performed by GTAC, Washington University, St. Louis. Mass spectrometry was performed by CREAM, University of Louisville, Louisville.

1406 ACELLULAR ARTIFICIAL ANTIGEN-PRESENTING CELLS INDUCE DISEASE-PROTECTIVE IMMUNOREGULATORY CELLS IN LUPUS

¹Antonio La Cava*, ²David A Horwitz. ¹University of California Los Angeles, Los Angeles, CA, USA; ²General Nanotherapeutics, Santa Monica, CA, USA

10.1136/lupus-2021-lupus21century.83

Background T regulatory cells (Tregs) provide beneficial effects in the suppression of autoimmune responses in systemic lupus erythematosus (SLE). We recently showed that poly(lactic-co-glycolic acid) PLGA nanoparticles (NPs) loaded with IL-2 and TGF- β could induce CD4⁺ and CD8⁺ Foxp3⁺ Tregs *in vitro* and *in vivo*, with resulting protective effects in mice that develop lupus-like disease following the transfer of donor T cells from DBA/2 mice into (C57BL/6 x DBA/2)F₁ (BDF1) recipients.

Methods We produced NPs encapsulating IL-2 as acellular tolerogenic artificial antigen-presenting cell (aAPC)s for the induction of CD4⁺ and CD8⁺ Foxp3⁺ Tregs *in vivo*. aAPC NPs were also engineered to promote the tolerogenic activity of NK cells. Those acellular aAPCs were then tested for therapeutic efficacy *in vivo* in lupus-prone BDF1 mice.

Results The aAPC NPs induced CD4⁺ and CD8⁺ Tregs *in vivo* in BDF1 mice and protected the animals from lupus

disease manifestations. At the NK cell level, protection from lupus-like disease could be ascribed to the production of TGF- β from the NP-induced NK cells. Finally, the use of aAPC NPs in humanized NSG mice expanded Tregs in numbers sufficient to protect mice from a rapidly evolving human anti-mouse graft-versus-host disease.

Conclusions These studies provide a proof-of-concept for the potential use of PLGA NPs as acellular aAPCs for the induction of therapeutically effective numbers of Tregs in conditions of pathologic deficit such as in SLE.

1500 – Genetics

1501 GENETICS OF AGE AT SYSTEMIC LUPUS ERYTHEMATOSUS DIAGNOSIS

¹Raffaella Carlomagno, ¹Fangming Liao, ²Jingjing Cao, ¹Daniela Dominguez, ³Dafna D Gladman, ⁴Mariko Ishimori, ⁴Caroline Jefferies, ⁵Diane L Kamen, ⁶Sylvia Kamphuis, ⁷Marisa S Klein-Gitelman, ¹Andrea M Knight, ⁴Chia-Chi J Lee, ¹Deborah M Levy, ⁸Karen B Onel, ²Andrew D Paterson, ⁹Christine A Peschken, ¹⁰Janet E Pope, ³Zahi Touma, ³Murray B Urowitz, ⁴Daniel J Wallace, ¹Declan Webber, ³Joan E Wither, ¹Earl D Silverman, ¹Linda T Hiraki*. ¹Division of Rheumatology; ²Genetics and Genome Biology, Research Institute, The Hospital for Sick Children; ³Schroeder Arthritis Institute, Krembil Research Institute, Toronto Western Hospital, Toronto, Canada; ⁴Division of Rheumatology, Department of Medicine, Cedars Sinai Medical Center, Los Angeles; ⁵Division of Rheumatology and Immunology, Medical University of South Carolina, Charleston, USA; ⁶Department of Pediatric Rheumatology, Sophia Children's Hospital, Erasmus University Medical Center, Rotterdam, Netherlands; ⁷Division of Rheumatology, Department of Pediatrics, Ann and Robert H. Lurie Children's Hospital of Chicago, Chicago; ⁸Pediatric Rheumatology, Hospital for Special Surgery, New York, USA; ⁹Departments of Medicine and Community Health Sciences, University of Manitoba, Winnipeg; ¹⁰Department of Medicine, University of Western Ontario, St. Joseph's Health Centre, London; ¹¹Division of Translational Medicine Research Institute, The Hospital for Sick Children, Toronto, Canada

10.1136/lupus-2021-lupus21century.84

Background Genome wide association studies (GWAS) have identified >100 SNPs associated with systemic lupus erythematosus (SLE) risk. There may be additional loci impacting the age of diagnosis. The purpose of this study was to identify genetic variants associated with age of SLE diagnosis.

Methods Our cohort included patients from tertiary care centres who met ACR and/or SLICC SLE classification criteria. We censored patients missing age at diagnosis. Patients were genotyped on the Illumina Multiethnic Array (MEGA) or Global Screen Array (GSA). Ungenotyped SNPs were imputed using the TopMed reference. We restricted to SNPs with a minor allele frequency (MAF) $\geq$ 0.01 and imputation quality $R^2 \geq$ 0.3. Ancestry was genetically inferred using principal components (PCs) and ADMIXTURE in reference to 1000 Genome Project. We completed genome-wide linear regression of log-transformed age at SLE diagnosis with GENESIS (significance $P < 5 \times 10^{-8}$). Multivariate models adjusted for sex and 5 PCs. We also conducted a GWAS of childhood-onset SLE

Abstract 1406 Figure 1