


# Cytokine autoantibodies in SARS-CoV-2 prepandemic and intrapandemic samples from an SLE cohort

May Y Choi <sup>1,2</sup>, Ann Elaine Clarke,<sup>1</sup> Katherine Buhler,<sup>1</sup> Michelle Jung,<sup>1</sup> Hannah Mathew,<sup>1</sup> Meifeng Zhang,<sup>1</sup> Francesca S Cardwell,<sup>3</sup> Heather Waldhauser,<sup>1</sup> Marvin J Fritzler<sup>1</sup>

**To cite:** Choi MY, Clarke AE, Buhler K, *et al.* Cytokine autoantibodies in SARS-CoV-2 prepandemic and intrapandemic samples from an SLE cohort. *Lupus Science & Medicine* 2022;**9**:e000667. doi:10.1136/lupus-2022-000667

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/lupus-2022-000667>).

AEC and MJF are joint senior authors.

Received 21 January 2022  
Accepted 29 March 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Medicine, University of Calgary Cumming School of Medicine, Calgary, Alberta, Canada

<sup>2</sup>McCaig Institute for Bone and Joint Health, Calgary, Alberta, Canada

<sup>3</sup>Geography and Environmental Management, University of Waterloo, Waterloo, Ontario, Canada

## Correspondence to

Dr May Y Choi; [may.choi@ucalgary.ca](mailto:may.choi@ucalgary.ca)

## ABSTRACT

Cytokine autoantibodies, particularly those directed to type I interferon (T1IFN), have been reported to portend an increased risk of severe COVID-19. Since SLE is one of the conditions historically associated with T1IFN autoantibodies, we sought to determine the prevalence of cytokine autoantibodies in our local cohort of 173 patients with SLE prepandemic and intrapandemic, of which nine had confirmed exposure to SARS-CoV-2. Autoantibodies to 16 different cytokines, including T1IFN, were measured by an addressable laser bead immunoassay. None of the 9 patients with confirmed exposure to SARS-CoV-2 had autoantibodies to T1IFN and none had severe COVID-19 symptoms, necessitating hospitalisation. Hence, we could not confirm that T1IFN autoantibodies increase the risk for severe COVID-19. In addition, the cytokine autoantibody pattern did not differ between those with and without evidence of SARS-CoV-2 exposure.

## INTRODUCTION

Autoantibodies directed to cytokines, including type I interferon (T1IFN), have been reported in a wide variety of infectious and autoinflammatory diseases including in up to 27% of patients with SLE.<sup>1–3</sup> A remarkable finding reported during the SARS-CoV-2 pandemic was that pre-existing autoantibodies directed to T1IFN were associated with a higher risk of developing severe COVID-19.<sup>4–6</sup> Given the importance of IFN in host viral defences,<sup>7</sup> these observations raised the question of whether anti-T1IFN autoantibodies are associated with an increased risk of severe COVID-19 in SLE, a disease in which T1IFN plays a key role.<sup>8</sup>

## METHODS

We examined the prevalence of cytokine autoantibodies in prepandemic and intrapandemic sera from a SLE cohort using a commercially available 15-PLEX anticytokine array (Luminex Corporation, Austin, Texas, USA), which included detection of

## Key messages

### What is already known about this subject?

- Recent evidence indicated that the presence of neutralising anti-type 1 interferon (T1INF) antibodies is associated with severe COVID-19 outcome. Previous studies reported antibodies to a wide range of cytokines in SLE, including T1INF.
- We sought to determine the prevalence of cytokine antibodies including anti-T1INF in patients with SLE, exposed versus unexposed to COVID-19.

### What does this study add?

- In a cohort of patients with SLE examining prepandemic and intrapandemic antibodies to cytokines, none of the COVID-19 exposed patients (although all had a mild infectious course) had antibodies to T1INF compared with 3.7% of those without COVID-19 exposure (not statistically significant).
- Autoantibodies to other interferons and cytokines were not associated with exposure to SARS-CoV-2.

### How might this impact on clinical practice?

- Measurement of antibodies to T1INF in SLE may be an important biomarker identifying SLE at risk of severe COVID-19. A larger study including patients with SLE with a more severe COVID-19 course and the impact of neutralising autoantibodies and vaccines are needed.

IgG autoantibodies directed to IFN-gamma ( $\gamma$ ), IFN-beta ( $\beta$ ), B cell activating factor (BAFF)/B lymphocyte stimulator (Blys), tumour necrosis factor alpha ( $\alpha$ ), granulocyte colony stimulating factor, interleukin (IL)-1 $\alpha$ , IL-6, IL-8, IL-10, IL-12 (p40), IL-15, IL-17A, IL-17F, IL-18 and IL-22. In addition, autoantibodies to IFN- $\alpha$  (IFN- $\alpha$ 2b) were detected by an addressable laser bead immunoassay (ALBIA) using the full-length purified human protein (Abcam, Cambridge, UK) following protocols as previously published.<sup>9</sup> A cut-off of 500 median fluorescence intensity was established at two standard deviations above the mean of age-matched healthy controls,

**Table 1** Frequency of cytokine autoantibodies among prepandemic and intrapandemic samples stratified by SARS-CoV-2 exposure (positive RT-PCR test or positive SARS-CoV-2 serology)

	Prepandemic			Intrapandemic		
	Exposed patients with SLE* n=9 (%)	Unexposed patients with SLE† n=164 (%)	P value‡	Exposed patients with SLE* n=9 (%)	Unexposed patients with SLE† n=164 (%)	P value‡
IFN $\gamma$	0 (0)	17 (10.4)	0.658	1 (11.1)	8 (4.9)	0.961
IFN- $\alpha$ 2b	0 (0)	6 (3.7)	1.000	0 (0)	6 (3.7)	1.000
IFN $\beta$	0 (0)	0 (0)	NA	0 (0)	0 (0)	NA
BAFF	0 (0)	8 (4.9)	1.000	1 (11.1)	10 (6.1)	1.000
TNF- $\alpha$	0 (0)	3 (1.8)	1.000	0 (0)	1 (0.6)	1.000
G-CSF	0 (0)	4 (2.4)	1.000	1 (11.1)	1 (0.6)	0.205
IL-1 $\alpha$	0 (0)	0 (0)	NA	0 (0)	0 (0)	NA
IL-6	2 (22.2)	13 (7.9)	0.381	1 (11.1)	8 (4.9)	0.961
IL-8	0 (0)	1 (0.6)	1.000	0 (0)	1 (0.6)	1.000
IL-10	0 (0)	10 (6.1)	0.976	1 (11.1)	9 (5.5)	1.000
IL-12	0 (0)	4 (2.4)	1.000	0 (0)	3 (1.8)	1.000
IL-15	0 (0)	0 (0)	NA	0 (0)	1 (0.6)	1.000
IL-17A	1 (11.1)	6 (3.7)	0.813	1 (11.1)	2 (1.2)	0.367
IL-17F	0 (0)	5 (3.0)	1.000	0 (0)	2 (1.2)	1.000
IL-18	0 (0)	0 (0)	NA	0 (0)	0 (0)	NA
IL-22	1 (11.1)	7 (4.3)	0.891	1 (11.1)	7 (4.3)	0.891
Any	3 (33.3)	46 (28.0)	1.000	2 (22.2)	38 (23.1)	1.000

\*Exposure to SARS-CoV-2 defined as positive RT-PCR test and/or positive serology (n=9).

†Not exposed to SARS-CoV-2 defined as negative RT-PCR test, negative serology or not assessed (n=164).

‡The p values for the respective  $\chi^2$  comparisons. A value of  $p < 0.003$  is considered statistically significant based on the Bonferroni correction. BAFF, B cell activating factor; G-CSF, granulocyte colony stimulating factor; IFN, interferon; IL, interleukin; NA, not applicable; RT-PCR, reverse transcriptase PCR assay for SARS-CoV-2 viral RNA; TNF, tumour necrosis factor.

following the requirements for validation of laboratory tests in the accredited laboratory (MitogenDx) that performed these tests. SLE-related autoantibodies (anti-double stranded DNA (dsDNA), anti-Smith (Sm), anti-U1-ribonucleoprotein (RNP), anti-ribosomal P, anti-SSA/Ro60, anti-Ro52/ (tripartite motif containing-21) TRIM21 and antichromatin) were detected in the prepandemic and intrapandemic sera by commercially available multianalyte immunoassays as previously published.<sup>9 10</sup>

Prepandemic serum samples from the SLE cohort were biobanked prior to 01 January 2020 and intrapandemic samples were collected from 15 March 2020 to 31 January 2021; all patients were unvaccinated at the time of serum collection.<sup>11</sup> All samples were tested for SARS-CoV-2 antibodies using an ELISA measuring IgA and IgG antispike 1 (S1) protein (Euroimmun AG, Lübeck, Germany) and an assay detecting IgG antibodies to nucleocapsid (N), S1 receptor binding domain (RBD) and S1 (XMAP: Luminex Corporation, Austin, Texas, USA).<sup>11</sup> RT-PCR tests were performed if clinically indicated and results collected retrospectively through to 31 January 2021.

Cytokine autoantibodies were compared between patients with SLE with SARS-CoV-2-positive serology and/or RT-PCR confirmed SARS-CoV-2 infection (SARS-CoV-2 serology methods previously published)<sup>10</sup> vs the remaining cohort using a  $\chi^2$  test. The Bonferroni method

was used to correct for multiple comparisons (17), with a corrected  $\alpha$  of 0.0029 being considered statistically significant. Potential associations between each cytokine autoantibody and each SLE-related autoantibody in the prepandemic and intrapandemic periods were assessed using a  $\chi^2$  test, with a Bonferroni corrected  $\alpha$  of 0.00045 being considered significant to account for the 112 comparisons.

## PATIENT AND PUBLIC INVOLVEMENT

Patients or the public were not involved in the design, conduct, reporting or dissemination plans of our research.


## RESULTS

The SLE cohort consisted of 173 patients (94.8% female, mean age 48.5 years, mean disease duration 11.7 years, 42.8% non-White race/ethnicity, 83.2% prescribed hydroxychloroquine, 28.9% corticosteroids and 43.9% other immunomodulators).<sup>11</sup> Intrapandemic samples from nine patients had SARS-CoV-2-positive serology and/or RT-PCR confirmed SARS-CoV-2 infection (SARS-CoV-2 serology + and RT-PCR+ (n=3); SARS-CoV-2 serology + and RT-PCR not done (n=3); SARS-CoV-2 serology- and RT-PCR+ (n=3)), yielding 164 patients with no laboratory evidence of SARS-CoV-2 exposure. The frequency of

**Table 2** Characteristics of SLE cohort patients with positive intrapandemic SARS-CoV-2 serology and/or positive RT-PCR

ID	Date of PCR test	Date serum collected	*Age, years	Sex	Race/ethnicity	Medications			Cytokine antibodies	
						At time of pre-pandemic serum sample	At time of Intrapandemic serum sample	Pre-pandemic sample	Intrapandemic sample	
<b>SARS-CoV-2 antibody + and RT-PCR+</b>										
SLE-A	Oct '20	Oct '20	40–45	M	Asian	Hydroxychloroquine	Hydroxychloroquine	Hydroxychloroquine	NRR	NRR
SLE-B	Dec '20	Jan '21	40–45	F	Asian	Hydroxychloroquine, mycophenolate mofetil, corticosteroids	Hydroxychloroquine, mycophenolate mofetil, belimumab	Hydroxychloroquine, mycophenolate	IL-6, IL-17a	BAFF
SLE-C	Apr '20	Oct '20	45–50	F	Asian	Hydroxychloroquine, mycophenolate mofetil, corticosteroids	Hydroxychloroquine, mycophenolate mofetil	Hydroxychloroquine, mycophenolate	NRR	NRR
<b>SARS-CoV-2 antibody + and RT-PCR not done</b>										
SLE-D	ND	Oct '20	60–65	F	White	None	None	None	IL-6	IFN $\gamma$ , G-CSF, IL-10, IL-6, IL-22, IL-17a
SLE-E	ND	Oct '20	45–50	F	White	Hydroxychloroquine, methotrexate	Hydroxychloroquine, methotrexate	Hydroxychloroquine, methotrexate	NRR	NRR
SLE-F	ND	Oct '20	70–75	F	White	Hydroxychloroquine	Hydroxychloroquine	Hydroxychloroquine	NRR	NRR
<b>SARS-CoV-2 antibody – and RT-PCR+</b>										
SLE-G	Dec '20	Nov '20	20–25	F	White	Hydroxychloroquine, azathioprine	Hydroxychloroquine, azathioprine	Hydroxychloroquine, mycophenolate mofetil	NRR	NRR
SLE-H	Oct '20	Jun '20	50–55	F	Asian	Mycophenolate mofetil, corticosteroids	Methotrexate	Methotrexate	NRR	NRR
SLE-I	Nov '20	Sep '20	45–50	F	White	Hydroxychloroquine	Hydroxychloroquine	Hydroxychloroquine	IL-22	NRR

\*Age shown as a 5-year interval to protect confidentiality of individual.  
 BAFF, B cell activating factor; G-CSF, granulocyte colony stimulating factor; IFN, interferon; IL, interleukin; ND, not done; NRR, normal reference range; RT-PCR, reverse transcriptase PCR assay for SARS-CoV-2 viral RNA.



**Figure 1** Heat map of cytokine autoantibody titres in pre-pandemic and intrapandemic patients with SLE who had evidence of SARS-CoV-2 exposure versus those who had no evidence of SARS-CoV-2 exposure. Apparently healthy controls are shown for comparison as a reference for established cut-offs. Each vertical column in the heat map represents an individual patient. The patients are sorted from left to right based on descending type I interferon (T1IFN) autoantibody titres. Therefore, the order of the patients in the pre-pandemic and intrapandemic panels are not the same. A supplemental figure where the pre-pandemic and intrapandemic results are vertically aligned is provided online supplemental figure 1. MFI, median fluorescence intensity.

cytokine autoantibodies in the pre-pandemic and intrapandemic samples stratified by SARS-CoV-2 exposure is shown in [table 1](#), the patient characteristics and cytokine autoantibodies for the nine exposed patients are shown in [table 2](#), and a graphical representation of the distribution of cytokine autoantibodies and their titers is shown in [figure 1](#) (patients are sorted from left to right based on descending T1IFN autoantibody titres separately for pre-pandemic and intrapandemic samples).

In the pre-pandemic samples, 33.3% (3/9) of patients with serological and/or RT-PCR evidence of SARS-CoV-2 exposure versus 28.0% (46/164) without exposure had at least one cytokine antibody ( $p=1.00$ ). In the intrapandemic samples, 22.2% (2/9) of exposed versus 23.1% (38/164) without exposure had at least one cytokine antibody ( $p=1.00$ ). Overall, there was no difference in the frequency of cytokine autoantibodies in pre-pandemic compared with intrapandemic samples (28.3% (49/173) vs 23.1% (40/173), respectively;  $p=0.33$ ). In the entire cohort of 173 patients, the most common cytokine autoantibodies in pre-pandemic samples were directed against IFN- $\gamma$  (9.8%), followed by IL-6 (8.7%), IL-10 (5.8%) and BAFF (4.6%) ([table 1](#) and [figure 1](#)) compared with 5.2% for IFN- $\gamma$  and anti-IL-6, 5.8% for IL-10% and 6.4% for BAFF in intrapandemic samples. In the pre-pandemic samples, none of the exposed patients had autoantibodies to IFN- $\gamma$  versus 10.4% (17/164) without exposure ( $p=0.66$ ). In the intrapandemic samples, 11.1% (1/9) of the exposed patients had autoantibodies to IFN- $\gamma$  versus 4.9% (8/164) without exposure ( $p=0.96$ ). In the pre-pandemic and intrapandemic samples, none of the exposed patients had autoantibodies to T1IFN (IFN- $\alpha$  or IFN- $\beta$ ) versus 3.7% (6/164) without exposure had autoantibodies

to IFN- $\alpha$  and 0% without exposure had autoantibodies to IFN- $\beta$  ( $p=1.00$  and not applicable, respectively).

When cytokine autoantibodies were analysed for associations with SLE-related autoantibodies, only in intrapandemic samples were there associations with  $p$  values less than 0.05: anti-IFN- $\alpha$  and anti-dsDNA (intrapandemic  $p=0.015$  vs pre-pandemic  $p=0.7035$ ); IL-12 and anti-dsDNA (intrapandemic  $p=0.043$  vs pre-pandemic  $p=0.816$ ); IL-12 and anti-ribosomal P (intrapandemic  $p=0.009$  vs pre-pandemic  $p=1.000$ ); and IL-12 and anti-Sm (intrapandemic  $p=0.005$  vs pre-pandemic  $p=1.000$ ). However, after correcting for multiple comparisons none of these associations met the reduced  $\alpha$  ( $\alpha=0.00045$ ) to be considered statistically significant.

## DISCUSSION

The frequency of cytokine autoantibodies observed in our study is in keeping with previous publications that used similar assay techniques reporting that up to 27% of patients with SLE had cytokine autoantibodies.<sup>1,3</sup> In the present study, none of the patients with SLE with confirmed exposure to SARS-CoV-2 had autoantibodies to T1IFN and, as we previously reported,<sup>11</sup> none of the 173 patients in our cohort, including the 9 patients with confirmed exposure, had severe COVID-19 symptoms necessitating hospitalisation. Hence, we could not confirm that T1IFN autoantibodies increase the risk for severe SARS-CoV-2. The overall observed frequency of anti-IFN $\gamma$  autoantibodies in our study is consistent with a previous report of 11% in patients with SLE.<sup>1</sup> In an earlier communication, we reported that this SLE cohort had a lower rate of pre-pandemic seropositivity and a slightly



lower to similar rate of intrapandemic seropositivity compared with contemporaneous controls.<sup>11</sup> Granted, in the current report we did not study the neutralising effect of cytokine autoantibodies, but it would be difficult to explain T1IFN neutralising activity in the absence of detectable T1IFN cytokine autoantibodies. In addition, we found a non-significant association between IFN $\gamma$  autoantibodies and dsDNA autoantibodies in intrapandemic samples, which is similar to a previous report that anti-IFN $\gamma$  autoantibodies were associated with anti-dsDNA.<sup>1</sup> Further, due to the size of our cohort and its geographical context, our results may not be generalisable to a broader population of patients with SLE. Because of interest in vaccine responses in patients with SLE,<sup>12–15</sup> current efforts are focusing on a possible relationship of cytokine autoantibodies to adverse or unusual reactions.

**Acknowledgements** The authors thank Haiyan Hou, Meifeng Zhang and Dr Paul Sciore (University of Calgary) for their laboratory technical assistance.

**Contributors** All authors were involved in the concept and design, data analysis and interpretation, and editing for intellectual content. MJF and AEC are joint senior authors.

**Funding** AEC holds The Arthritis Society Chair in Rheumatic Diseases at the University of Calgary and some of these funds were used to support HW and FSC.

**Competing interests** MJF is a consultant to and has received honoraria and/or travel support from Werfen (Barcelona, Spain, San Diego, CA). MJF is also Medical Director of Mitogen Diagnostics Corporation. MYC has received consulting fees from Janssen, AstraZeneca and MitogenDx. AEC has received consulting fees from AstraZeneca, Bristol Myers, Squibb and Glaxo Smith Kline. All other authors declare no conflict of interest.

**Patient consent for publication** Consent obtained directly from patient(s).

**Ethics approval** This study involves human participants and was approved by the University of Calgary Ethics Board (REB13-0425). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is

properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

May Y Choi <http://orcid.org/0000-0003-3760-2737>

#### REFERENCES

- Gupta S, Tatouli IP, Rosen LB, *et al*. Distinct functions of autoantibodies against interferon in systemic lupus erythematosus: a comprehensive analysis of anticytokine autoantibodies in common rheumatic diseases. *Arthritis Rheumatol* 2016;68:1677–87.
- Zhang Q, Bastard P, Liu Z, *et al*. Inborn errors of type I IFN immunity in patients with life-threatening COVID-19. *Science* 2020;370.
- Slavikova M, Schmeisser H, Kontsejkova E, *et al*. Incidence of autoantibodies against type I and type II interferons in a cohort of systemic lupus erythematosus patients in Slovakia. *J Interferon Cytokine Res* 2003;23:143–7.
- Bastard P, Rosen LB, Zhang Q, *et al*. Autoantibodies against type I IFNs in patients with life-threatening COVID-19. *Science* 2020;370:eabd4585.
- Damoiseau J, Dotan A, Fritzler MJ, *et al*. Autoantibodies and SARS-CoV2 infection: the spectrum from association to clinical implication: report of the 15th Dresden Symposium on autoantibodies. *Autoimmun Rev* 2022;21:103012.
- Knight JS, Caricchio R, Casanova J-L, *et al*. The intersection of COVID-19 and autoimmunity. *J Clin Invest* 2021;131:e154886.
- Hoffmann H-H, Schneider WM, Rice CM. Interferons and viruses: an evolutionary arms race of molecular interactions. *Trends Immunol* 2015;36:124–38.
- Crow MK, Ronnblom L. Type I interferons in host defence and inflammatory diseases. *Lupus Sci Med* 2019;6:e000336–e36.
- Trahtenberg U, Fritzler MJ, On behalf of the COVID-19 chapter of the “Longitudinal Biomarkers in Lung Injury” study group. COVID-19-associated autoimmunity as a feature of acute respiratory failure. *Intensive Care Med* 2021;47:801–4.
- Choi MY, Clarke AE, St. Pierre Y, *et al*. Antinuclear Antibody–Negative systemic lupus erythematosus in an international inception cohort. *Arthritis Care Res* 2019;71:893–902.
- Mathew HR, Choi MY, Buhler K, *et al*. SARS-CoV-2 seroprevalence, seroconversion and neutralizing antibodies in a systemic lupus erythematosus cohort and comparison to controls. *Lupus* 2021;30:2318–20.
- Sattui SE, Liew JW, Kennedy K, *et al*. Early experience of COVID-19 vaccination in adults with systemic rheumatic diseases: results from the COVID-19 global rheumatology alliance vaccine survey. *RMD Open* 2021;7:e001814–e14.
- Connolly CM, Boyarsky BJ, Ruddy JA, *et al*. Absence of humoral response after two-dose SARS-CoV-2 messenger RNA vaccination in patients with rheumatic and musculoskeletal diseases: a case series. *Ann Intern Med* 2021;174:1332–4.
- Tani C, Pratesi F, Talarico R, *et al*. Efficacy of anti-SARS-CoV-2 mRNA vaccine in systemic autoimmune disorders: induction of high avidity and neutralising anti-RBD antibodies. *RMD Open* 2021;7:e001914–e14.
- Kroon FPB, Najm A, Alunno A, *et al*. Risk and prognosis of SARS-CoV-2 infection and vaccination against SARS-CoV-2 in rheumatic and musculoskeletal diseases: a systematic literature review to inform EULAR recommendations. *Ann Rheum Dis* 2022;81:422–432.