

Abstract PO.7.160 Table 1 Enlight-LN registry inclusion and exclusion criteria

Inclusion Criteria	Exclusion Criteria
<ul style="list-style-type: none"> Adults ≥18 years of age Biopsy-confirmed lupus nephritis Initiating or have initiated treatment with voclosporin within the 3 months prior to consent Ability to understand and provide written consent 	<ul style="list-style-type: none"> Off-label use (use of voclosporin outside of the FDA-approved labeling)

FDA, food and drug administration

utilization. The registry will enroll patients who are initiating or who have already initiated treatment with commercial voclosporin within 3 months prior to consent. Patients ≥ 18 years of age with biopsy-confirmed lupus nephritis are eligible (Table 1). Secondary objectives include describing at baseline and during the study period the clinical characteristics, treatment and response patterns of patients treated with voclosporin.

To date, 36 sites in 16 states have been selected to participate in the registry; Enlight-LN is currently enrolling patients.

PO.7.161 CLINICAL AND IMMUNOLOGICAL CHARACTERIZATION OF SYSTEMIC LUPUS ERYTHEMATOSUS (SLE) PATIENTS IN ESTONIA. A PROSPECTIVE COHORT STUDY OF 40 SLE PATIENTS IN ESTONIA

¹S Meisalu*, ²K Kisand. ¹East-Tallinn Central Hospital ~ Tallinn ~ Estonia; ²University of Tartu ~ Estonia

10.1136/lupus-2022-elm2022.181

Background SLE is a rare chronic autoimmune disease with polymorphic clinical manifestation and wide-ranging disease course with treatment tactics dependent on disease activity and organ involvement. In 2017, a study to estimate prevalence and incidence of SLE in Estonia was done, but there is no data published to describe the Estonian SLE population. The aim of the present study is to analyze a sample of Estonian SLE patients.

Methods Consecutive outpatient and inpatient patients with rheumatologist diagnosed SLE (≥ 20 years) were enrolled in East-Tallinn Central Hospital. Two study visits were done with 6 months apart to evaluate disease activity, current treatment, organ involvement, immunological findings and comorbidities. In addition, data from medical records were collected: organ involvement and immunological findings at the time of diagnosis and initial treatment. SLE disease activity was measured using SLEDAI 2K (Systemic Lupus Erythematosus Disease Activity Index 2K) score.

Results Among 40 patients (mean age 50 (standard deviation ± 12.4) years, mean disease duration 12 (± 9.9) years, mean SLEDAI 2K at diagnosis 10 (± 3.9)) 92.5% were females. Mean SLEDAI 2K value at entering into the study was 4 (± 3.4) similar to the value after six months 4 (± 4.9). 82.5% of patients received hydroxychloroquine and 75% glycocorticosteroid treatment, 27.5% of patients were treated with rituximab. During their disease course 90% had joint and 50% skin involvement, 35% had leucopenia, all patients were positive for antinuclear antibody (ANA), 80% were anti-double-stranded DNA antibody (anti-dsDNA) positive and 70% of patients had low complement levels.

Conclusion The first analysis of Estonian SLE patients' clinical and laboratory parameters indicates that the disease is overall well managed in most of the patients. Further studies are in

progress on collected serum and PBMC samples to find immunological causes for poor treatment response.

PO.7.162 EPIDEMIOLOGY OF SYSTEMIC LUPUS ERYTHEMATOSUS IN CENTRAL SWEDEN: A POPULATION-BASED COHORT STUDY FROM THE ÖSTERGÖTLAND COUNTY OVER 14 YEARS

¹EV Arkema, ²M Saleh, ³JF Simard, ²C Sjöwall*. ¹Department of Medicine Solna, Division of Clinical Epidemiology, Karolinska Institutet ~ Stockholm ~ Sweden; ²Department of Biomedical and Clinical Sciences, Division of Inflammation and Infection/Rheumatology, Linköping University ~ Linköping ~ Sweden; ³Department of Medicine, Division of Immunology and Rheumatology, Stanford School of Medicine, Stanford ~ Palo Alto, California ~ USA


10.1136/lupus-2022-elm2022.182

Purpose We examined variations in incidence and prevalence of systemic lupus erythematosus (SLE) within a geographically defined area of central Sweden over a time period of 14 years. We described longitudinal differences in disease activity measures (e.g., the SLE disease activity index-2000 [SLEDAI-2K] and the Physician's Global Assessment), laboratory measurements and disease manifestations included among the American College of Rheumatology (ACR) criteria.

Methods We identified adults (≥ 18 years) residing in Östergötland County between 2008 and 2021 (mean adult population: 357 000 citizens) with a clinical diagnosis of SLE. Cases were defined as those with an SLE diagnosis set by a rheumatologist combined with fulfillment of the 1982 ACR classification criteria and/or the Fries' diagnostic principle (presence of antinuclear antibodies [ANA] by immunofluorescence microscopy at least once plus involvement of at least two defined organ systems). All subjects were included in the quality and research register 'Clinical Lupus Register in North-Eastern Gothia' (Swedish acronym KLURING). Individuals were followed prospectively until death, December 31, 2021, or emigration. We estimated incidence per 100 000 inhabitants stratified by sex and age. We used linear regression with calendar year of diagnosis as the outcome to assess whether each clinical measurement at diagnosis varied over time.

Results 126 new SLE cases (80% females) were diagnosed during the period 2008–2021, yielding a mean annual incidence of 3.0 per 100 000 inhabitants; higher in females (4.8 per 100 000) than in males (1.2 per 100 000). The mean age at diagnosis was 43.7 (Standard deviation [SD]

Incidence of SLE per 100 000 per year by age and sex in Östergötland County 2008–2021

**Abstract PO.7.162 Figure 1**