

203

INSIGHTS INTO LUPUS BIOLOGY FROM INBORN ERRORS OF IMMUNITY: IMMUNOPATHOGENESIS OF STAT1 GAIN-OF-FUNCTION AUTOIMMUNITY

¹Largent Andrea, ²Kathi Lambert, ¹Chiang Kristy, ¹Shumlak Natali, ³Quan-Zhen Li, ⁴Kelly Hudkins, ⁵Denny Liggett, ⁶Mohammed Oukka, ⁷Troy Torgerson, ²Jane Buckner, ^{1,6}Eric Allenspach, ^{1,6,8}David J Rawlings, ^{1,4,6}Shaun W Jackson. ¹Seattle Children's Research Institute, Seattle, WA; ²Benaroya Research Institute, Seattle, WA; ³Department of Immunology, University of Texas Southwestern Medical Center, Dallas, TX; ⁷Laboratory Medicine and Pathology; ⁸Comparative Medicine, University of Washington School of Medicine; ⁵Departments of Pediatrics; ⁴Allen Institute for Immunology, Seattle, WA 98109, USA; ⁶Immunology

10.1136/lupus-2022-lupus21century.7

Background Human genetic variations resulting in immunodeficiency and/or the propensity for autoimmunity represent “experiments of nature” that can advance our understanding of polygenic human diseases. Heterozygous gain-of-function (GOF) mutations in *STAT1* induce immune dysregulation characterized by autosomal dominant chronic mucocutaneous candidiasis (CMC) and the propensity for humoral autoimmunity. However, given widespread expression of *STAT1* in immune and non-immune lineages and engagement by multiple cytokine receptors, the immune mechanisms driving breaks in immune tolerance in *STAT1* GOF syndrome remain poorly understood. In addition, *STAT1* GOF variants are thought to enhance cytokine signaling by increasing total *STAT1* protein levels, but the cause of this phenotype has not been identified.

Methods To gain insights into the complex roles for *STAT1* in human immunity, we performed mass cytometry using PBMCs from *STAT1* GOF patients. Affected subjects were studied prior to treatment with JAK inhibitors, allowing unique insight into the immune landscape of *STAT1*-driven immune dysregulation. In parallel, we generated a novel murine knock-in strain allowing cell-intrinsic expression of a pathogenic human *STAT1* GOF mutation

Results We performed multiparameter immunophenotyping of pediatric *STAT1* GOF patients and age-matched controls to identify immune characteristics of *STAT1*-driven inflammation. Affected patients exhibited expansion of CXCR3-expressing CD4⁺ T and B cell populations exhibiting surface markers indicative of B cell helper function and extrafollicular activation, respectively. Moreover, relative expansion of these adaptive immune populations correlated with serum autoantibody titers. To study underlying mechanisms, we generated a new *Stat1* GOF transgenic model and confirmed the development of spontaneous humoral autoimmunity recapitulating the human phenotype. Despite clinical resemblance to human regulatory T cell (Treg) deficiency (IPEX syndrome), *Stat1*^{GOF} mice and humans exhibited normal Treg development and function. Rather, *STAT1* GOF autoimmunity was driven by dysregulated *STAT1*-dependent signals downstream of the type 1 and type 2 interferon (IFN) receptors. Surprisingly, autoimmunity in *Stat1*^{GOF} mice lacking the type 1 IFN receptor (IFNAR) was only partially ameliorated, whereas loss of type 2 IFN (IFN- γ) signals prevented disease. Strikingly, IFN- γ R deletion abolished the known increase in total *STAT1* expression resulting in normalization of *STAT1*-dependent systemic inflammation.

Conclusions Since *STAT1* regulates its own transcription, these findings highlight IFN- γ as the critical driver of a feedforward inflammatory cascade in *STAT1* GOF syndrome. More broadly, our data provide new insights into the *STAT1*-

dependent cellular mechanisms underlying both rare monogenic and more common polygenic autoimmune diseases, such as systemic lupus erythematosus.

204

IL-4 AS A NEGATIVE REGULATOR OF PATHOGENIC EXTRAFOLLICULAR DN2 B CELLS IN SLE

John D Mountz, Changming Lu, Shanrun Liu, Min Gao, Winn Chatham, Hui-Chen Hsu. University of Alabama at Birmingham

10.1136/lupus-2022-lupus21century.8

Background Pathogenic extrafollicular double negative 2 (DN2) B cells in SLE have a phenotype of IgD⁻CD27⁻Tbet⁺CD11c⁺, and are the precursor of ribonuclear protein (RNP) autoantibody producing B cells. This trajectory of development is promoted by a synergistic effect of type I interferon (IFN) and TLR7-induced activation of B cells at the transitional (Tr) and naïve (NAV) developmental stages. In contrast, IL-4 and IL-4R expression is low in SLE patients and higher expression of IL-4 has been associated with a milder disease course in SLE. The present studies were carried out to determine the mechanism associated with IL-4-mediated B-cell quiescence program *in vitro* in SLE B cells and *in vivo* in the BXD2 mouse model of SLE.

Methods All SLE patients met the American College of Rheumatology 1997 revised criteria and the 2017 ACR/EULAR classification criteria for SLE. Peripheral blood mononuclear cells were analyzed by FACS for surface expression of IL-4R, intracellular IFN- β , and intranuclear T-bet and IRF7. DN2 B cell differentiation *in vitro* was stimulated with or without IL-4 50 ng/ml pre-culture for 1 hr. B-cell phenotypes, autoantibody profiles, and their association with the expression of IL-4R and IFN- β were analyzed in 47 SLE patients. The transcriptomics program associated with B-cell fate decision at the Tr, NAV, and activated naïve (aNAV) stages of B cell development in healthy control (HC) subjects and SLE subjects was analyzed using single cell RNA-sequencing (scRNA-seq) analysis. BXD2 mice were injected weekly with IL-4 complex or carrier anti-IL-4 antibody followed by R848, a TLR7 ligand. Four weeks later, mice were treated an additional IV injection of IL-4 complex or carrier. One week later, mice were sacrificed. The development of autoantibodies, GC B cells, DN2 B cells, and B-cell transcriptome were analyzed.

Results Section scRNA-seq analysis showed that type I interferon (IFN) stimulated genes (ISGs) were upregulated at Tr, rNAV, and aNAV stages of SLE B cells. In contrast, Tr and NAV B cells from SLE patients exhibited downregulation of an IL-4R quiescent gene program consisting of *IL4R*, *BACH2*, and *FCRE2A* (CD23). In HC, aNAV B cells exhibited upregulation of gene signatures of germinal center (GC) and classical memory (cMEM) B cells including *LTB*, *GPR183*, *CD27*, *CD44*, and *CD83*. IKAROS was identified as a top transcription factor associated with the upregulated genes in B cells from HC. In contrast, in SLE, aNAV B cells expressed signature genes of DN2 B cells including *FCRL3*, *FCRL5*, and *ZEB2*. Pseudotime analysis revealed in SLE, 63% of B cells developed into the DN2 pathway compared to only 3% in healthy controls. In contrast, only 14% of SLE B cells developed into the GC and classical memory pathway compared to 20% for healthy controls. IL-4 pre-treatment resulted in a significant increase of the CD11c⁻Tbet⁻ DN1 subpopulation and a decrease in the CD11c⁺Tbet⁺ DN2 B cells from SLE